


①  
S. ...  
...  
...  
...



*Handwritten text, possibly a signature or title, in cursive script.*



AN

INTRODUCTION

TO

**ALGEBRA,**

BEING THE

FIRST PART

OF A

**COURSE OF MATHEMATICS,**

ADAPTED TO

THE METHOD OF INSTRUCTION IN THE HIGHER SCHOOLS AND  
ACADEMIES IN THE UNITED STATES.

\*\*\*\*\*

By JEREMIAH DAY, LL.D.  
President of Yale College.



PUBLISHED BY HOWE & SPALDING.

\*\*\*\*\*

FLAGG & GRAY, PRINTERS.

1819.


QA153

D39

25443




THE following elementary compendium is the same as the Introduction to Algebra intended for the use of Colleges, with the omission of a few of the sections at the latter end of the book. These it was supposed would not generally be needed, in private schools and academies. The parts retained contain the *elements* of the science. The first two hundred pages are precisely the same as in the second edition of the larger work. The subjects in the first edition, which are omitted in this abridgment, are Variation, Progression, Infinity, the Binomial Theorem, Resolution of the higher Equations, Application of Algebra to Geometry, and Equations of Curves.



# CONTENTS.

	PAGE
Introductory Observations, on the Mathematics in general,	1
ALGEBRA.	
Section I. Notation, Positive and Negative quantities, Axioms, &c.	8
II. Addition,	21
III. Subtraction,	27
IV. Multiplication,	31
V. Division,	41
VI. Algebraic Fractions,	49
VII. Reduction of Equations, by Transposition, Multiplication, and Division. Solution of Problems,	65
VIII. Involution. Notation, addition, subtraction, multiplication, and division of Powers,	83
IX. Evolution. Notation, reduction, addition, subtraction, multiplication, division, and involution, of Radical Quantities,	99
X. Reduction of Equations by Involution and Evolution. Affected Quadratic Equations,	128
XI. Solution of Problems which contain two or more Unknown Quantities,	153
XII. Ratio and Proportion,	173
XIII. Division by Compound Divisors,	205
XIV. Evolution of Compound Quantities,	210
XV. Infinite Series,	214



# INTRODUCTORY OBSERVATIONS

ON THE

## Mathematics in General.

---

ART. 1. **M**ATHEMATICS is the science of QUANTITY.

Any thing which can be multiplied, divided, or measured, is called *quantity*. Thus, a *line* is a quantity, because it can be doubled, trebled, or halved; and can be measured, by applying to it another line, as a foot, a yard, or an ell. *Weight* is a quantity, which can be measured, in pounds, ounces, and grains. *Time* is a species of quantity, whose measure can be expressed, in hours, minutes, and seconds. But *colour* is not a quantity. It cannot be said, with propriety, that one colour is twice as great, or half as great as another. The operations of the *mind*, such as thought, choice, desire, hatred, &c. are not quantities. They are incapable of mensuration.\*

2. Those parts of the Mathematics, on which all the others are founded, are *Arithmetic*, *Algebra*, and *Geometry*.

3. ARITHMETIC is the science of *numbers*. Its aid is required, to complete and apply the calculations, in almost every other department of the mathematics.

4. ALGEBRA is a method of computing by *letters* and other symbols. FLUXIONS, or the Differential and Integral Calculus, may be considered as belonging to the higher branches of algebra.†

5. GEOMETRY is that part of the mathematics, which treats of *magnitude*. By magnitude, in the appropriate sense of

\* See note A.

† See note B.

the term, is meant that species of quantity, which is *extended*; that is, which has one or more of the three dimensions, *length*, *breadth*, and *thickness*. Thus a *line* is a magnitude, because it is extended, in length. A *surface* is a magnitude, having length and breadth. A *solid* is a magnitude, having length, breadth, and thickness. But *motion*, though a quantity, is not, strictly speaking, a magnitude. It has neither length, breadth, nor thickness.\*

6. TRIGONOMETRY and CONIC SECTIONS are branches of the mathematics, in which, the principles of geometry are applied to *triangles*, and the sections of a *cone*.

7. Mathematics are either pure, or mixed. In *pure* mathematics, quantities are considered, independently of any substances actually existing. But, in *mixed* mathematics, the relations of quantities are investigated, in connection with some of the properties of matter, or with reference to the common transactions of business. Thus, in *Surveying*, mathematical principles are applied to the measuring of land; in *Optics*, to the properties of light; and in *Astronomy*, to the motions of the heavenly bodies.

8. The science of the pure mathematics has long been distinguished, for the clearness and distinctness of its principles; and the irresistible conviction, which they carry to the mind of every one who is once made acquainted with them. This is to be ascribed, partly to the nature of the subjects, and partly to the exactness of the definitions, the axioms, and the demonstrations.

9. The foundation of all mathematical knowledge must be laid in definitions. A *definition* is an explanation of what is meant, by any word or phrase. Thus, an equilateral triangle is defined, by saying, that it is a figure bounded by three equal sides.

It is essential to a complete definition, that it perfectly distinguish the thing defined, from every thing else. On many subjects, it is difficult to give such precision to language, that it shall convey, to every hearer or reader, exactly the same ideas. But, in the mathematics, the principal terms may be so defined, as not to leave room for the least difference of apprehension, respecting their meaning. All must be agreed, as to the nature of a circle, a square, and a triangle, when they have once learned the definitions of these figures.

\* Some writers, however, use magnitude, as synonymous with quantity.

Under the head of definitions, may be included explanations of the *characters* which are used to denote the relations of quantities. Thus, the character  $\sqrt{\quad}$  is explained or defined, by saying that it signifies the same as the words square root.

10. The next step, after becoming acquainted with the meaning of mathematical terms, is to bring them together, in the form of propositions. Some of the relations of quantities require no process of reasoning, to render them evident. To be understood, they need only to be proposed. That a square is a different figure from a circle; that the whole of a thing is greater, than one of its parts; and, that two straight lines cannot inclose a space, are propositions so manifestly true, that no reasoning upon them could make them more certain. They are, therefore, called self-evident truths, or *axioms*.

11. There are, however, comparatively few mathematical truths which are self-evident. Most require to be proved, by a chain of reasoning. Propositions of this nature are denominated *theorems*; and the process, by which, they are shown to be true, is called *demonstration*. This is a mode of arguing, in which, every inference is immediately derived, either from definitions, or from principles which have been previously demonstrated. In this way, complete certainty is made to accompany every step, in a long course of reasoning.

12. Demonstration is either *direct*, or *indirect*. The former is the common, obvious mode of conducting a demonstrative argument. But, in some instances, it is necessary to resort to indirect demonstration; which is a method of establishing a proposition, by proving that to suppose it *not* true, would lead to an absurdity. This is frequently called *reductio ad absurdum*. Thus, in certain cases in geometry, two lines may be proved to be equal, by shewing that to suppose them unequal, would involve an absurdity.

13. Besides the principal theorems in the mathematics, there are also *Lemmas*, and *Corollaries*. A Lemma is a proposition which is demonstrated, for the purpose of using it, in the demonstration of some other proposition. This preparatory step is taken to prevent the proof of the principal theorem from becoming complicated and tedious.

14. A *Corollary* is an *inference* from a preceding proposition. A Scholium is a remark of any kind, suggested by something which has gone before, though not, like a corollary, immediately depending on it.

15. The immediate object of inquiry, in the mathematics, is, frequently, not the demonstration of a general truth, but a method of performing some operation, such as reducing a vulgar fraction to a decimal, extracting the cube root, or inscribing a circle in a square. This is called solving a problem. A *theorem* is something to be *proved*. A *problem* is something to be *done*.

16. When that which is required to be done, is so easy, as to be obvious to every one, without an explanation, it is called a *postulate*. Of this nature, is the drawing of a straight line, from one point to another.

17. A quantity is said to be *given*, when it is either supposed to be already *known*, or is made a *condition*, in the statement of any theorem or problem. In the rule of proportion in arithmetic, for instance, three terms must be given, to enable us to find a fourth. These three terms are the *data*, upon which the calculation is founded. If we are required to find the number of acres, in a circular island ten miles in circumference, the circular figure, and the length of the circumference, are the data. They are said to be given *by supposition*, that is, by the conditions of the problem. A quantity is also said to be given, when it may be directly and easily *inferred*, from something else which is given. Thus, if two numbers are given, their *sum* is given; because it is obtained, by merely adding the numbers together.

In Geometry, a quantity may be given, either in *position*, or *magnitude*, or both. A line is given in position, when its *situation* and *direction* are known. It is given in magnitude, when its *length* is known. A circle is given in *position*, when the place of its centre is known. It is given in *magnitude*, when the length of its diameter is known.

18. One proposition is *contrary*, or contradictory to another, when, what is affirmed, in the one, is denied, in the other. A proposition and its contrary, can never *both* be true. It cannot be true, that two given lines are equal, and that they are *not* equal, at the same time.

19. One proposition is the *converse* of another, when the order is inverted; so that, what is *given* or supposed, in the first, becomes the *conclusion*, in the last; and what is given in the last, is the conclusion, in the first. Thus, it can be proved, first, that if the *sides* of a triangle are equal, the *angles* are equal; and secondly, that if the *angles* are equal, the *sides* are equal. Here, in the first proposition, the equal-

ity of the *sides is given* ; and the equality of the *angles inferred* : in the second, the equality of the *angles is given*, and the equality of the *sides inferred*. In many instances, a proposition and its converse are both true ; as in the preceding example. But this is not always the case. A circle is a figure bounded by a curve ; but a figure bounded by a curve is not of course a circle.

20. The practical applications of the mathematics, in the common concerns of business, in the useful arts, and in the various branches of physical science, are almost innumerable. Mathematical principles are necessary, in *mercantile transactions*, for keeping, arranging, and settling accounts, adjusting the prices of commodities, and calculating the profits of trade : in *Navigation*, for directing the course of a ship on the ocean, adapting the position of her sails to the direction of the wind, finding her latitude and longitude, and determining the bearings and distances of objects on shore : in *Surveying*, for measuring, dividing, and laying out grounds, taking the elevation of hills, and fixing the boundaries of fields, estates, and public territories : in *Mechanics*, for understanding the laws of motion, the composition of forces, the equilibrium of the mechanical powers, and the structure of machines : in *Architecture*, for calculating the comparative strength of timbers, the pressure which each will be required to sustain, the forms of arches, the proportions of columns, &c. : in *Fortification*, for adjusting the position, lines, and angles, of the several parts of the works : in *Gunnery*, for regulating the elevation of the cannon, the force of the powder, and the velocity and range of the shot : in *Optics*, for tracing the direction of the rays of light, understanding the formation of images, the laws of vision, the separation of colours, the nature of the rainbow, and the construction of microscopes and telescopes : in *Astronomy*, for computing the distances, magnitudes, and revolutions of the heavenly bodies ; and the influence of the law of gravitation, in raising the tides, disturbing the motions of the moon, causing the return of the comets, and retaining the planets in their orbits : in *Geography*, for determining the figure and dimensions of the earth, the extent of oceans, islands, continents, and countries ; the latitude and longitude of places, the courses of rivers, the height of mountains, and the boundaries of kingdoms : in *History*, for fixing the chronology of remarkable events, and estimating the strength of armies, the wealth of nations, the value of their revenues, and the amount of their population : and, in the concerns of

*Government*, for apportioning taxes, arranging schemes of finance, and regulating national expenses. The mathematics have also important applications to Chemistry, Mineralogy, Music, Painting, Sculpture, and indeed to a great proportion of the whole circle of arts and sciences.

21. It is true, that, in many of the branches which have been mentioned, the ordinary business is frequently transacted, and the mechanical operations performed, by persons who have not been regularly instructed in a course of mathematics. Machines are framed, lands are surveyed, and ships are steered, by men who have never thoroughly investigated the principles, which lie at the foundation of their respective arts. The reason of this is, that the methods of proceeding, in their several occupations, have been pointed out to them, by the genius and labour of others. The mechanic often works by rules, which men of science have provided for his use, and of which he knows nothing more, than the practical application. The mariner calculates his longitude by tables, for which he is indebted to mathematicians and astronomers of no ordinary attainments. In this manner, even the abstruse parts of the mathematics are made to contribute their aid to the common arts of life.

22. But an additional and more important advantage, to persons of a liberal education, is to be found, in the enlargement and improvement of the reasoning powers. The mind, like the body, acquires strength by exertion. The art of reasoning, like other arts, is learned by practice. It is perfected, only by long continued exercise. Mathematical studies are peculiarly fitted for this discipline of the mind. They are calculated to form it to habits of fixed attention; of sagacity, in detecting sophistry; of caution, in the admission of proof; of dexterity, in the arrangement of arguments; and of skill, in making all the parts of a long continued process tend to a result, in which the truth is clearly and firmly established. When a habit of close and accurate thinking is thus acquired; it may be applied to any subject, on which a man of letters or of business may be called to employ his talents. "The youth," says Plato, "who are furnished with mathematical knowledge, are prompt and quick, at all other sciences."

It is not pretended, that an attention to other objects of inquiry, is rendered unnecessary, by the study of the mathematics. It is not their office, to lay before us historical facts; to teach the principles of morals; to store the fancy with



brilliant images; or to enable us to speak and write with rhetorical vigour and elegance. The beneficial effects which they produce on the mind, are to be seen, principally, in the regulation and increased energy of the *reasoning powers*. These they are calculated to call into frequent and vigorous exercise. At the same time, mathematical studies may be so conducted, as not often to require excessive exertion and fatigue. Beginning with the more simple subjects, and ascending gradually to those which are more complicated; the mind acquires strength, as it advances; and by a succession of steps, rising regularly one above another, is enabled to surmount the obstacles which lie in its way. In a course of mathematics, the parts succeed each other in such a connected series, that the preceding propositions are preparatory to those which follow. The student who has made himself master of the former, is qualified for a successful investigation of the latter. But he who has passed over any of the ground superficially, will find that the obstructions to his future progress are yet to be removed. In mathematics, as in war, it should be made a principle, not to advance, while any thing is left unconquered behind. It is important that the student should be deeply impressed with a conviction of the necessity of this. Neither is it sufficient that he understands the *nature* of one proposition or method of operation, before proceeding to another. He ought also to make himself *familiar* with every step, by a careful attention to the examples. He must not expect to become thoroughly versed in the science, by merely *reading* the main principles, rules and observations. It is practice only, which can put these completely in his possession. The method of studying here recommended, is not only that which promises success, but that which will be found, in the end, to be the most expeditious, and by far the most pleasant. While a superficial attention occasions perplexity and consequent aversion; a thorough investigation is rewarded with a high degree of gratification. The peculiar entertainment which mathematical studies are calculated to furnish to the mind, is reserved for those who make themselves masters of the subjects to which their attention is called.

NOTE. The principal definitions, theorems, rules, &c. which it is necessary to *commit to memory*, are distinguished by being put in Italics or Capitals.

{ Henry or HENRY

# ALGEBRA.

---

## SECTION I.

### NOTATION, NEGATIVE QUANTITIES, AXIOMS, &c.

ART. 23. **ALGEBRA** may be defined, A GENERAL METHOD OF INVESTIGATING THE RELATIONS OF QUANTITIES, BY LETTERS, AND OTHER SYMBOLS. This, it must be acknowledged, is an imperfect account of the subject; as every account must necessarily be, which is comprised in the compass of a definition. Its real nature is to be learned, rather by an attentive examination of its parts, than from any summary description.

The solutions in Algebra, are of a more *general* nature, than those in common Arithmetic. The latter relate to particular numbers; the former, to whole *classes* of quantities. On this account, Algebra has been termed a kind of *universal Arithmetic*. The generality of its solutions is principally owing to the use of *letters*, instead of numeral figures, to express the several quantities which are subjected to calculation. In Arithmetic, when a problem is solved, the answer is limited to the particular numbers which are specified, in the statement of the question. But an algebraic solution may be equally applicable to all other quantities which have the same relations. This important advantage is owing to the difference between the customary use of figures, and the manner in which letters are employed in Algebra. One of the nine digits invariably expresses the same number: but a letter may be put for any number whatever. The figure 8 always signifies eight; the figure 5, five, &c. And, though one of the digits, in connection with others, may have a *local* value, different from its simple value when alone; yet the same *combination* always expresses the same number. Thus 263 has one uniform signification. And this is the case with every other combination of figures. But in Algebra, a letter may stand for any quantity which we wish it to represent. Thus *b* may be put for 2, or 10, or 50, or 1000. It must not be understood from this, however, that the letter has no

determinate value. Its value is fixed for the occasion. For the present purpose, it remains unaltered. But on a different occasion, the same letter may be put for any other number.

A calculation may be greatly abridged by the use of letters; especially when very large numbers are concerned. And when several such numbers are to be combined, as in multiplication, the process becomes extremely tedious. But a single letter may be put for a large number, as well as for a small one. The numbers 26347297, 68347823, and 27462498, for instance, may be expressed by the letters  $b$ ,  $c$  and  $d$ . The multiplying them together, as will be seen hereafter, will be nothing more than writing them, one after another, in the form of a word, and the product will be simply  $bcd$ . Thus, in Algebra, much of the labour of calculation may be saved, by the rapidity of the operations. Solutions are sometimes effected, in the compass of a few lines, which, in common Arithmetic, must be extended through many pages.

24. Another advantage obtained from the notation by letters instead of figures, is, that the several quantities which are brought into calculation, may be preserved *distinct from each other*, though carried through a number of complicated processes; whereas, in arithmetic, they are so blended together, that no trace is left of what they were, before the operation began.

25. Algebra differs farther from arithmetic, in making use of *unknown* quantities, in carrying on its operations. In arithmetic, all the quantities which enter into a calculation must be known. For they are expressed *in numbers*. And every number must necessarily be a determinate quantity. But in algebra, a letter may be put for a quantity, before its value has been ascertained. And yet it may have such relations to other quantities, with which it is connected, as to answer an important purpose in the calculation.

### NOTATION.

26. To facilitate the investigations in algebra, the several steps of the reasoning, instead of being expressed in *words*, are translated into the language of signs and symbols, which may be considered as a species of *short-hand*. This serves to place the quantities and their relations distinctly before the eye, and to bring them all into view at once. They are thus more readily compared and understood, than when re-

moved at a distance from each other, as in the common mode of writing. But before any one can avail himself of this advantage, he must become perfectly familiar with the new language.

27. The *quantities* in algebra, as has been already observed, are generally expressed by *letters*. The *first* letters of the alphabet are used, to represent *known* quantities; and the *last* letters, those which are unknown. Sometimes the quantities, instead of being expressed by letters, are set down in figures, as in common arithmetic.

28. Besides the letters and figures, there are certain characters used, to indicate the *relations* of the quantities, or the *operations* which are performed with them. Among these are the signs  $+$  and  $-$ , which are read *plus* and *minus*, or *more* and *less*. The former is prefixed to quantities which are to be *added*; the latter, to those which are to be *subtracted*. Thus  $a + b$  signifies that  $b$  is to be added to  $a$ . It is read  $a$  plus  $b$ , or  $a$  added to  $b$ , or  $a$  and  $b$ . If the expression be  $a - b$ , i. e.  $a$  minus  $b$ ; it indicates that  $b$  is to be subtracted from  $a$ .

29. The sign  $+$  is prefixed to quantities which are considered as *affirmative* or *positive*; and the sign  $-$ , to those which are supposed to be negative. For the nature of this distinction, see art. 54.

All the quantities which enter into an algebraic process, are considered, for the purposes of calculation, as either positive or negative. Before the *first* one, unless it be negative, the sign is generally omitted. But it is always to be understood. Thus  $a + b$ , is the same as  $+ a + b$ .

30. Sometimes *both*  $+$  and  $-$  are prefixed to the same letter. The sign is then said to be *ambiguous*. Thus  $a \pm b$  signifies that in certain cases, comprehended in a general solution,  $b$  is to be added to  $a$ , and, in other cases, subtracted from it.

31. When it is intended to express the difference between two quantities without deciding which is the one to be subtracted, the character  $\oslash$  or  $\smile$  is used. Thus  $a \oslash b$ , or  $a \smile b$  denotes the difference between  $a$  and  $b$ , without determining whether  $a$  is to be subtracted from  $b$ , or  $b$  from  $a$ .

32. The *equality* between two quantities or sets of quantities is expressed, by parallel lines  $=$ . Thus  $a + b = d$  signifies that  $a$  and  $b$  together are equal to  $d$ . And  $a + d = c$

$= b + g = h$  signifies that  $a$  and  $d$  equal  $c$ , which is equal to  $b$  and  $g$ , which are equal to  $h$ . So  $8 + 4 = 16 - 4 = 10 + 2 = 7 + 2 + 3 = 12$ .

33. When the first of the two quantities compared is *greater*, than the other, the character  $>$  is placed between them. Thus  $a > b$  signifies that  $a$  is greater than  $b$ .

If the first is *less* than the other, the character  $<$  is used; as  $a < b$ ; i. e.  $a$  is less than  $b$ . In both cases, the quantity towards which the character *opens*, is greater than the other.

34. A numeral figure is often prefixed to a letter. This is called a *co-efficient*. It shows how often the quantity expressed by the letter is to be taken. Thus  $2b$  signifies twice  $b$ , and  $9b$ , 9 times  $b$ , or 9 multiplied into  $b$ .

The co-efficient may be either a whole number or a fraction. Thus  $\frac{2}{3}b$  is two thirds of  $b$ . When the co-efficient is not expressed, 1 is always to be understood. Thus  $a$  is the same as  $1 a$ ; i. e. once  $a$ .

35. The co-efficient may be a *letter*, as well as a figure. In the quantity  $mb$ ,  $m$  may be considered the co-efficient of  $b$ ; because  $b$  is to be taken as many times as there are units in  $m$ . If  $m$  stands for 6, then  $mb$  is 6 times  $b$ . In  $3abc$ , 3 may be considered as the co-efficient of  $abc$ ;  $3a$ , the co-efficient of  $bc$ ; or  $3ab$ , the co-efficient of  $c$ . See art. 42.

36. A *simple* quantity is either a single letter or number, or several letters connected together, without the signs  $+$  and  $-$ . Thus  $a$ ,  $ab$ ,  $abd$ , and  $8b$  are each of them simple quantities. A *compound* quantity consists of a number of simple quantities, connected by the sign  $+$  or  $-$ . Thus  $a + b$ ,  $d - y$ ,  $b - d + 3h$ , are each compound quantities. The members of which it is composed, are called *terms*.

37. If there are *two* terms in a compound quantity, it is called a *binomial*. Thus  $a + b$  and  $a - b$  are binomials. The latter is also called a *residual* quantity, because it expresses the difference of two quantities, or the remainder, after one is taken from the other. A compound quantity consisting of *three* terms, is sometimes called a *trinomial*; one of four terms, a *quadrinomial*, &c.

38. When the several members of a compound quantity are to be subjected to the same operation, they are frequently connected by a line called a *vinculum*. Thus  $a - \overline{b + c}$  shows that the *sum* of  $b$  and  $c$  is to be subtracted from  $a$ . But  $a - b + c$  signifies that  $b$  only is to be subtracted from  $a$ ,

while  $c$  is to be added. The sum of  $c$  and  $d$ , subtracted from the sum of  $a$  and  $b$ , is  $\overline{a + b - c + d}$ . The marks used for parentheses, ( ) are often substituted, instead of a line, for a vinculum. Thus  $x - (a + c)$  is the same as  $x - \overline{a + c}$ . The equality of two sets of quantities is expressed, without using a vinculum. Thus  $a + b = c + d$  signifies, not that  $b$  is equal to  $c$ ; but that the sum of  $a$  and  $b$  is equal to the sum of  $c$  and  $d$ .

39. A single letter, or a number of letters, representing any quantities with their relations, is called an algebraic expression; and sometimes a formula. Thus  $a + b + 3d$  is an algebraic expression.

40. The character  $\times$  denotes multiplication. Thus  $a \times b$  is  $a$  multiplied into  $b$ : and  $6 \times 3$  is 6 times 3, or 6 into 3. Sometimes a point is used to indicate multiplication. Thus  $a.b$  is the same as  $a \times b$ . But the sign of multiplication is more commonly omitted, between simple quantities; and the letters are connected together, in the form of a word or syllable. Thus  $ab$  is the same as  $a.b$  or  $a \times b$ . And  $bcde$  is the same as  $b \times c \times d \times e$ . When a compound quantity is to be multiplied, a vinculum is used, as in the case of subtraction. Thus the sum of  $a$  and  $b$ , multiplied into the sum of  $c$  and  $d$ , is  $\overline{a + b} \times \overline{c + d}$ , or  $(a + b) \times (c + d)$ . And  $(6 + 2) \times 5$  is  $8 \times 5$  or 40. But  $6 + 2 \times 5$  is  $6 + 10$  or 16. When the marks of parentheses are used, the sign of multiplication is frequently omitted. Thus  $(x + y)(x - y)$  is  $(x + y) \times (x - y)$ .

41. When two or more quantities are multiplied together, each of them is called a factor. In the product  $ab$ ,  $a$  is a factor, and so is  $b$ . In the product  $x \times a + m$ ,  $x$  is one of the factors, and  $a + m$ , the other. Hence every co-efficient may be considered a factor. (Art. 35.) In the product  $3y$ , 3 is a factor, as well as  $y$ .

42. A quantity is said to be resolved into factors, when any factors are taken, which, being multiplied together, will produce the given quantity. Thus  $3ab$  may be resolved into the two factors  $3a$  and  $b$ , because  $3a \times b$  is  $3ab$ . And  $5amn$  may be resolved into the three factors  $5a$ , and  $m$ , and  $n$ . And 48 may be resolved, into the two factors  $2 \times 24$ , or  $3 \times 16$ , or  $4 \times 12$ , or  $6 \times 8$ ; or into the three factors  $2 \times 3 \times 8$ , or  $4 \times 6 \times 2$ , &c.

43. The character  $\div$  is used to show, that the quantity which precedes it, is to be *divided*, by that which follows. Thus  $a \div c$  is  $a$  divided by  $c$ : and  $a + b \div c + d$  is the sum of  $a$  and  $b$ , divided by the sum of  $c$  and  $d$ . But in algebra, division is more commonly expressed, by writing the divisor under the dividend, in the form of a vulgar fraction. Thus  $\frac{a}{b}$  is the same as  $a \div b$ : and  $\frac{c \text{ } \smile \text{ } b}{d + h}$  is the difference of  $c$  and  $b$  divided by the sum of  $d$  and  $h$ . A character prefixed to the dividing line of a fractional expression, is to be understood as referring to all the parts taken collectively; that is, to the whole value of the quotient. Thus  $a - \frac{b + c}{m + n}$  signifies that the quotient of  $b + c$  divided by  $m + n$  is to be subtracted from  $a$ . And  $\frac{c - d}{a + m} \times \frac{h + n}{x - y}$  denotes that the first quotient is to be multiplied into the second.

44. When four quantities are *proportional*, the proportion is expressed by points, in the same manner, as in the Rule of Three in arithmetic. Thus  $a : b :: c : d$  signifies that  $a$  has to  $b$ , the same ratio, which  $c$  has to  $d$ . And  $ab : cd :: a + m : b + n$ , means, that  $ab$  is to  $cd$ ; as the sum of  $a$  and  $m$ , to the sum of  $b$  and  $n$ .

45. Algebraic quantities are said to be *alike*, when they are expressed by the same *letters*, and are of the same *power*: and *unlike*, when the letters are different, or when the same letter is raised to different powers.\* Thus  $ab$ ,  $3ab$ ,  $-ab$ , and  $-6ab$ , are like quantities, because the letters are the same in each, although the signs and co-efficients are different. But  $3a$ ,  $3y$ , and  $3bx$ , are unlike quantities, because the letters are unlike, although there is no difference in the signs and co-efficients.

46. One quantity is said to be a *multiple* of another, when the former *contains* the latter a certain number of times without a remainder. Thus  $10a$  is a multiple of  $2a$ ; and  $24$  is a multiple of  $6$ .

47. One quantity is said to be a *measure* of another, when the former *is contained* in the latter, any number of times, without a remainder. Thus  $3b$  is a measure of  $15b$ : and  $7$  is a measure of  $35$ .

\* For the notation of *powers* and *roots*, see the sections on those subjects.

48. The *value* of an expression, is the number or quantity, for which the expression stands. Thus the value of  $3 + 4$  is 7; of  $3 \times 4$  is 12; of  $\frac{1}{2}$  is 2.

49. The **RECIPROCAL** of a quantity, is the quotient arising from dividing A UNIT by that quantity. Thus the reciprocal of  $a$  is  $\frac{1}{a}$ ; the reciprocal of  $a + b$  is  $\frac{1}{a + b}$ ; the reciprocal of 4 is  $\frac{1}{4}$ .

50. The relations of quantities, which, in ordinary language, are signified by *words*, are represented, in the algebraic notation, by *signs*. The latter mode of expressing these relations, ought to be made so familiar to the mathematical student, that he can, at any time, substitute the one for the other. A few examples are here added, in which, words are to be converted into signs.

1. What is the algebraic expression for the following statement, in which, the letters  $a, b, c,$  &c. may be supposed to represent any given quantities?

The product of  $a, b,$  and  $c,$  divided by the difference of  $c$  and  $d,$  is equal to the sum of  $b$  and  $c$  added to 15 times  $h.$

$$\text{Ans. } \frac{abc}{c-d} = b + c + 15h.$$

2. The product of the difference of  $a$  and  $h$  into the sum of  $b, c,$  and  $d,$  is equal to 37 times  $m,$  added to the quotient of  $b$  divided by the sum of  $h$  and  $b.$  Ans.

3. The sum of  $a$  and  $b,$  is to the quotient of  $b$  divided by  $e;$  as the product of  $a$  into  $c,$  to 12 times  $h.$  Ans.

4. The sum of  $a, b,$  and  $c$  divided by six times their product, is equal to four times their sum diminished by  $d.$  Ans.

5. The quotient of 6 divided by the sum of  $a$  and  $b,$  is equal to 7 times  $d,$  diminished by the quotient of  $b,$  divided by 36. Ans.

51. It is necessary also, to be able to reverse what is done in the preceding examples, that is, to translate the algebraic signs into common language.

What will the following expressions become, when words are substituted for the signs?

$$1. \frac{a+b}{h} = abc - 6m + \frac{a}{a+c}.$$

Ans. The sum of  $a$  and  $b$  divided by  $h,$  is equal to the product of  $a, b,$  and  $c$  diminished by 6 times  $m,$  and increased by the quotient of  $a$  divided by the sum of  $a$  and  $c.$





$$2. ab + \frac{3h - c}{x + y} = d \times \frac{a + b + c}{6 + b} - \frac{h}{6 + b}$$

$$3. a + 7(h + x) - \frac{c - 6d}{2a + 4} = (a + h) \times (b - c)$$

$$4. a - b : a c :: \frac{d - 4}{m} : 3 \times h + d + y$$

$$5. \frac{a - h}{3 + b - c} + \frac{d + ab}{2m} = \frac{ba \times d + h}{am} - \frac{cd}{h + dm}$$

52. At the close of an algebraic process, it is frequently necessary to restore the *numbers*, for which letters had been substituted, at the beginning. In doing this, the sign of multiplication must not be omitted, as it generally is, between factors, expressed by letters. Thus, if *a* stands for 3, and *b*, for four; the product *ab* is not 34, but  $3 \times 4$  i. e. 12.

In the following examples,

Let  $a = 3$

$b = 4$

$c = 2$

And  $d = 6$ .

$m = 8$ .

$n = 10$ .

Then, 1.  $\frac{a + m}{cd} + \frac{bc - n}{3d} = \frac{3 + 8}{2 \times 6} + \frac{4 \times 2 - 10}{3 \times 6}$ .

2.  $\frac{b + ad}{c + dm} - b c m n + \frac{d - 4cn}{5ab} = \text{---}$

3.  $b m d + \frac{ab - 3d}{cdm} - \frac{3bn - bc}{4a + 3cd} + \frac{b}{a} = \text{---}$

53. An algebraic expression, in which numbers have been substituted for letters, may often be rendered much more simple, by reducing several terms to one. This can not generally be done, while the letters remain. If  $a + b$  is used for the sum of two quantities, *a* can not be united in the same term with *b*. But if *a* stands for 3, and *b*, for 4, then  $a + b = 3 + 4 = 7$ . The value of an expression consisting of many terms may thus be found, by actually performing, with the numbers, the operations of addition, subtraction, multiplication, &c. indicated by the algebraic characters.

Find the value of the following expressions, in which the letters are supposed to stand for the same numbers, as in the preceding article.

$$1. \frac{ad}{c} + a + mn = \frac{3 \times 6}{2} + 3 + 8 \times 10 = 9 + 3 + 80 = 92.$$

$$2. abm + \frac{2b}{m-d} + 2n = 3 \times 4 \times 8 + \frac{2 \times 4}{8-6} + 2 \times 10 =$$

$$3. \frac{a+c}{n-m} + \frac{m-b}{m-d} - a \times \frac{n-m}{n-m} =$$

$$4. \frac{a \times d + c}{n-d} + abc - \frac{c+b \times m-d}{n-bc} = \text{---}$$

$$5. \frac{ac+5m}{2n+3} + m - cb + \frac{4d-b \times a-c}{n} =$$

### POSITIVE AND NEGATIVE QUANTITIES.\*

54. To one who has just entered on the study of algebra, there is generally nothing more perplexing, than the use of what are called *negative* quantities. He supposes he is about to be introduced to a class of quantities, which are entirely new; a sort of mathematical *nothings*, of which he can form no distinct conception. As positive quantities are *real*, he concludes that those which are negative must be *imaginary*. But this is owing to a misapprehension of the term negative, as used in the mathematics.

55. A NEGATIVE QUANTITY IS ONE WHICH IS REQUIRED TO BE SUBTRACTED. When several quantities enter into a calculation, it is frequently necessary that some of them should be *added* together, while others are *subtracted*. The former are called affirmative or positive, and are marked with the sign +; the latter are termed negative, and distinguished by the sign -. If, for instance, the profits of trade are the subject of calculation, and the *gain* is considered positive; the *loss* will be negative; because the latter must be *subtracted* from the former, to determine the clear profit. If the sums of a book account, are brought into an algebraic process, the debt and the credit are distinguished by opposite signs. If a man on a journey is, by any accident, necessitated to return several miles, this backward motion is to be considered *negative*, because that, in determining his real progress, it must

\* On the subject of Negative quantities, see Newton's Universal Arithmetic, Maseres on the Negative Sign, Mansfield's Mathematical Essays, and Maclaurin's, Simpson's, Euler's, Saunderson's and Ludlam's Algebra.

be subtracted from the distance which he has travelled in the opposite direction. If the *ascent* of a body from the earth be called positive, its *descent* will be negative. These are only different examples of the same general principle. In each of the instances, one of the quantities is to be *subtracted* from the other.

56. The terms positive and negative, as used in the mathematics, are merely *relative*. They imply that there is, either in the nature of the quantities, or in their circumstances, or in the purposes which they are to answer in calculation, some such *opposition* as requires that one should be *subtracted* from the other. But this opposition is not that of existence and non-existence, nor of one thing greater than nothing, and another less than nothing. For, in many cases, either of the signs may be, indifferently and at pleasure, applied to the very same quantity; that is, the two characters may change places. In determining the progress of a ship, for instance, her easting may be marked +, and her westing -; or the westing may be +, and the easting -. All that is necessary is, that the two signs be prefixed to the quantities, in such a manner as to show, which are to be added, and which subtracted. In different processes, they may be differently applied. On one occasion, a downward motion may be called positive, and on another occasion, negative.

57. In every algebraic calculation, some one of the quantities must be fixed upon, to be considered positive. All other quantities which will *increase* this, must be positive also. But those which will tend to *diminish* it, must be negative. In a mercantile concern, if the *stock* is supposed to be positive, the *profits* will be positive; for they *increase* the stock; they are to be *added* to it. But the *losses* will be negative; for they *diminish* the stock; they are to be *subtracted* from it. When a boat, in attempting to ascend a river, is occasionally driven back by the current; if the progress up the stream, to any particular point, is considered positive, every succeeding instance of *forward* motion will be positive, while the *backward* motion will be negative.

58. A negative quantity is frequently *greater*, than the positive one with which it is connected. But how, it may be asked, can the former be *subtracted* from the latter? The greater is certainly not *contained* in the less: how then can it be taken out of it? The answer to this is, that the

greater may be supposed first to *exhaust* the less, and then to leave a remainder equal to the difference between the two. If a man has in his possession, 1000 dollars, and has contracted a debt of 1500; the latter subtracted from the former, not only exhausts the whole of it, but leaves a balance of 500 against him. In common language, he is 500 dollars worse than nothing.

59. In this way, it frequently happens, in the course of an algebraic process, that a negative quantity is brought to *stand alone*. It has the sign of subtraction, without being connected with any other quantity, from which it is to be subtracted. This denotes that a previous subtraction has left a remainder, which is a part of the quantity subtracted. If the latitude of a ship which is 20 degrees north of the equator, is considered positive, and if she sails south 25 degrees; her motion first *diminishes* her latitude, then reduces it to *nothing*, and finally gives her 5 degrees of *south* latitude. The sign  $-$  prefixed to the 25 degrees, is retained before the 5, to show that this is what remains of the *southward* motion, after balancing the 20 degrees of north latitude. If the motion southward is only fifteen degrees, the remainder must be  $+ 5$ , instead of  $- 5$ , to show that it is a part of the ship's *northern* latitude, which has been thus far diminished, but not reduced to nothing. The balance of a book account will be positive or negative, according as the debt or the credit is the greater of the two. To determine to which side the remainder belongs, the sign must be retained, though there is no other quantity, from which this is again to be subtracted, or to which it is to be added.

60. When a quantity continually decreasing is reduced to nothing, it is sometimes said to become afterwards *less than nothing*. But this is an exceptionable manner of speaking.\* No quantity can be really less than nothing. It may be diminished, till it vanishes, and gives place to an *opposite* quantity. The latitude of a ship crossing the equator, is first made less, then nothing, and afterwards *contrary* to what it was before. The north and south latitudes may therefore be properly distinguished, by the signs  $+$  and  $-$ ; all the

\* NOTE. The expression "*less than nothing*," may not be wholly improper; if it is intended to be understood, not literally, but merely as a convenient phrase adopted for the sake of avoiding a tedious circumlocution; as we say "the sun rises," instead of saying "the earth rolls round, and brings the sun into view." The use of it in this manner, is warranted by Newton, Euler, and others.

positive degrees being on one side of 0, and all the negative, on the other; thus,

+ 6, + 5, + 4, + 3, + 1, 0, - 1, - 2, - 3, - 4, - 5, &c.

The numbers belonging to any other series of opposite quantities, may be arranged in a similar manner. So that 0 may be conceived to be a kind of *dividing point* between positive and negative numbers. On a thermometer, the degrees *above* 0 may be considered positive, and those *below* 0, negative.

61. A quantity is sometimes said to be *subtracted from* 0. By this is meant, that it belongs on the negative side of 0. But a quantity is said to be *added to* 0, when it belongs on the positive side. Thus, in speaking of the degrees of a thermometer,  $0 + 6$  means 6 degrees *above* 0; and  $0 - 6$ , 6 degrees *below* 0.

### AXIOMS.

62. The object of mathematical inquiry is, generally, to investigate some unknown quantity, and discover how *great* it is. This is effected, by comparing it with some other quantity or quantities already known. The dimensions of a stick of timber are found, by applying to it a measuring rule of known length. The *weight* of a body is ascertained, by placing it in one scale of a balance, and observing how many pounds in the opposite scale, will equal it. And any quantity is determined, when it is found to be equal to some known quantity or quantities.

Let  $a$  and  $b$  be known quantities, and  $y$ , one which is unknown. Then  $y$  will become known, if it is discovered to be equal to the sum of  $a$  and  $b$ ; that is, if

$$y = a + b.$$

An expression like this, representing the equality between one quantity or set of quantities, and another, is called an *equation*. It will be seen hereafter, that much of the business of algebra consists in finding equations, in which some unknown quantity is shown to be equal to others which are known. But it is not often the fact, that the first comparison of the quantities, furnishes the equation required. It will generally be necessary to make a number of additions, subtractions, multiplications, &c. before the unknown quantity is discovered. But in all these changes, a constant equality must be preserved, between the two sets of quantities compared. This will be done, if, in making the altera-

tions, we are guided by the following *axioms*. These are not inserted here, for the purpose of being proved; for they are self-evident. (Art. 10.) But as they must be continually introduced or implied, in demonstrations and the solutions of problems, they are placed together, for the convenience of reference.

63. Axiom 1. If the same quantity or equal quantities be *added* to equal quantities, their *sums* will be equal.

2. If the same quantity or equal quantities be *subtracted* from equal quantities, the *remainders* will be equal.

3. If equal quantities be *multiplied* into the same, or equal quantities, the *products* will be equal.

4. If equal quantities be *divided* by the same or equal quantities, the *quotients* will be equal.

5. If the same quantity be both *added to* and *subtracted from* another, the value of the latter will not be altered.

6. If a quantity be both *multiplied* and *divided* by another, the value of the former will not be altered.

7. If to unequal quantities, equals be added, the greater will give the greater sum.

8. If from unequal quantities, equals be subtracted, the greater will give the greater remainder.

9. If unequal quantities be multiplied by equals, the greater will give the greater product.

10. If unequal quantities be divided by equals, the greater will give the greater quotient.

11. Quantities which are respectively equal to any other quantity, are equal to each other.

12. The whole of a quantity is greater than a part.

This is, by no means, a *complete* list of the self-evident propositions, which are furnished by the mathematics. It is not necessary to enumerate them all. Those have been selected, to which we shall have the most frequent occasion to refer.

64. The investigations in algebra are carried on principally, by means of a series of *equations* and *proportions*. But instead of entering directly upon these, it will be necessary to attend, in the first place, to a number of processes, on which the management of equations and proportions depends. These preparatory operations are similar to the calculations under the common rules of arithmetic. We have addition, multiplication, division, involution, &c. in algebra, as well as in arithmetic. But this application of a common

name, to operations in these two branches of the mathematics, is often the occasion of perplexity and mistake. The learner naturally expects to find addition in algebra the same as addition in arithmetic. They are in fact the same, in many respects : in *all* respects perhaps, in which the steps of the one will admit of a direct comparison, with those of the other. But addition in algebra is more *extensive*, than in arithmetic. The same observation may be made, concerning several other operations in algebra. They are, in many points of view, the same as those which bear the same names in arithmetic. But they are frequently extended farther, and comprehend processes which are unknown to arithmetic. This is commonly owing to the introduction of negative quantities. The management of these requires steps which are unnecessary, where quantities of one class only are concerned. It will be important therefore, as we pass along, to mark the *difference*, as well as the *resemblance*, between arithmetic and algebra ; and, in some instances, to give a new definition, accommodated to the latter.



## SECTION II.



## ADDITION.

ART. 65. **I**N entering on an algebraic calculation, the first thing to be done, is evidently to *collect the materials*. Several distinct quantities are to be concerned in the process. These must be brought together. They must be connected in some form of expression, which will present them at once to our view, and show the relations which they have to each other. This collecting of quantities is what, in algebra, is called **ADDITION**. It may be defined, **THE CONNECTING OF SEVERAL QUANTITIES, WITH THEIR SIGNS, IN ONE ALGEBRAIC EXPRESSION.**

66. It is common to include in the definition, "uniting in one term, such quantities, as will admit of being united." But this is not so much a part of the addition itself, as a *reduction*, which accompanies or follows it. The addition may, in all cases, be performed, by merely connecting the quantities, by their proper signs. Thus  $a$  added to  $b$  is, evidently,  $a$  and  $b$ : that is, according to the algebraic notation  $a+b$ . And  $a$ , added to the sum of  $b$  and  $c$ , is  $a+b+c$ . And  $a+b$ , added to  $c+d$ , is  $a+b+c+d$ . In the same manner, if the sum of any quantities whatever, be added to the sum of any others, the expression for the whole, will contain all these quantities connected by the sign  $+$ .

67. Again, if the *difference* of  $a$  and  $b$  be added to  $c$ ; the sum will be  $a-b$  added to  $c$ , that is,  $a-b+c$ . And if  $a-b$  be added to  $c-d$ , the sum will be  $a-b+c-d$ . In one of the compound quantities added here,  $a$  is to be diminished by  $b$ , and in the other,  $c$  is to be diminished by  $d$ ; the *sum* of  $a$  and  $c$  must therefore be diminished, both by  $b$ , and by  $d$ , that is, the expression for the sum total, must contain  $-b$  and  $-d$ . On the same principle, all the quantities which, in the parts to be added, have the negative sign, must *retain* this sign, in the amount. Thus  $a+2b-c$ , added to  $d-h-m$ , is  $a+2b-c+d-h-m$ .

68. The sign must be retained also, when a positive quantity is to be added, to a *single* negative quantity. If  $a$  be added to  $-b$ , the sum will be  $-b+a$ . Here it may be objected, that the negative sign prefixed to  $b$ , shows that it is to be *subtracted*. What propriety then can there be in *adding* it? In reply to this, it may be observed, that the sign prefixed to  $b$  while standing alone, signifies that  $b$  is to be subtracted, *not from*  $a$ , but from some *other* quantity, which is not here expressed. Thus  $-b$  may represent the *loss*, which is to be subtracted from the *stock* in trade. (Art. 55.) The object of the calculation, however, may not require that the value of this stock should be specified. But the loss is to be connected with a *profit* on some other article. Suppose the profit is 2000 dollars, and the loss 400. The inquiry then is, what is the value of 2000 dollars profit, when connected with 400 dollars loss?

The answer is, evidently,  $2000-400$ , which shows that 2000 dollars are to be *added* to the stock, and 400 *subtracted* from it; or, which will amount to the same, that the *difference* between 2000 and 400 is to be added to the stock.



69. QUANTITIES ARE ADDED, then, BY WRITING THEM ONE AFTER ANOTHER, WITHOUT ALTERING THEIR SIGNS; observing always, that a quantity, to which *no* sign is prefixed, is to be considered positive. (Art. 29.)

The sum of  $a+m$ , and  $b-8$ , and  $2h-3m+d$ , and  $h-n$ , and  $r+3m-y$ , is

$$a+m+b-8+2h-3m+d+h-n+r+3m-y.$$

70. It is immaterial in what *order* the terms are arranged. The sum of  $a$  and  $b$  and  $c$  is either  $a+b+c$ , or  $a+c+b$ , or  $c+b+a$ . For it evidently makes no difference, which of the quantities is added *first*. The sum of 6 and 3 and 9, is the same as 3 and 9 and 6, or 9 and 6 and 3.

And  $a+m-n$ , is the same as  $a-n+m$ . For it is plainly of no consequence, whether we first add  $m$  to  $a$ , and afterwards subtract  $n$ ; or first subtract  $n$ , and then add  $m$ .

71. Though connecting quantities by their signs, is all which is *essential* to addition; yet it is desirable to make the expression as simple as may be, by *reducing several terms to one*. The amount of  $3a$ , and  $6b$ , and  $4a$ , and  $5b$ , is

$$3a+6b+4a+5b.$$

But this may be abridged. The first and third terms may be brought into one; and so may the second and fourth. For 3 times  $a$ , and 4 times  $a$ , make 7 times  $a$ . And 6 times  $b$ , and 5 times  $b$ , make 11 times  $b$ . The sum, when reduced, is therefore  $7a+11b$ .

For making the reductions connected with addition, two rules are given, adapted to the two cases, in one of which, the quantities and signs are alike, and in the other, the quantities are alike, but the signs are unlike. Like quantities are the same *powers* of the same *letters*. (Art. 45.) But as the addition of powers and radical quantities will be considered, in a future section, the examples given in this place, will be all of the first power.

72. CASE I. TO REDUCE SEVERAL TERMS TO ONE, WHEN THE QUANTITIES ARE ALIKE AND THE SIGNS ALIKE, ADD THE CO-EFFICIENTS, ANNEX THE COMMON LETTER OR LETTERS, AND PREFIX THE COMMON SIGN.

Thus, to reduce  $3b+7b$ , that is  $+3b+7b$  to one term, add the co-efficients 3 and 7, to the sum 10, annex the common letter  $b$ , and prefix the sign  $+$ . The expression will then be  $+10b$ . That 3 times any quantity, and 7 times the same quantity, make 10 times that quantity, needs no proof.

*Examples.*

$bc$	$3xy$	$7b + xy$	$ry + 3abh$	$cdxy + 3mg$
$2bc$	$7xy$	$8b + 3xy$	$3ry + abh$	$2cdxy + mg$
$9bc$	$xy$	$2b + 2xy$	$6ry + 4abh$	$5cdxy + 7mg$
$3bc$	$2xy$	$6b + 5xy$	$2ry + abh$	$7cdxy + 8mg$
$15bc$		$23b + 11xy$		$15cdxy + 19mg$

The mode of proceeding will be the same, if the signs are *negative*.

Thus  $-3bc - bc - 5bc$ , becomes, when reduced,  $-9bc$ .

And  $-ax - 3ax - 2ax = -6ax$ . Or thus,

$-3bc$	$-ax$	$-2ab - my$	$-3ach - 8bdy$
$-bc$	$-3ax$	$-ab - 3my$	$-ach - bdy$
$-5bc$	$-2ax$	$-7ab - 8my$	$-5ach - 7bdy$
$-9bc$		$-10ab - 12my$	

73. It may perhaps be asked here, as in art. 68, what propriety there is, in *adding* quantities, to which the negative sign is prefixed; a sign which denotes *subtraction*? The answer to this is, that when the negative sign is applied to several quantities, it is intended to indicate that these quantities are to be subtracted, *not from each other*, but from some *other* quantity, marked with the contrary sign. Suppose that, in estimating a man's property, the sum of money in his possession is marked +, and the debts which he owes are marked -. If these debts are 200, 300, 500 and 700 dollars, and if  $a$  is put for 100; they will together be  $-2a - 3a - 5a - 7a$ . And the several terms reduced to one, will evidently be  $-17a$ , that is, 1700 dollars.

74. CASE II. TO REDUCE SEVERAL TERMS TO ONE, WHEN THE QUANTITIES ARE ALIKE, BUT THE SIGNS UNLIKE, TAKE THE LESS CO-EFFICIENT FROM THE GREATER; TO THE DIFFERENCE, ANNEX THE COMMON LETTER OR LETTERS, AND PREFIX THE SIGN OF THE GREATER CO-EFFICIENT.

Thus, instead of  $8a - 6a$ , we may write  $2a$ .

And instead of  $7b - 2b$ , we may put  $5b$ .

For the simple expression, in each of these instances, is equivalent to the compound one, for which it is substituted.

To	+6b	+4b	5bc	- 2hm	- dy + 6m	3h - dx
Add	-4b	-6b	-7bc	-9hm	4dy - m	5h + 4dx
Sum	+2b		-2bc		3dy + 5m	

75. Here again, it may excite surprise, that what appears to be subtraction, should be introduced under addition. But according to what has been observed, (Art. 66,) this subtraction is, strictly speaking, no part of the addition. It belongs to a consequent *reduction*. Suppose  $6b$  is to be added to  $a-4b$ . The sum is  $a-4b+6b$ . (Art. 69.)

But this expression may be rendered more simple. As it now stands,  $4b$  is to be subtracted from  $a$ , and  $6b$  added. But the amount will be the same, if, without subtracting any thing, we add  $2b$ , making the whole  $a+2b$ . And in all similar instances, the *balance* of two or more quantities, may be substituted for the quantities themselves.

77. If two *equal* quantities have *contrary signs*, they destroy each other, and may be cancelled. Thus  $+6b-6b=0$ : And  $3 \times 6 - 18 = 0$ : And  $7bc-7bc=0$ .

Let there be any two quantities whatever, of which  $a$  is the greater, and  $b$  the less.

Their sum will be  $a+b$   
And their difference  $a-b$

The sum and difference added, will be  $2a+0$ , or simply  $2a$ . That is, if the *sum* and *difference* of any two quantities be added together, the *whole* will be *twice* the greater quantity. This is one instance, among multitudes, of the rapidity with which *general* truths are discovered and demonstrated in algebra. (Art. 23.)

78. If several positive, and several negative quantities are to be reduced to one term; first reduce those which are positive, next those which are negative, and then take the *difference* of the co-efficients, of the two terms thus found.

Ex. 1. Reduce  $13b+6b+b-4b-5b-7b$ , to one term.

$$\begin{array}{l} \text{By art. 72, } 13b+6b+b = 20b \\ \text{And } \quad \quad -4b-5b-7b = -16b \end{array} \left. \vphantom{\begin{array}{l} 13b+6b+b \\ -4b-5b-7b \end{array}} \right\}$$

By art. 74,  $20b-16b=4b$ , which is the value of all the given quantities, taken together.

Ex. 2. Reduce  $3xy - xy + 2xy - 7xy + 4xy - 9xy + 7xy - 6xy$ .

The positive terms are $3xy$	The negative terms are $-xy$
$2xy$	$-7xy$
$4xy$	$-9xy$
$7xy$	$-6xy$
<hr style="width: 50%; margin: 0 auto;"/>	<hr style="width: 50%; margin: 0 auto;"/>

And their sum is  $16xy$   $-23xy$

Then  $16xy - 23xy = -7xy$ .

Ex. 3.  $3ad - 6ad + ad + 7ad - 2ad + 9ad - 8ad - 4ad = 0$ .

4.  $2abm - abm + 7abm - 3abm + 7abm =$

5.  $axy - 7axy + 8axy - axy - 8axy + 9axy =$

79. If the *letters*, in the several terms to be added, are different, they can only be placed after each other, with their proper signs. They cannot be united in one simple term. If  $4b$ , and  $-6y$ , and  $3x$ , and  $17h$ , and  $-5d$ , and  $6$ , be added; their sum will be

$4b - 6y + 3x + 17h - 5d + 6$ . (Art 69.)

Different letters can no more be united in the same term, than dollars and guineas can be added, so as to make a single sum. Six guineas and 4 dollars are neither ten guineas nor ten dollars. Seven hundred, and five dozen, are neither 12 hundred nor 12 dozen. But, in such cases, the algebraic signs serve to show how the different quantities stand related to each other; and to indicate future operations, which are to be performed, whenever the letters are converted into numbers. In the expression  $a + 6$ , the two terms can not be united in one. But if  $a$  stands for 15, and if, in the course of a calculation, this number is restored; then  $a + 6$  will become  $15 + 6$ , which is equivalent to the single term 21. In the same manner,  $a - 6$  becomes  $15 - 6$ , which is equal to 9. The signs keep in view the relations of the quantities, till an opportunity occurs of reducing several terms to one.

80. When the quantities to be added contain several terms which are *alike*, and several which are *unlike*, it will be convenient to arrange them in such a manner, that the similar terms may stand one under another.

To	$3bc - 6d + 2b - 3y$	}	These may be arranged thus,
Add	$-3bc + x - 3d + bg$		$3bc - 6d + 2b - 3y$
And	$2d + y + 3x + b$		$-3bc - 3d \quad + x + bg$
			$2d \quad + y + 3x \quad + b$
			<hr style="width: 50%; margin: 0 auto;"/>

The sum will be

$-7d + 2b - 2y + 4x + bg + b$ .

*Examples.*

1. Add and reduce  $ab+8$  to  $cd-3$  and  $5ab-4m+2$ .  
The sum is  $6ab+7+cd-4m$ .

2. Add  $x+3y-dx$ , to  $7-x-8+hm$ .  
Ans.  $3y-dx-1+hm$ .

3. Add  $abm-3x+bm$ , to  $y-x+7$ , and  $5x-6y+9$ .

4. Add  $3am+6-7xy-8$ , to  $10xy-9+5am$ .

5. Add  $6ahy+7d-1+mxy$ , to  $3ahy-7d+17-mxy$ .

6. Add  $7ad-h+8xy-ad$ , to  $5ad+h-7xy$ .

7. Add  $3ab-2ay+x$ , to  $ab-ay+bx-h$ .

8. Add  $2by-3ax+2a$ , to  $3bx-by+a$ .

*The sum is infinity.*



## SECTION III.



## SUBTRACTION.

ART. 81. **A**DDITION is bringing quantities together, to find their amount. On the contrary, SUBTRACTION IS FINDING THE DIFFERENCE OF TWO QUANTITIES, OR SETS OF QUANTITIES.

Particular rules might be given, for the several cases in subtraction. But it is more convenient to have one general rule, founded on the principle, that *taking away a positive quantity*, from an algebraic expression, is the same in effect, as *annexing an equal negative quantity*; and *taking away a negative quantity* is the same, as *annexing an equal positive one*.

Suppose $+b$ is to be subtracted from	$a+b$ .
Taking away $+b$ , from $a+b$ , leaves	$a$ .
And annexing $-b$ , to $a+b$ , gives	$a+b-b$ .
But by axiom 5th, $a+b-b$ is equal to	$a$

That is, *taking away a positive term*, from an algebraic expression, is the same in effect, as *annexing an equal negative term*.

Again, suppose $-b$ is to be subtracted from	$a-b$
Taking away $-b$ , from $a-b$ , leaves	$a$
And annexing $+b$ , to $a-b$ , gives	$a-b+b$
But $a-b+b$ is equal to	$a$

That is, *taking away a negative term*, is equivalent to *annexing a positive one*. If an estate is encumbered with a debt; to cancel this debt, is to add so much to the value of the estate. Subtracting an item from one side of a book account, will produce the same alteration in the balance, as adding an equal sum to the opposite side.

To place this in another point of view.

If $m$ is added to $b$ , the sum is by the notation,	$b+m$ }
But if $m$ is subtracted from $b$ , the remainder is	$b-m$ }
So if $m$ and $h$ are each added to $b$ , the sum is	$b+m+h$ }
But if $m$ and $h$ are each subtracted from $b$ , the remainder is	$b-m-h$ }

The only difference then between adding a positive quantity and subtracting it, is, that the *sign* is changed from  $+$  to  $-$ .

Again, if  $m-n$  is subtracted from  $b$ , the remainder is,

$$b-m+n.$$

For the *less* the quantity subtracted, the *greater* will be the remainder. But in the expression  $m-n$ ,  $m$  is diminished by  $n$ ; therefore,  $b-m$  must be *increased* by  $n$ ; so as to become  $b-m+n$ : that is,  $m-n$  is subtracted from  $b$ , by changing  $+m$  into  $-m$ , and  $-n$  into  $+n$ , and then writing them after  $b$ , as in addition. The explanation will be the same, if there are *several* quantities which have the negative sign. Hence,

82. TO PERFORM SUBTRACTION IN ALGEBRA, CHANGE THE SIGNS OF ALL THE QUANTITIES TO BE SUBTRACTED, OR SUPPOSE THEM TO BE CHANGED, FROM  $+$  TO  $-$ , OR FROM  $-$  TO  $+$ , AND THEN PROCEED AS IN ADDITION.

The signs are to be changed, in the *subtrahend* only. Those in the *minuend* are not to be altered. Although the rule here given is adapted to every case of subtraction; yet there may be an advantage in giving some of the examples in distinct classes.

83. In the first place, the signs may be *alike*, and the *minuend greater* than the *subtrahend*.

From	+28	16 <i>b</i>	14 <i>da</i>	-28	-16 <i>b</i>	-14 <i>da</i>
Subtract	+16	12 <i>b</i>	6 <i>da</i>	-16	-12 <i>b</i>	-6 <i>da</i>
	<hr/>	<hr/>	<hr/>	<hr/>	<hr/>	<hr/>
Difference	+12	4 <i>b</i>	8 <i>da</i>	-12	-4 <i>b</i>	-8 <i>da</i>

Here, in the first example, the + before 16 is supposed to be changed into -, and then, the signs being unlike, the two terms are brought into one, by the second case of reduction in addition. (Art. 74.) The two next examples are subtracted in the same way. In the three last, the - in the subtrahend, is supposed to be changed into +. It may be well for the learner, at first, to write out the examples; and actually to change the signs, instead of merely conceiving them to be changed. When he has become familiar with the operation, he can save himself the trouble of transcribing.

This case is the same as subtraction in *arithmetic*. The two next cases do not occur in common arithmetic.

84. In the second place, the signs may be alike, and the minuend *less* than the subtrahend.

From	+16 <i>b</i>	12 <i>b</i>	6 <i>da</i>	-16	-12 <i>b</i>	-6 <i>da</i>
Sub.	+28 <i>b</i>	16 <i>b</i>	14 <i>da</i>	-28	-16 <i>b</i>	-14 <i>da</i>
	<hr/>	<hr/>	<hr/>	<hr/>	<hr/>	<hr/>
Dif.	-12 <i>b</i>	-4 <i>b</i>	-8 <i>da</i>	+12	4 <i>b</i>	8 <i>da</i>

The same quantities are given here, as in the preceding article, for the purpose of comparing them together. But the minuend and subtrahend are made to change places. The mode of subtracting is the same. In this class, a *greater* quantity is taken from a *less*: in the preceding, a *less* from a *greater*. By comparing them, it will be seen, that there is no difference in the answers, except that the *signs* are *opposite*. Thus  $16b - 12b$  is the same as  $12b - 16b$ , except that one is  $+4b$ , and the other  $-4b$ : That is, a greater quantity subtracted from a less, gives the same result, as a less subtracted from a greater, except that the one is positive, and the other negative. See art. 58 and 59.

85. In the third place, the *signs* may be *unlike*.

From	+28	+16 <i>b</i>	+14 <i>da</i>	-28	-16 <i>b</i>	-14 <i>da</i>
Sub.	-16	-12 <i>b</i>	-6 <i>da</i>	+16	+12 <i>b</i>	+6 <i>da</i>
	<hr/>	<hr/>	<hr/>	<hr/>	<hr/>	<hr/>
Dif.	+44	28 <i>b</i>	20 <i>da</i>	-44	-28 <i>b</i>	-20 <i>da</i>

From these examples, it will be seen that the *difference* between a positive and a negative quantity, may be *greater* than either of the two quantities. In a thermometer, the difference between 28 degrees above cypher, and 16 below, is 44 degrees. The difference between gaining 1000 dollars in trade, and losing 500, is equivalent to 1500 dollars.

86. Subtraction may be *proved*, as in arithmetic, by adding the remainder to the subtrahend. The sum ought to be equal to the minuend, upon the obvious principle, that the difference of two quantities added to one of them, is equal to the other. This serves not only to correct any particular error, but to verify the general rule.

From	$2xy - 1$	$h + 3bx$	$hy - ah$	$nd - 7by$
Sub.	$-xy + 7$	$3h - 9bx$	$5hy - 6ah$	$5nd - by$
	$3xy - 8$		$-4hy + 5ah$	
Dif.				
From	$3abm - xy$	$-17 + 4ax$	$ax + 7b$	$3ah + axy$
Sub.	$-7abm + 6xy$	$-20 - ax$	$-4ax + 15b$	$-7ah + axy$
	$10abm - 7xy$		$5ax - 8b$	
Rem.				

87. When there are *several terms alike*, they may be reduced as in addition.

1. From  $ab$ , subtract  $3am + am + 7am + 2am + 6am$ .

Ans.  $ab - 3am - am - 7am - 2am - 6am = ab - 19am$ . (Art. 72.)

2. From  $y$ , subtract  $-a - a - a - a$ .

Ans.  $y + a + a + a + a = y + 4a$ .

3. From  $ax - bc + 3ax + 7bc$ , subtract  $4bc - 2ax + bc + 4ax$ .

Ans.  $ax - bc + 3ax + 7bc - 4bc + 2ax - bc - 4ax = 2ax + bc$ .  
(Art. 78.)

4. From  $ad + 3dc - bx$ , subtract  $3ad + 7bx - dc + ad$ .

88. When the *letters* in the minuend are different from those in the subtrahend, the latter are subtracted, by first changing the signs, and then placing the several terms one after another, as in addition. (Art. 79.)

From  $3ab + 8 - my + dh$ , subtract  $x - dr + 4hy - bmx$ .

Ans.  $3ab + 8 - my + dh - x + dr - 4hy + bmx$ .



88.b. The sign  $-$ , placed before the marks of *parenthesis* which include a number of quantities, requires, that when these marks are removed, the signs of *all* the quantities thus included, should be changed.

Thus  $a - (b - c + d)$  signifies that the quantities  $b$ ,  $-c$ , and  $+d$ , are to be subtracted from  $a$ . The expression will then become  $a - b + c - d$ .

2.  $13ad + xy + d - (7ad - xy + d + hm - ry) = 6ad + 2xy - hm + ry.$

3.  $7abc - 8 + 7x - (3abc - 8 - dx + r) = 4abc + 7x + dx - r.$


4.  $3ad + h - 2y - (7y + 3h - mx + 4ad - hy - ad) =$

5.  $6am - dy + 8 - (16 + 3dy - 8 + am - e + r) =$

6.  $7ay - 2x + 5 - (4 + h - ay + x + 3b) =$

88.c. On the other hand, when a number of quantities are introduced within the marks of parenthesis, with  $-$  immediately preceding; the signs must be changed.

Thus  $-m + b - dx + 3h = -(m - b + dx - 3h)$

*All human things are subject to change!*  


SECTION IV.



MULTIPLICATION.\*

ART. 89. **I**N addition, one quantity is connected with another. It is frequently the case, that the quantities brought together are *equal*; that is, a quantity is added to *itself*.

As  $a + a = 2a$

$a + a + a + a = 4a$

$a + a + a = 3a$

$a + a + a + a + a = 5a$ , &c.

This repeated addition of a quantity to itself, is what was originally called *multiplication*. But the term, as it is now

\* Newton's Universal Arithmetic, p. 4. Maseres on the Negative Sign, Sec. II. Camus' Arithmetic, Book II. Chap. 3. Euler's Algebra, Sec. I. and II. Chap 3. Simpson's Algebra, Sec. IV. Maclaurin, Saunderson, Lacroix, Ludlam.

used, has a more extensive signification. We have frequent occasion to repeat, not only the *whole* of a quantity, but a certain *portion* of it. If the stock of an incorporated company is divided into shares, one man may own ten of them, another five, and another a *part* only of a share, say two fifths. When a dividend is made, of a certain sum on a share, the first is entitled to *ten* times this sum, the second to *five* times, and the third to only *two fifths* of it. As the apportioning of the dividend, in each of these instances, is upon the same principle, it is called multiplication in the last, as well as in the two first.

Again, suppose a man is obligated to pay an annuity of 100 dollars a year. As this is to be *subtracted* from his estate, it may be represented by  $-a$ . As it is to be subtracted *year after year*, it will become, in four years,  $-a-a-a-a=-4a$ . This *repeated subtraction* is also called multiplication. According to this view of the subject ;

90. MULTIPLYING BY A WHOLE NUMBER IS TAKING THE MULTIPLICAND AS MANY TIMES, AS THERE ARE UNITS IN THE MULTIPLIER.

Multiplying by 1, is taking the multiplicand *once*, as  $a$ .

Multiplying by 2, is taking the multiplicand *twice*, as  $a+a$ .

Multiplying by 3, is taking the multiplicand *three times*, as  $a+a+a$ , &c.

MULTIPLYING BY A FRACTION IS TAKING A CERTAIN PORTION OF THE MULTIPLICAND AS MANY TIMES, AS THERE ARE LIKE PORTIONS OF AN UNIT IN THE MULTIPLIER.\*

Multiplying by  $\frac{1}{5}$ , is taking  $\frac{1}{5}$  of the multiplicand, *once*, as  $\frac{1}{5}a$ .

Multiplying by  $\frac{2}{5}$ , is taking  $\frac{1}{5}$  of the multiplicand, *twice*, as  $\frac{1}{5}a+\frac{1}{5}a$ .

Multiplying by  $\frac{3}{5}$ , is taking  $\frac{1}{5}$  of the multiplicand *three times*.

Hence, if the multiplier is *a unit*, the product is *equal* to the multiplicand : If the multiplier is *greater* than a unit, the product is *greater* than the multiplicand : And if the multiplier is *less* than a unit, the product is *less* than the multiplicand.

MULTIPLICATION BY A NEGATIVE QUANTITY, HAS THE SAME RELATION TO MULTIPLICATION BY A POSITIVE QUANTITY, WHICH SUBTRACTION HAS TO ADDITION. In the one, the sum of the repetitions of the multiplicand is to be *added*, to the other quantities with which this multiplier is connected. In the other, the sum of these repetitions is to be *sub-*

\* See Note C.

tracted from the other quantities. This subtraction is performed at the time of multiplying, by changing the sign of the product. See Art. 107 and 108.

91. Every multiplier is to be considered a *number*. We sometimes speak of multiplying by a given *weight*, or *measure*, a sum of *money*, &c. But this is abbreviated language. If construed literally, it is absurd. Multiplying is taking either the whole or a part of a quantity, a certain *number of times*. To say that one quantity is repeated as many times, as another is *heavy*, is nonsense. But if a part of the weight of a body be fixed upon as a *unit*, a quantity may be multiplied by a *number* equal to the number of these parts contained in the body. If a diamond is sold by weight, a particular price may be agreed upon for each *grain*. A grain is here *the unit*; and it is evident that the value of the diamond, is equal to the given price repeated as many times, as there are grains in the whole weight. We say concisely, that the price is multiplied by the *weight*; meaning that it is multiplied by a *number* equal to the number of grains in the weight. In a similar manner, any quantity whatever may be supposed to be made up of parts, each being considered a *unit*, and any number of these may become a multiplier.

92. As multiplying is taking the whole or a part of a quantity a certain number of times, it is evident that the *product* must be of the same nature as the *multiplicand*.

If the multiplicand is an abstract *number*; the product will be a number.

If the multiplicand is *weight*, the product will be weight. If the multiplicand is a *line*, the product will be a line. Repeating a quantity does not alter its nature. It is frequently said, that the product of two lines is a *surface*, and that the product of three lines is a *solid*. But these are abbreviated expressions, which if interpreted literally are not correct. See Section xxi.

93. The multiplication of *fractions* will be the subject of a future section. We have first to attend to multiplication by positive whole numbers. This, according to the definition (Art. 90.) is taking the multiplicand as many times, as there are units in the multiplier. Suppose *a* is to be multiplied by *b*, and that *b* stands for 3. There are, then, three units in the multiplier *b*. The multiplicand must therefore be taken three times; thus,  $a + a + a = 3a$ , or  $ba$ .

So that, multiplying two letters together is nothing more, than writing them one after the other, either with, or without the sign of multiplication between them. Thus  $b$  multiplied into  $c$ , is  $b \times c$ , or  $bc$ . And  $x$  into  $y$ , is  $x \times y$ , or  $x.y$ , or  $xy$ .

94. If more than two letters are to be multiplied, they must be connected in the same manner. Thus  $a$  into  $b$  and  $c$ , is  $cba$ . For by the last article,  $a$  into  $b$ , is  $ba$ . This product is now to be multiplied into  $c$ . If  $c$  stands for 5, then  $ba$  is to be taken five times, thus,

$$ba + ba + ba + ba + ba = 5ba, \text{ or } cba.$$

The same explanation may be applied to any number of letters. Thus  $am$  into  $xy$ , is  $amxy$ . And  $bh$  into  $mrx$ , is  $bhmrx$ .

95. It is immaterial in what order the letters are arranged. The product  $ba$  is the same as  $ab$ . Three times five is equal to five times three. Let the number 5 be represented by as many points, in a horizontal line; and the number 3, by as many points in a perpendicular line.

```

      . . . . .
      . . . . .
      . . . . .
      . . . . .
  
```

Here it is evident that the whole number of points is equal, either to the number in the horizontal row three times repeated, or to the number in the perpendicular row five times repeated; that is, to  $5 \times 3$ , or  $3 \times 5$ . This explanation may be extended to a series of factors consisting of any numbers whatever. For the product of two of the factors may be considered as one number. This may be placed before or after a third factor: the product of three, before or after a fourth, &c.

Thus  $24 = 4 \times 6$  or  $6 \times 4 = 4 \times 3 \times 2$  or  $4 \times 2 \times 3$  or  $2 \times 3 \times 4$ .

The product of  $a, b, c$ , and  $d$ , is  $abcd$ , or  $acdb$ , or  $dcba$ , or  $badc$ .

It will generally be convenient, however, to place the letters in alphabetical order.

96. WHEN THE LETTERS HAVE NUMERICAL CO-EFFICIENTS, THESE MUST BE MULTIPLIED TOGETHER, AND PREFIXED TO THE PRODUCT OF THE LETTERS.

Thus  $3a$  into  $2b$  is  $6ab$ . For if  $a$  into  $b$  is  $ab$ , then 3 times  $a$  into  $b$ , is evidently  $3ab$ : and if, instead of multiplying by  $b$ , we multiply by twice  $b$ , the product must be twice as great, that is  $2 \times 3ab$  or  $6ab$ .

Mult.	$9ab$	$12hy$	$3dh$	$2ad$	$7bdh$	$3ay$
Into	$3xy$	$2rx$	$my$	$13hmg$	$x$	$8mx$
Prod.	$27abxy$		$3dhmy$		$7bdhx$	

97. If either of the factors consist of figures *only*, these must be multiplied into the co-efficients and letters of the other factors.

Thus  $3ab$  into  $4$ , is  $12ab$ . And  $36$  into  $2x$ , is  $72x$ . And  $24$  into  $hy$ , is  $24hy$ .

98. If the multiplicand is a *compound* quantity, *each of its terms* must be multiplied into the multiplier. Thus  $b+c+d$  into  $a$  is  $ab+ac+ad$ . For the whole of the multiplicand is to be taken as many times, as there are units in the multiplier. If then  $a$  stands for  $3$ , the repetitions of the multiplicand are

$$\begin{array}{r} b+c+d \\ b+c+d \\ b+c+d \\ \hline \end{array}$$

And their sum is  $3b+3c+3d$ , that is,  $ab+ac+ad$ .

Mult.	$d+2xy$	$2h+m$	$3hl+1$	$2hm+3+dr$
Into	$3b$	$6dy$	$my$	$4b$
Prod.	$3bd+6bxy$		$3hlmy+my$	

99. The preceding instances must not be confounded with those in which several *factors* are connected by the sign  $\times$ , or by a point. In the latter case, the multiplier is to be written before the other factors *without being repeated*. The product of  $b \times d$  into  $a$ , is  $ab \times d$ , and not  $ab \times ad$ . For  $b \times d$  is  $bd$ , and this into  $a$ , is  $abd$ . (Art. 94.) The expression  $b \times d$  is not to be considered, like  $b+d$ , a *compound quantity* consisting of two terms. Different terms are always separated by  $+$  or  $-$ . (Art. 36.) The product of  $b \times h \times m \times y$  into  $a$ , is  $a \times b \times h \times m \times y$  or  $abhmy$ . But  $b+h+m+y$  into  $a$ , is  $ab+ah+am+ay$ .

100. If *both* the factors are compound quantities, *each term in the multiplier must be multiplied into each in the multiplicand*.

Thus  $a+b$  into  $c+d$  is  $ac+ad+bc+bd$ .

For the units in the multiplier  $a+b$  are equal to the units in  $a$  added to the units in  $b$ . Therefore the product produced by  $a$ , must be added to the product produced by  $b$ .

The product of  $c+d$  into  $a$  is  $ac+ad$  } ART. 98.  
 The product of  $c+d$  into  $b$  is  $bc+bd$  }

The product of  $c+d$  into  $a+b$  is therefore  $ac+ad+bc+bd$ .

Mult. $3x+d$	$4ay+2b$	$a+1$
Into $2a+hm$	$3c+rx$	$3x+4$
Prod. $6ax+2ad+3hmx+dhm$		
		$3ax+3x+4a+4$

Mult.  $2h+7$  into  $6d+1$ . Prod.  $12dh+42d+2h+7$ .

Mult.  $dy+rx+h$  into  $6m+4+7y$ . Prod.

Mult.  $7+6b+ad$  into  $3r+4+2h$ . Prod.

101. When several terms in the product are *alike*, it will be expedient to *set one under the other*, and then to unite them, by the rules for the reduction in addition.

Mult. $b+a$	$b+c+2$	$a+y+1$
Into $b+a$	$b+c+3$	$3b+2x+7$
Prod. $bb+ab$		
	$bb+bc+2b$	
	$+ab+aa$	$bc+cc+2c$
		$+3b+3c+6$
Prod. $bb+2ab+aa$		
		$bb+2bc+5b+cc+5c+6$

Mult.  $3a+d+4$  into  $2a+3d+1$ . Prod.

Mult.  $b+cd+2$  into  $3b+4cd+7$ . Prod.

Mult.  $3b+2x+h$  into  $a \times d \times 2x$ . Prod.

103. It will be easy to see that when the multiplier and multiplicand consist of any quantity *repeated as a factor*, this factor will be repeated in the product, as many times as in the multiplier and multiplicand together.

Mult.  $a \times a \times a$  Here  $a$  is repeated *three times* as a factor.

Into  $a \times a$  Here it is repeated *twice*.

Prod.  $a \times a \times a \times a \times a$ . Here it is repeated *five times*.

The product of  $bbbb$  into  $bbb$ , is  $bbbbbbb$ .

The product of  $2x \times 3x \times 4x$  into  $5x \times 6x$ , is  $2x \times 3x \times 4x \times 5x \times 6x$ .

104. But the *numeral co-efficients* of several fellow-factors may be brought together by multiplication.

Thus  $2a \times 3b$  into  $4a \times 5b$  is  $2a \times 3b \times 4a \times 5b$ , or  $120aabb$ .

For the co-efficients are *factors*, (Art. 41.) and it is immaterial in what *order* these are arranged. (Art. 95.) So that

$$2a \times 3b \times 4a \times 5b = 2 \times 3 \times 4 \times 5 \times a \times a \times b \times b = 120aabb.$$

The product of  $3a \times 4bh$  into  $5m \times 6y$ , is  $360abhmy$ .

The product of  $4b \times 6d$  into  $2x+1$ , is  $48bdx+24bd$ .

105. The examples in multiplication thus far have been confined to *positive* quantities. It will now be necessary to consider in what manner the result will be affected, by multiplying positive and negative quantities together. We shall find,

That + into + produces +  
 - into + -  
 + into - -  
 - into - +

All these may be comprised in one general rule, which it will be important to have always familiar. IF THE SIGNS OF THE FACTORS ARE ALIKE, THE SIGN OF THE PRODUCT WILL BE AFFIRMATIVE; BUT IF THE SIGNS OF THE FACTORS ARE UNLIKE, THE SIGN OF THE PRODUCT WILL BE NEGATIVE.

106. The first case, that of + into +, needs no farther illustration. The second is - into +, that is, the multiplicand is negative, and the multiplier positive. Here  $-a$  into  $+4$  is  $-4a$ . For the repetitions of the multiplicand are,  
 $-a - a - a - a = -4a$ .

Mult. $b-3a$	$2a-m$	$h-3d-4$	$a-2-7d-x$
Into $6y$	$3h+x$	$2y$	$3b+h$
Prod. $6by-18ay$		$2hy-6dy-8y$	

107. In the two preceding cases, the affirmative sign prefixed to the multiplier shows, that the repetitions of the multiplicand are to be *added*, to the other quantities with which the multiplier is connected. But in the two remaining cases, the negative sign prefixed to the multiplier, indicates that the sum of the repetitions of the multiplicand are to be *subtracted* from the other quantities. (Art. 90.) And this subtraction is performed, at the time of multiplying, by making the sign of the product opposite to that of the multiplicand.

Thus  $+a$  into  $-4$ , is  $-4a$ . For the repetitions of the multiplicand are,

$$+a+a+a+a=+4a.$$

But this sum is to be *subtracted*, from the other quantities with which the multiplier is connected. It will then become  $-4a$ . (Art. 82.)

Thus, in the expression  $b-(4 \times a)$ , it is manifest that  $4 \times a$  is to be subtracted from  $b$ . Now  $4 \times a$  is  $4a$ , that is,  $+4a$ . But, to subtract this from  $b$ , the sign  $+$  must be changed into  $-$ . So that  $b-(4 \times a)$  is  $b-4a$ . And  $a \times -4$  is therefore  $-4a$ .

Again, suppose the multiplicand is  $a$ , and the multiplier  $(6-4)$ . As  $(6-4)$  is equal to 2, the product will be equal to  $2a$ . This is *less* than the product of 6 into  $a$ . To obtain then the product of the compound multiplier  $(6-4)$  into  $a$ , we must *subtract* the product of the negative part, from that of the positive part.

$$\begin{array}{l} \text{Multiplying} \\ \text{Into} \end{array} \left. \begin{array}{c} a \\ 6-4 \end{array} \right\} \text{ is the same as } \left\{ \begin{array}{l} \text{Multiplying} \\ \text{Into} \end{array} \right. \left. \begin{array}{c} a \\ 2 \end{array} \right\}$$

And the product  $6a-4a$ , is the same as the product  $2a$ .

Therefore  $a$  into  $-4$ , is  $-4a$ .

But if the multiplier had been  $(6+4)$ , the two products must have been *added*.

$$\begin{array}{l} \text{Multiplying} \\ \text{Into} \end{array} \left. \begin{array}{c} a \\ 6+4 \end{array} \right\} \text{ is the same as } \left\{ \begin{array}{l} \text{Multiplying} \\ \text{Into} \end{array} \right. \left. \begin{array}{c} a \\ 10 \end{array} \right\}$$

And the prod.  $6a+4a$  is the same as the product  $10a$

This shows at once the difference between multiplying by a *positive* factor, and multiplying by a *negative* one. In the former case, the sum of the repetitions of the multiplicand is to be *added to*, in the latter, *subtracted from*, the other quantities, with which the multiplier is connected. For every negative quantity must be supposed to have a reference to some other which is positive; though the two may not always stand in connection, when the multiplication is to be performed.

$$\begin{array}{l} \text{Mult. } a+b \\ \text{Into } b-x \end{array}$$

$$\begin{array}{l} 3dy+hx+2 \\ mr-ab \end{array}$$

$$\begin{array}{l} 3h+3 \\ ad-6 \end{array}$$

$$\text{Prod. } ab+bb-ax-bx$$

$$3adh+3ad-18h-18$$

108. If *two negatives* be multiplied together, the product will be affirmative:  $-4 \times -a = +4a$ . In this case, as in the



preceding, the repetitions of the multiplicand are to be *subtracted*, because the multiplier has the negative sign. These repetitions, if the multiplicand is  $-a$ , and the multiplier  $-4$ , are  $-a - a - a - a = -4a$ . But this is to be subtracted by changing the sign. It then becomes  $+4a$ .

Suppose  $-a$  is multiplied into  $(6-4)$ . As  $6-4=2$ , the product is evidently, *twice* the multiplicand, that is  $-2a$ . But if we multiply  $-a$  into 6 and 4 separately;  $-a$  into 6 is  $-6a$ , and  $-a$  into 4 is  $-4a$ . (Art. 106.) As, in the multiplier, 4 is to be subtracted from 6; so, in the product,  $-4a$  must be subtracted from  $-6a$ . Now  $-4a$  becomes by subtraction  $+4a$ . The whole product then is  $-6a+4a$ , which is equal to  $-2a$ . Or thus,

$$\begin{array}{l} \text{Multiplying} \\ \text{Into} \end{array} \left. \begin{array}{l} -a \\ 6-4 \end{array} \right\} \text{ is the same as } \left\{ \begin{array}{l} \text{Multiplying} \\ \text{Into} \end{array} \right. \left. \begin{array}{l} -a \\ 2 \end{array} \right.$$

And the prod.  $-6a+4a$ , is equal to the product  $-2a$ .

It is often considered a great mystery, that the product of two negatives should be affirmative. But it amounts to nothing more than this, that the subtraction of a negative quantity, is equivalent to the addition of an affirmative one; (Art. 81.) and, therefore, that the *repeated* subtraction of a negative quantity, is equivalent to a *repeated* addition of an affirmative one. Taking off from a man's hands a debt of ten dollars every month, is adding ten dollars a month to the value of his property.

Mult. $a-4$	$3d-hy-2x$	$3ay-b$
Into $3b-6$	$4b-7$	$6x-1$
Prod. $3ab-12b-6a+24$		$18axy-6bx-3ay+b$

Multiply  $3ad-ah-7$  into  $4-dy-hr$ .  
 Multiply  $2hy+3m-1$  into  $4d-2x+3$ .

109. As a negative multiplier changes the sign of the quantity which it multiplies; if there are *several* negative factors to be multiplied together,

The *two first* will make the product *positive*;

The *third* will make it *negative*;

The *fourth* will make it *positive*, &c.

Thus $-a \times -b = +ab$	} the product of	{	<i>two factors</i>
$+ab \times -c = -abc$			<i>three,</i>
$-abc \times -d = +abcd$			<i>four,</i>
$+abcd \times -e = -abcde$			<i>five.</i>

That is, the product of any *even* number of negative factors is *positive*; but the product of any *odd* number of negative factors is *negative*.

Thus  $-a \times -a = aa$       And  $-a \times -a \times -a \times -a = aaaa$   
 $-a \times -a \times -a = -aaa$      $-a \times -a \times -a \times -a \times -a = -aaaaa$

The product of several factors which are *all positive*, is invariably positive.

110. Positive and negative terms may frequently *balance* each other, so as to disappear in the product. (Art. 77.) A star is sometimes put in the place of the deficient term.

Mult. $a - b$	$mm - yy$	$aa + ab + bb$
Into $a + b$	$mm + yy$	$a - b$
<hr style="width: 100%;"/>	<hr style="width: 100%;"/>	<hr style="width: 100%;"/>
$aa - ab$		$aaa + aab + abb$
$+ ab - bb$		$- aab - abb - bbb$
<hr style="width: 100%;"/>	<hr style="width: 100%;"/>	<hr style="width: 100%;"/>
Prod. $aa$ * $-bb$		$aaa$ * * $-bbb$
<hr style="width: 100%;"/>	<hr style="width: 100%;"/>	<hr style="width: 100%;"/>

111. For many purposes, it is sufficient merely to *indicate* the multiplication of compound quantities, without actually multiplying the several terms. Thus the product of  $a + b + c$  into  $h + m + y$ , is  $(a + b + c) \times (h + m + y)$ . (Art 40.) The product of

$a + m$  into  $h + x$  and  $d + y$ , is  $(a + m) \times (h + x) \times (d + y)$ .  
 By this method of representing multiplication, an important advantage is often gained, in preserving the factors distinct from each other.

When the several terms are multiplied in form, the expression is said to be *expanded*. Thus

$(a + b) \times (c + d)$  becomes when expanded  $ac + ad + bc + bd$ .

112. With a given multiplicand, the less the multiplier, the less will be the product. If then the multiplier be reduced to *nothing*, the *product* will be nothing. Thus  $a \times 0 = 0$ . And if 0 be one of *any number* of fellow-factors, the product of the whole will be nothing.

Thus  $ab \times c \times 3d \times 0 = 3abcd \times 0 = 0$ .

And  $(a + b) \times (c + d) \times (h - m) \times 0 = 0$ .

113. Although, for the sake of illustrating the different points in multiplication, the subject has been drawn out into a considerable number of particulars; yet it will scarcely be

necessary for the learner, after he has become familiar with the examples, to burden his memory with any thing more than the following general rule.

MULTIPLY THE LETTERS AND CO-EFFICIENTS OF EACH TERM IN THE MULTIPLICAND, INTO THE LETTERS AND CO-EFFICIENTS OF EACH TERM IN THE MULTIPLIER ; AND PREFIX, TO EACH TERM OF THE PRODUCT, THE SIGN REQUIRED BY THE PRINCIPLE, THAT LIKE SIGNS PRODUCE +, AND DIFFERENT SIGNS -.

Mult.  $a+3b-2$  into  $4a-6b-4$ .

Mult.  $4ab \times x \times 2$  into  $3my-1+h$ .

Mult.  $(7ah-y) \times 4$  into  $4x \times 3 \times 5 \times d$ . — — —

Mult.  $(6ab-hd+1) \times 2$  into  $(8+4x-1) \times d$ .

Mult.  $3ay+y-4+h$  into  $(d+x) \times (h+y)$ .

Mult.  $6ax-(4h-d)$  into  $(b+1) \times (h+1)$ .

Mult.  $7ay-1+h \times (d-x)$  into  $-(r+3-4m)$ .



## SECTION V.



## DIVISION.

ART. 114. **I**N multiplication, we have two factors given, and are required to find their product. By multiplying the factors 4 and 6, we obtain the product 24. But it is frequently necessary to reverse this process. The number 24, and *one* of the factors may be given, to enable us to find the other. The operation by which this is effected, is called *Division*. We obtain the number 4, by dividing 24 by 6. The quantity to be divided is called the *dividend* ; the *given* factor, the *divisor* ; and that which is *required*, the *quotient*.

115. DIVISION IS FINDING A QUOTIENT, WHICH MULTIPLIED INTO THE DIVISOR WILL PRODUCE THE DIVIDEND.\*

\* The *remainder* is here supposed to be included in the quotient, as is commonly the case in algebra.

In multiplication, the *multiplier* is always a *number*. (Art. 91.) And the *product* is a quantity of the same kind, as the multiplicand. (Art. 92.) The product of 3 rods into 4, is 12 rods. When we come to division, the product and *either* of the factors may be given, to find the other: that is,

The divisor may be a *number*, and then the quotient will be a quantity of the same kind as the dividend; or

The *divisor* may be a quantity of the same kind as the dividend; and then the *quotient* will be a number.

Thus  $12 \text{ rods} \div 4 = 3 \text{ rods}$ . But  $12 \text{ rods} \div 3 \text{ rods} = 4$   
 And  $12 \text{ rods} \div 24 = \frac{1}{2} \text{ rod}$ . And  $12 \text{ rods} \div 24 \text{ rods} = \frac{1}{2}$

In the first case, the divisor, being a *number*, shows into *how many* parts the dividend is to be separated; and the quotient shows what these parts are.

If 12 rods be divided into 3 parts, each will be 4 rods long. And if 12 rods be divided into 24 parts, each will be *half* a rod long.

In the other case, if the divisor is *less* than the dividend, the former shows into *what* parts the latter is to be divided; and the quotient shows *how many* of these parts are contained in the dividend. In other words, division in this case consists in finding *how often one quantity is contained in another*.

A line of 3 rods, is contained in one of 12 rods, *four times*.

But if the divisor is *greater* than the dividend, and yet a quantity of the same kind, the quotient shows *what part* of the divisor is equal to the dividend.

Thus *one half* of 24 rods is equal to 12 rods.

116. As the product of the divisor and quotient is equal to the dividend, the quotient may be found, by resolving the dividend into two such factors, that one of them shall be the divisor. The other will of course, be the quotient.

Suppose  $abd$  is to be divided by  $a$ . The factors  $a$  and  $bd$  will produce the dividend. The first of these, being a divisor, may be set aside. The other is the quotient. Hence,

WHEN THE DIVISOR IS FOUND AS A FACTOR, IN THE DIVIDEND, THE DIVISION IS PERFORMED, BY CANCELLING THIS FACTOR.

Divide	$cx$	$dh$	$drx$	$hmy$	$dhxy$	$abcd$	$abxy$
By	$c$	$d$	$dr$	$hm$	$dy$	$b$	$ax$
Quot.	$x$		$x$		$hx$		$by$

In each of these examples, the letters which are common to the divisor and dividend, are set aside, and the other letters form the quotient. It will be seen at once, that the product of the quotient and divisor is equal to the dividend.

117. If a letter is *repeated* in the dividend, care must be taken that the factor rejected be only equal to the divisor.

Div.	$aab$	$bbx$	$aadddx$	$aammyy$	$aaaxxxh$	$yyy$
By	$a$	$b$	$ad$	$amy$	$auxx$	$yy$
Quot.	$ab$	<del><math>bx</math></del>	$addx$	<del><math>ammy</math></del>	$axh$	<del><math>yy</math></del>

In such instances, it is obvious that we are not to reject *every* letter in the dividend which is the same with one in the divisor.

118. If the dividend consists of *any factors whatever*, expunging one of them is dividing by it.

Div.	$a(b+d)$	$a(b+d)$	$(b+x)(c+d)$	$(b+y) \times (d-h)x$
By	$a$	$b+d$	$b+x$	$d-h$
Quot.	$b+d$	$a$	$c+d$	$(b+y) \times x$

In all these instances the product of the quotient and divisor is equal to the dividend by Art. 111.

119. In performing multiplication, if the factors contain *numeral figures*, these are multiplied into each other. (Art. 96.) Thus  $3a$  into  $7b$  is  $21ab$ . Now if this process is to be *reversed*, it is evident that dividing the number in the product, by the number in one of the factors, will give the number in the other factor. The quotient of  $21ab \div 3a$  is  $7b$ . Hence,

In division, if there are *numeral co-efficients* prefixed to the letters, the *co-efficient of the dividend must be divided, by the co-efficient of the divisor*.

Div.	$6ab$	$16dxy$	$25dhr$	$12xy$	$34drx$	$20hm$
By	$2b$	$4dx$	$dh$	$6$	$34$	$m$
Quot.	$3a$	<del><math>4x</math></del>	$25r$	<del><math>2xy</math></del>	$drx$	<del><math>hm</math></del>

120. When a simple factor is multiplied into a *compound* one, the former enters into *every* term of the latter. (Art. 98.) Thus  $a$  into  $b+d$ , is  $ab+ad$ . Such a product is easily resolved again into its original factors.

Thus  $ab+ad=a \times (b+d)$

$$ab+ac+ah=a \times (b+c+h)$$

$$amh+amx+amy=am \times (h+x+y)$$

$$4ad+8ah+12am+4ay=4a \times (d+2h+3m+y)$$

Now if the whole quantity be divided by one of these factors, according to Art. 118, the quotient will be the other factor.

Thus  $(cb+ad) \div a = b+d$ . And  $(ab+ad) \div (b+d) = a$ .  
Hence,

If the divisor is contained in every term of a compound dividend, it must be cancelled in each.

Div.	$ab+ac$	$bdh+bdy$	$aah+ay$	$drx+dhx+dxy$
By	$a$	$bd$	$a$	$dx$
Quot.	$b+c$	$h+y$	$ah+y$	$d+h+y$

And if there are *co-efficients*, these must be divided, in each term also.

Div.	$6ab+12ac$	$10dry+16d$	$12hx+8$	$35dm+14dx$
By	$3a$	$2d$	$4$	$7d$
Quot.	$2b+4c$	$5ry+8$	$3hx+2$	$5m+2x$

121. On the other hand, if a compound expression contain any factor in every term, be divided by the other quantities connected by their signs, the quotient will be that factor. See the first part of the preceding article.

Div.	$ab+ac+ah$	$amh+amx+amy$	$4ab+8ay$	$ahm+ahy$
By	$b+c+h$	$h+x+y$	$b+2y$	$m+y$
Quot.	$a$		$4a$	

122. In division, as well as in multiplication, the caution must be observed, not to confound *terms* with *factors*. See Art. 99.

Thus  $(ab+ac) \div a = b+c$ . (Art. 120.)

But  $(ab \times ac) \div a = aabc \div a = abc$ .

And  $(ab+ac) \div (b+c) = a$ . (Art. 121.)

But  $(ab \times ac) \div (b \times c) = aabc \div bc = aa$ .

123. IN DIVISION, THE SAME RULE IS TO BE OBSERVED RESPECTING THE SIGNS, AS IN MULTIPLICATION; THAT IS, IF THE DIVISOR AND DIVIDEND ARE BOTH POSITIVE, OR BOTH NEGATIVE, THE QUOTIENT MUST BE POSITIVE: IF ONE IS POSITIVE AND THE OTHER NEGATIVE, THE QUOTIENT MUST BE NEGATIVE. (Art. 105.)

This is manifest from the consideration that the product of the divisor and quotient must be the same as the dividend.

$$\left. \begin{array}{l} \text{If } +a \times +b = +ab \\ \quad -a \times +b = -ab \\ \quad +a \times -b = -ab \\ \quad -a \times -b = +ab \end{array} \right\} \text{ then } \left\{ \begin{array}{l} +ab \div +b = +a \\ -ab \div +b = -a \\ -ab \div -b = +a \\ +ab \div -b = -a \end{array} \right.$$

Div.	$abx$	$8a - 10ay$	$3ax - 6ay$	$6am \times dh$
By	$-a$	$-2a$	$3a$	$-2a$
	$-bx$	$-4 + 5y$	<del><math>x - 2y</math></del>	$-3m \times dh = -3hdm$

124. IF THE LETTERS OF THE DIVISOR ARE NOT TO BE FOUND IN THE DIVIDEND, THE DIVISION IS EXPRESSED BY WRITING THE DIVISOR UNDER THE DIVIDEND, IN THE FORM OF A VULGAR FRACTION.

Thus  $xy \div a = \frac{xy}{a}$ ; and  $(d-x) \div -h = \frac{d-x}{h}$ .

This is a method of *denoting* division, rather than an actual performing of the operation. But the *purposes* of division may frequently be answered, by these fractional expressions. As they are of the same nature with other vulgar fractions, they may be added, subtracted, multiplied, &c. See the next Section.

125. When the dividend is a compound quantity, the divisor may either be placed under the *whole* dividend, as in the preceding instances, or it may be repeated under *each term*, taken separately. There are occasions when it will be convenient to exchange one of these forms of expression for the other.

Thus  $b+c$  divided by  $x$ , is either  $\frac{b+c}{x}$ , or  $\frac{b}{x} + \frac{c}{x}$ .

And  $a+b$  divided by 2, is either  $\frac{a+b}{2}$  that is, half the

sum of  $a$  and  $b$ ; or  $\frac{a}{2} + \frac{b}{2}$  that is, the sum of half  $a$  and half  $b$ . For it is evident that *half the sum* of two or more quantities, is equal to the *sum of their halves*. And the same principle is applicable, to a third, fourth, fifth, or any other portion of the dividend.

So also  $a-b$  divided by 2, is either  $\frac{a-b}{2}$ , or  $\frac{a}{2} - \frac{b}{2}$ .

For *half the difference* of two quantities, is equal to the *difference of their halves*.

So  $\frac{a-2b+h}{m} = \frac{a}{m} - \frac{2b}{m} + \frac{h}{m}$ . And  $\frac{3a-c}{-x} = \frac{3a}{-x} - \frac{c}{-x}$ .

126. If *some* of the letters in the divisor are in each term of the dividend, the fractional expression may be rendered more simple, by rejecting equal factors from the numerator and denominator.

Div.	$\frac{ab}{ac}$	$\frac{dhx}{dy}$	$\frac{ahm-3ay}{ab}$	$\frac{ab+bx}{by}$	$\frac{2am}{2xy}$
By	$\frac{ab}{ac}$	$\frac{b}{c}$	$\frac{hm-3y}{b}$	$\frac{am}{xy}$	$\frac{am}{xy}$
Quot.	$\frac{ab}{ac}$ or $\frac{b}{c}$				

These reductions are made upon the principle, that a given divisor is contained in a given dividend, just as many times, as double the divisor in double the dividend; triple the divisor in triple the dividend, &c. See the reduction of fractions.

127. If the divisor is in some of the terms of the dividend, but not in all; those which contain the divisor may be divided as in Art. 116, and the others set down in the form of a fraction.

Thus  $(ab+d) \div a$  is either  $\frac{ab+d}{a}$ , or  $\frac{ab}{a} + \frac{d}{a}$  or  $b + \frac{d}{a}$ .

Div.	$\frac{dxy+rx-hd}{x}$	$\frac{2ah+ad+x}{a}$	$\frac{bm+3y}{-b}$	$\frac{2my+dh}{2m}$
By	$\frac{dxy+rx-hd}{x}$	$\frac{2ah+ad+x}{a}$	$\frac{bm+3y}{-b}$	$\frac{2my+dh}{2m}$
Quot.	$dy+r-\frac{hd}{x}$		$-m+\frac{3y}{-b}$	



128. The quotient of any quantity divided by *itself* or its *equal*, is obviously *a unit*.

Thus  $\frac{a}{a} = 1$ . And  $\frac{3ax}{3ax} = 1$ . And  $\frac{6}{4+2} = 1$ . And  $\frac{a+b-3h}{a+b-3h} = 1$ .

Div.	$ax+x$	$3bd-3d$	$4axy-4a+8ad$	$3ab+3-6m$
By	$x$	$3d$	$4a$	$3$
Quot.	$a+1$	$b-1$	$xy-1+2d$	$ab+2m$

Cor. If the dividend is *greater* than the divisor, the quotient must be *greater than a unit*: But if the dividend is *less* than the divisor, the quotient must be *less than a unit*.

PROMISCUOUS EXAMPLES.

1. Divide  $12aby+6abx-18bbm+24b$ , by  $6b$ .
2. Divide  $16a-12+8y+4-20adx+m$ , by  $4$ .
3. Divide  $(a-2h) \times (3m+y) \times x$ , by  $(a-2h) \times (3m+y)$ .
4. Divide  $ahd-4ad+3ay-a$ , by  $hd-4d+3y-1$ .
5. Divide  $ax-ry+ad-4my-6+a$ , by  $-a$ .
6. Divide  $amy+3my-mxy+am-d$ , by  $-dmy$ .
7. Divide  $ard-6a+2r-hd+6$ , by  $2ard$ .
8. Divide  $6ax-8+2xy+4-6hy$ , by  $4axy$ .

129. From the nature of division it is evident, that the value of the quotient depends both on the divisor and the dividend. With a given divisor, the greater the dividend, the greater the quotient. And with a given dividend, the greater the divisor, the less the quotient. In several of the succeeding parts of algebra, particularly the subjects of fractions, ratios, and proportion, it will be important to be able to determine what change will be produced in the quotient, by increasing or diminishing either the divisor or the dividend.

If the given dividend be 24, and the divisor 6; the quotient will be 4. But this same dividend may be supposed to be multiplied or divided by some *other* number, before it is divided by 6. Or the *divisor* may be multiplied or divided by some other number, before it is used in dividing 24. In each of these cases, the quotient will be altered.

130. In the first place, if the given divisor is contained in the given dividend a certain number of times, it is obvious that the same divisor is contained,

In *double* that dividend, *twice* as many times;

In *triple* the dividend, *thrice* as many times, &c.

That is, if the divisor remains the same, *multiplying the dividend* by any quantity, is, in effect, *multiplying the quotient* by that quantity.

Thus, if the constant divisor is 6, then  $24 \div 6 = 4$  the quotient.

Multiplying the dividend by 2,  $2 \times 24 \div 6 = 2 \times 4$

Multiplying by any number  $n$   $n \times 24 \div 6 = n \times 4$

131. Secondly, if the given divisor is contained in the given dividend a certain number of times, the same divisor is contained,

In *half* that dividend, half as many times ;

In *one third* of the dividend, one third as many times, &c.

That is, if the divisor remains the same, *dividing the dividend* by any other quantity, is, in effect, *dividing the quotient* by that quantity.

Thus

$$24 \div 6 = 4$$

Dividing the dividend by 2,

$$\frac{1}{2} 24 \div 6 = \frac{1}{2} 4$$

Dividing by  $n$ ,

$$\frac{1}{n} 24 \div 6 = \frac{1}{n} 4.$$

132. Thirdly, if the given divisor is contained in the given dividend a certain number of times, then, in the same dividend,

*Twice* that divisor is contained only *half* as many times ;

*Three times* the divisor is contained, *one third* as many times.

That is, if the dividend remains the same, *multiplying the divisor* by any quantity, is, in effect, *dividing the quotient* by that quantity.

Thus

$$24 \div 6 = 4$$

Multiplying the divisor by 2,

$$24 \div 2 \times 6 = \frac{4}{2}$$

Multiplying by  $n$

$$24 \div n \times 6 = \frac{4}{n}.$$

133. Lastly, if the given divisor is contained in the given dividend a certain number of times, then, in the same dividend,

*Half* that divisor is contained, *twice* as many times.

*One third* of the divisor is contained *thrice* as many times ;

That is, if the dividend remains the same, *dividing the divisor* by any other quantity, is, in effect, *multiplying the quotient* by that quantity.

Thus

$$24 \div 6 = 4$$

Dividing the divisor by 2,

$$24 \div \frac{1}{2} 6 = 2 \times 4$$

Dividing by  $n$ ,

$$24 \div \frac{1}{n} 6 = n \times 4$$

For the method of performing division, when the divisor and dividend are *both compound quantities*, see one of the following sections.

## SECTION V.




---

 FRACTIONS.\*

ART. 134. **E**XPRESSIONS in the form of fractions occur more frequently in algebra than in arithmetic. Most instances in division belong to this class. Indeed the numerator of *every* fraction may be considered as a *dividend*, of which the denominator is a *divisor*.

According to the common definition in arithmetic, the denominator shows into what parts an integral unit is supposed to be divided; and the numerator shows how many of these parts belong to the fraction. But it makes no difference, whether the *whole* of the numerator is divided by the denominator; or only *one* of the integral units is divided, and then the quotient taken as many times, as the number of units in the numerator. Thus  $\frac{3}{4}$  is the same as  $\frac{1}{4} + \frac{1}{4} + \frac{1}{4}$ . A fourth part of *three* dollars, is equal to three fourths of *one* dollar.

135. The *value* of a fraction, is the *quotient* of the numerator divided by the denominator.

Thus the value of  $\frac{6}{2}$  is 3. The value of  $\frac{ab}{b}$  is  $a$ .

From this it is evident, that whatever changes are made in the *terms* of a fraction; if the *quotient* is not altered, the value remains the same. For any fraction, therefore, we may substitute any *other* fraction which will give the same quotient.

Thus  $\frac{4}{2} = \frac{10}{5} = \frac{4ba}{2ba} = \frac{3drx}{4drx} = \frac{6+2}{3+1}$  &c. For the quotient in each of these instances is 2.

136. As the value of a fraction is the quotient of the numerator divided by the denominator, it is evident, from Art. 128, that when the numerator is *equal* to the denominator, the value of the fraction is a *unit*; when the numerator is

\* Horsley's Mathematics, Camus' Arithmetic, Emerson, Euler, Saunderson, and Ludlam.

less than the denominator, the value is *less than a unit*; and when the numerator is *greater* than the denominator, the value is *greater than a unit*.

The calculations in fractions depend on a few general principles, which will here be stated in connection with each other.

137. *If the denominator of a fraction remains the same, multiplying the NUMERATOR by any quantity, is multiplying the VALUE by that quantity; and dividing the numerator, is dividing the value.* For the numerator and denominator are a dividend and divisor, of which the value of the fraction is the quotient. And by Art. 130 and 131, multiplying the dividend is in effect multiplying the quotient, and dividing the dividend is dividing the quotient.

Thus, in the fractions  $\frac{ab}{a}$ ,  $\frac{3ab}{a}$ ,  $\frac{7abd}{a}$ ,  $\frac{\frac{1}{3}ab}{a}$ , &c.

The quotients or values are  $b$ ,  $3b$ ,  $7bd$ ,  $\frac{1}{3}b$ , &c.

Here it will be seen that, while the denominator is not altered, the value of the fraction is multiplied or divided by the same quantity as the numerator.

Cor. With a given denominator, the greater the numerator, the greater will be the *value* of the fraction; and, on the other hand, the greater the value, the greater the numerator.

138. *If the numerator remains the same, multiplying the denominator by any quantity, is dividing the value by that quantity; and dividing the denominator, is multiplying the value.* For multiplying the divisor is dividing the quotient; and dividing the divisor is multiplying the quotient. (Art. 132, 133.)

In the fractions  $\frac{24ab}{6b}$ ,  $\frac{24ab}{12b}$ ,  $\frac{24ab}{3b}$ ,  $\frac{24ab}{b}$ , &c.

The values are  $4a$ ,  $2a$ ,  $8a$ ,  $24a$ , &c.

Cor. With a given numerator, the *greater* the denominator, the *less* will be the value of the fraction; and the *less* the value, the *greater* the denominator.

139. From the two last articles it follows, that *dividing the numerator* by any quantity, will have the same effect on the value of the fraction, as *multiplying the denominator* by that quantity; and *multiplying the numerator* will have the same effect, as *dividing the denominator*.

140. It is also evident, from the preceding articles, that IF THE NUMERATOR AND DENOMINATOR BE BOTH MULTIPLIED, OR BOTH DIVIDED, BY THE SAME QUANTITY, THE VALUE OF THE FRACTION WILL NOT BE ALTERED.

Thus  $\frac{bx}{b} = \frac{abx}{ab} = \frac{3bx}{3b} = \frac{\frac{1}{2}bx}{\frac{1}{2}b} = \frac{\frac{1}{3}abx}{\frac{1}{3}ab}$  &c. For in each of these instances the quotient is  $x$ .

141. Any integral quantity may, without altering its value, be thrown into the form of a fraction, by multiplying the quantity into the proposed denominator, and taking the product for a numerator.

Thus  $a = \frac{a}{1} = \frac{ab}{b} = \frac{ad+ah}{d+h} = \frac{6adh}{6dh}$ , &c. For the quotient of each of these is  $a$ .

So  $d+h = \frac{dx+hx}{x}$ . And  $r+1 = \frac{2drr+2dr}{2dr}$ .

142. There is nothing perhaps, in the calculation of algebraic fractions, which occasions more perplexity to a learner, than the positive and negative signs. The changes in these are so frequent, that it is necessary to become familiar with the principles on which they are made. The use of the sign which is prefixed to the dividing line, is to show whether the value of the *whole fraction* is to be added to, or subtracted from, the other quantities with which it is connected. (Art. 43.) This sign, therefore, has an influence on the several terms taken collectively. But in the numerator and denominator, each sign affects only the single term to which it is applied.

The value of  $\frac{ab}{b}$  is  $a$ . (Art. 135.) But this will become negative, if the sign  $-$  be prefixed to the fraction.

Thus  $y + \frac{ab}{b} = y + a$ . But  $y - \frac{ab}{b} = y - a$ .

So that changing the sign which is before the whole fraction, has the effect of changing the *value* from positive to negative, or from negative to positive.

Next, suppose the sign or signs of the *numerator* to be changed.

By Art. 123,  $\frac{ab}{b} = +a$ . But  $\frac{-ab}{b} = -a$ .

$$\text{And } \frac{ab-bc}{b} = +a-c. \quad \text{But } \frac{-ab+bc}{b} = -a+c.$$

That is, by changing all the signs of the numerator, the value of the fraction is changed from positive to negative, or the contrary.

Again, suppose the sign of the *denominator* to be changed.

$$\text{As before } \frac{ab}{b} = +a. \quad \text{But } \frac{ab}{-b} = -a.$$

143. We have, then, this general proposition; IF THE SIGN PREFIXED TO A FRACTION, OR ALL THE SIGNS OF THE NUMERATOR, OR ALL THE SIGNS OF THE DENOMINATOR BE CHANGED; THE VALUE OF THE FRACTION WILL BE CHANGED, FROM POSITIVE TO NEGATIVE, OR FROM NEGATIVE TO POSITIVE.

From this is derived another important principle. As each of the changes mentioned here is from positive to negative, or the contrary; if any *two* of them be made at the same time, *they will balance each other.*

Thus, by changing the sign of the numerator,

$$\frac{ab}{b} = +a \text{ becomes } \frac{-ab}{b} = -a.$$

But, by changing both the numerator and denominator, it becomes  $\frac{-ab}{-b} = +a$ , where the positive value is restored.

By changing the sign before the fraction,

$$y + \frac{ab}{b} = y + a \text{ becomes } y - \frac{ab}{b} = y - a.$$

But, by changing the sign of the numerator also, it becomes  $y - \frac{-ab}{b}$  where the quotient  $-a$  is to be *subtracted* from  $y$ , or which is the same thing, (Art. 81,)  $+a$  is to be *added*, making  $y + a$  as at first. Hence,

144. IF ALL THE SIGNS BOTH OF THE NUMERATOR AND DENOMINATOR, OR THE SIGNS OF ONE OF THESE WITH THE SIGN PREFIXED TO THE WHOLE FRACTION, BE CHANGED AT THE SAME TIME, THE VALUE OF THE FRACTION WILL NOT BE ALTERED.

$$\text{Thus } \frac{6}{2} = \frac{-6}{-2} = -\frac{-6}{2} = -\frac{6}{-2} = +3.$$

$$\text{And } \frac{6}{-2} = \frac{-6}{2} = -\frac{6}{2} = -\frac{-6}{-2} = -3.$$

Hence the quotient in division may be set down in different ways. Thus  $(a-c) \div b$ , is either  $\frac{a}{b} + \frac{-c}{b}$ , or  $\frac{a}{b} - \frac{c}{b}$ .

The latter method is the most common. See the examples in Art. 127.

### REDUCTION OF FRACTIONS.

145. From the principles which have been stated, are derived the rules for the *Reduction* of fractions, which are substantially the same in algebra, as in arithmetic.

A FRACTION MAY BE REDUCED TO LOWER TERMS, BY DIVIDING BOTH THE NUMERATOR AND DENOMINATOR, BY ANY QUANTITY WHICH WILL DIVIDE THEM WITHOUT A REMAINDER. According to Art. 140, this will not alter the *value* of the fraction.

$$\text{Thus } \frac{ab}{cb} = \frac{a}{c}. \quad \text{And } \frac{6dm}{3dy} = \frac{3m}{4y}. \quad \text{And } \frac{7m}{7mr} = \frac{1}{r}.$$

In the last example, both parts of the fraction are divided by the numerator.

$$\text{Again } \frac{a+bc}{(a+bc) \times m} = \frac{1}{m} \quad (\text{Art. 118.}) \quad \text{And } \frac{am+ay}{bm+by} = \frac{a}{b}.$$

If a letter is in *every* term both of the numerator and denominator, it may be *cancelled*, for this is *dividing* by that letter. (Art. 120.)

$$\text{Thus } \frac{3am+ay}{ad+ah} = \frac{3m+y}{d+h} \quad \text{And } \frac{dry+dy}{dhy-dy} = \frac{r+1}{h-1}.$$

If the numerator and denominator be divided by the *greatest common measure*, it is evident that the fraction will be reduced to the *lowest* terms. For the method of finding the greatest common measure, see Sec. xvi.

146. FRACTIONS OF DIFFERENT DENOMINATORS MAY BE REDUCED TO A COMMON DENOMINATOR, BY MULTIPLYING EACH NUMERATOR INTO ALL THE DENOMINATORS EXCEPT ITS OWN,

FOR A NEW NUMERATOR; AND ALL THE DENOMINATORS TOGETHER, FOR A COMMON DENOMINATOR.

Ex. 1. Reduce  $\frac{a}{b}$ , and  $\frac{c}{d}$ , and  $\frac{m}{y}$  to a common denominator.

$$\left. \begin{array}{l} a \times d \times y = ady \\ c \times b \times y = cby \\ m \times b \times d = mbd \\ b \times d \times y = bdy \end{array} \right\} \begin{array}{l} \text{the three numerators.} \\ \text{the common denominator.} \end{array}$$

The fractions reduced are  $\frac{ady}{bdy}$ , and  $\frac{bcy}{bdy}$ , and  $\frac{bdm}{bdy}$ .

Here it will be seen, that the reduction consists in multiplying the numerator and denominator of each fraction, into all the other denominators. This does not alter the value. (Art. 140.)

2. Reduce  $\frac{dr}{3m}$ , and  $\frac{2h}{g}$ , and  $\frac{6c}{y}$ .

3. Reduce  $\frac{2}{3}$ , and  $\frac{a}{x}$ , and  $\frac{r+1}{d+h}$ .

4. Reduce  $\frac{1}{a+b}$ , and  $\frac{1}{a-b}$ .

After the fractions have been reduced to a common denominator, they may be brought to lower terms, by the rule in the last article, if there is any quantity, which will divide the denominator, and *all* the numerators, without a remainder.

An *integer* and a fraction are easily reduced to a common denominator. (Art. 141.)

Thus  $a$  and  $\frac{b}{c}$  are equal to  $\frac{a}{1}$  and  $\frac{b}{c}$ , or  $\frac{ac}{c}$  and  $\frac{b}{c}$ .

And  $a, b, \frac{h}{m}, \frac{d}{y}$  are equal to  $\frac{amy}{my}, \frac{bmy}{my}, \frac{hy}{my}, \frac{dm}{my}$ .

147. TO REDUCE AN IMPROPER FRACTION TO A MIXED QUANTITY, DIVIDE THE NUMERATOR BY THE DENOMINATOR, as in Art. 127.

Thus  $\frac{ab+bm+d}{b} = a + m + \frac{d}{b}$ .



Reduce  $\frac{am - a + ady - hr}{a}$ , to a mixed quantity.

For the reduction of a *mixed quantity* to an improper fraction, see Art. 150. And for the reduction of a *compound fraction* to a simple one, see Art. 160.

### ADDITION OF FRACTIONS.

148. In adding fractions, we may either write them one after the other, with their signs, as in the addition of integers, or we may incorporate them into a single fraction, by the following rule :

REDUCE THE FRACTIONS TO A COMMON DENOMINATOR, MAKE THE SIGNS BEFORE THEM ALL POSITIVE, AND THEN ADD THEIR NUMERATORS.

The common denominator shows into what parts the integral unit is supposed to be divided ; and the numerators show the *number* of these parts belonging to each of the fractions. (Art. 134.) Therefore the numerators *taken together* show the whole number of parts in all the fractions.

$$\text{Thus } \frac{2}{7} = \frac{1}{7} + \frac{1}{7} \quad \text{And } \frac{3}{7} = \frac{1}{7} + \frac{1}{7} + \frac{1}{7},$$

$$\text{Therefore } \frac{2}{7} + \frac{3}{7} = \frac{1}{7} + \frac{1}{7} + \frac{1}{7} + \frac{1}{7} + \frac{1}{7} = \frac{5}{7}.$$

The numerators are added, according to the rules for the addition of integers. (Art. 69, &c.) It is obvious that the sum is to be placed over the common denominator. To avoid the perplexity which might be occasioned by the signs, it will be expedient to make those *prefixed* to the fractions uniformly positive. But in doing this, care must be taken not to alter the value. This will be preserved, if all the signs in the numerator, are changed at the same time with that before the fraction. (Art. 144.)

$$\text{Ex. 1. Add } \frac{2}{16} \text{ and } \frac{4}{16} \text{ of a pound. Ans. } \frac{2+4}{16} \text{ or } \frac{6}{16}.$$

It is as evident that  $\frac{2}{16}$ , and  $\frac{4}{16}$  of a pound, are  $\frac{6}{16}$  of a pound, as that 2 ounces, and 4 ounces, are 6 ounces.

$$2. \text{ Add } \frac{a}{b} \text{ and } \frac{c}{d}. \text{ First reduce them to a common denom-}$$

inator. They will then be  $\frac{ad}{bd}$  and  $\frac{bc}{bd}$ , and their sum  $\frac{ad+bc}{bd}$ .

3. Given  $\frac{m}{d}$  and  $-\frac{2r+d}{3h}$ , to find their sum.

$$\text{Ans. } \frac{m}{d} \text{ and } -\frac{2r+d}{3h} = \frac{3hm}{3dh} \text{ and } -\frac{2dr+dd}{3dh} = \frac{3hm-2dr-dd}{3dh}$$

$$4. \frac{a}{d} \text{ and } -\frac{b-m}{y} = \frac{a}{d} + \frac{-b+m}{y} = \frac{ay-bd+dm}{dy}$$

$$5. \frac{a}{y} \text{ and } \frac{d}{-m} = \frac{-am}{-my} + \frac{dy}{-my} = \frac{-am+dy}{-my} \text{ or } \frac{am-dy}{my}$$

$$6. \frac{a}{a+b} \text{ and } \frac{b}{a-b} = \frac{aa-ab+ab+bb}{aa+ab-ab-bb} = \frac{aa+bb}{aa-bb}. \quad (\text{Art. 77.})$$

$$7. \text{ Add } \frac{-a}{d} \text{ to } \frac{-h}{m-r}. \quad 8. \text{ Add } \frac{-4}{2} \text{ to } \frac{-16}{7-3}. \quad \text{Ans. } -6.$$

149. For many purposes, it is sufficient to add fractions in the same manner as integers are added, by writing them one after another with their signs. (Art. 69.)

$$\text{Thus the sum of } \frac{a}{b} \text{ and } \frac{3}{y} \text{ and } -\frac{d}{2m}, \text{ is } \frac{a}{b} + \frac{3}{y} - \frac{d}{2m}.$$

In the same manner, fractions and integers may be added.

$$\text{The sum of } a \text{ and } \frac{d}{y} \text{ and } 3m \text{ and } -\frac{h}{r}, \text{ is } a+3m+\frac{d}{y}-\frac{h}{r}.$$

150. Or the integer may be *incorporated* with the fraction, by converting the former into a fraction, and then adding the numerators. See Art. 141.

$$\text{The sum of } a \text{ and } \frac{b}{m}, \text{ is } \frac{a}{1} + \frac{b}{m} = \frac{am}{m} + \frac{b}{m} = \frac{am+b}{m}$$

$$\text{The sum of } 3d \text{ and } \frac{h+d}{m-y}, \text{ is } \frac{3dm-3dy+h+d}{m-y}$$

Incorporating an integer with a fraction, is the same as *reducing a mixed quantity* to an improper fraction. For a mixed quantity is an integer and a fraction. In arithmetic, these are generally placed together, without any sign be-

tween them. But in algebra, they are distinct terms. Thus  $2\frac{1}{3}$  is 2 and  $\frac{1}{3}$ , which is the same as  $2+\frac{1}{3}$ .

Ex. 1. Reduce  $a+\frac{1}{b}$  to an improper fraction. Ans.  $\frac{ab+1}{b}$ .

2. Reduce  $m+d-\frac{r}{h-d}$ . Ans.  $\frac{hm-dm+dh-dd-r}{h-d}$ .

3. Reduce  $1+\frac{d}{b}$ . Ans.  $\frac{b+d}{b}$ . 4. Reduce  $1-\frac{h}{m}$ .

5. Reduce  $b+\frac{c}{d-y}$ . 6. Reduce  $3+\frac{2d-4}{3a}$ .

SUBTRACTION OF FRACTIONS.

151. The methods of performing subtraction in algebra, depend on the principle, that adding a negative quantity is equivalent to subtracting a positive one; and *v. v.* (Art. 81.) For the subtraction of fractions, then, we have the following simple rule. CHANGE THE FRACTION TO BE SUBTRACTED, FROM POSITIVE TO NEGATIVE, OR THE CONTRARY, AND THEN PROCEED AS IN ADDITION. (Art. 148.) In making the required change, it will be expedient to alter, in some instances, the signs of the numerator, and in others, the sign before the dividing line, (Art. 143,) so as to leave the latter always affirmative.

Ex. 1. From  $\frac{a}{b}$ , subtract  $\frac{h}{m}$ .

First change  $\frac{h}{m}$ , the fraction to be subtracted, to  $\frac{-h}{m}$ .

Secondly, reduce the two fractions to a common denominator, making  $\frac{am}{bm}$  and  $\frac{-bh}{bm}$ .

Thirdly, the sum of the numerators  $am-bh$ , placed over the common denominator, gives the answer,  $\frac{am-bh}{bm}$ .

2. From  $\frac{a+y}{r}$ , subtract  $\frac{h}{d}$ . Ans.  $\frac{ad+dy-hr}{dr}$ .

3. From  $\frac{a}{m}$  subtract  $\frac{d-b}{y}$ . Ans.  $\frac{ay-dm+bm}{my}$ .

4. From  $\frac{a+3d}{4}$ , subtract  $\frac{3a-2d}{3}$ . Ans.  $\frac{17d-9a}{12}$ .

5. From  $\frac{b-d}{m}$  subtract  $-\frac{b}{y}$ . Ans.  $\frac{by-dy+bm}{my}$ .

6. From  $\frac{a+1}{d}$  subtract  $\frac{d-1}{m}$ . 7. From  $\frac{3}{a}$  subtract  $\frac{4}{b}$ .

152. Fractions may also be subtracted, like integers, by setting them down, after their signs are changed, without reducing them to a common denominator.

From  $\frac{h}{m}$  subtract  $-\frac{h+d}{y}$ . Ans.  $\frac{h}{m} + \frac{h+d}{y}$ .

In the same manner, an integer may be subtracted from a fraction, or a fraction from an integer.

From  $a$  subtract  $\frac{b}{m}$ . Ans.  $a - \frac{b}{m}$ .

153. Or the integer may be incorporated with the fraction, as in Art. 150.

Ex. 1. From  $\frac{h}{y}$  subtract  $m$ . Ans.  $\frac{h}{y} - m = \frac{h-my}{y}$ .

2. From  $4a + \frac{b}{c}$  subtract  $3a - \frac{h}{d}$ . Ans.  $\frac{acd+bd+hc}{cd}$ .

3. From  $1 + \frac{b-c}{d}$  subtract  $\frac{c-b}{d}$ . Ans.  $\frac{d+2b-2c}{d}$ .

3. From  $a+3h - \frac{d-b}{2}$  subtract  $3a - h + \frac{d+b}{3}$ .

### MULTIPLICATION OF FRACTIONS.

154. By the definition of multiplication, multiplying by a fraction is taking a *part* of the multiplicand, as many times, as there are like parts of an unit in the multiplier. (Art. 90.) Now the denominator of a fraction shows into what parts the integral unit is supposed to be divided; and the numerator shows how many of those parts belong to the given fraction. In multiplying by a fraction, therefore, the multiplicand is to be divided into such parts, as are denoted by the denominator; and then one of these parts is to be repeated, as many times, as is required by the numerator.

Suppose  $a$  is to be multiplied by  $\frac{3}{4}$ .

A fourth part of  $a$  is  $\frac{a}{4}$ .

This taken 3 times is  $\frac{a}{4} + \frac{a}{4} + \frac{a}{4} = \frac{3a}{4}$ . (Art. 148.)

Again, suppose  $\frac{a}{b}$  is to be multiplied by  $\frac{3}{4}$ .

One fourth of  $\frac{a}{b}$  is  $\frac{a}{4b}$ . (Art. 138.)

This taken 3 times is  $\frac{a}{4b} + \frac{a}{4b} + \frac{a}{4b} = \frac{3a}{4b}$

the product required.

In a similar manner, any fractional multiplicand may be divided into parts, by multiplying the denominator; and one of the parts may be repeated, by multiplying the numerator. We have then the following rule:

155. TO MULTIPLY FRACTIONS, MULTIPLY THE NUMERATORS TOGETHER, FOR A NEW NUMERATOR, AND THE DENOMINATORS TOGETHER, FOR A NEW DENOMINATOR.

Ex. 1. Multiply  $\frac{3b}{c}$  into  $\frac{d}{2m}$ . Product  $\frac{3bd}{2cm}$ .

2. Multiply  $\frac{a+d}{y}$  into  $\frac{4h}{m-2}$ . Product  $\frac{4ah+4dh}{my-2y}$ .

3. Multiply  $\frac{(a+m) \times h}{3}$  into  $\frac{4}{(a-n)}$  Product  $\frac{(a+m) \times 4h}{3 \times (a-n)}$ .

4. Mult.  $\frac{a+h}{3+y}$  into  $\frac{4-m}{c+y}$ . 5. Mult.  $\frac{1}{a+3r}$  into  $\frac{3}{8}$ .

156. The method of multiplying is the same, when there are more than two fractions to be multiplied together.

1. Multiply together  $\frac{a}{b}$ ,  $\frac{c}{d}$ , and  $\frac{m}{y}$ . Product  $\frac{acm}{bdy}$ .

For  $\frac{a}{b} \times \frac{c}{d}$  is, by that article  $\frac{ac}{bd}$ , and this into  $\frac{m}{y}$  is  $\frac{acm}{bdy}$ .

2. Multiply  $\frac{2a}{m}$ ,  $\frac{h}{y}$ , and  $\frac{1}{r-1}$ . Product  $\frac{2abh-2abd}{cmry-cmy}$ .

3. Mult.  $\frac{3+b}{n}$ ,  $\frac{1}{h}$  and  $\frac{d}{r+2}$ . 4. Mult.  $\frac{ad}{hy}$ ,  $\frac{a-6}{d+1}$ , and  $\frac{3}{7}$ .

157. The multiplication may sometimes be shortened, by rejecting equal factors, from the numerators and denominators.

1. Multiply  $\frac{a}{r}$  into  $\frac{h}{a}$  and  $\frac{d}{y}$ . Product  $\frac{dh}{ry}$ .

Here  $a$  being in one of the numerators, and in one of the denominators, may be omitted. If it be retained, the product will be  $\frac{adh}{ary}$ . But this reduced to lower terms, by

Art. 145, will become  $\frac{dh}{ry}$  as before.

2. Multiply  $\frac{ad}{m}$  into  $\frac{m}{3a}$  and  $\frac{ah}{2d}$ . Product  $\frac{ah}{6}$ .

It is necessary that the factors rejected from the numerators be exactly equal to those which are rejected from the denominators. In the last example,  $a$  being in two of the numerators, and in only one of the denominators, must be retained in one of the numerators.

3. Multiply  $\frac{a+d}{y}$  into  $\frac{my}{ah}$ . Product  $\frac{am+dm}{ah}$ .

Here, though the same letter  $a$  is in one of the numerators, and in one of the denominators, yet as it is not in every term of the numerator, it must not be cancelled.

4. Multiply  $\frac{am+d}{h}$  into  $\frac{h}{m}$  and  $\frac{3r}{5a}$ .

If any difficulty is found, in making these contractions, it will be better to perform the multiplication, without omitting any of the factors; and to reduce the product to lower terms afterwards.

158. When a fraction and an integer are multiplied together, the numerator of the fraction is multiplied into the integer. The denominator is not altered; except in cases where division of the denominator is substituted for multiplication of the numerator, according to Art. 139.

Thus  $a \times \frac{m}{y} = \frac{am}{y}$ . For  $a = \frac{a}{1}$ ; and  $\frac{a}{1} \times \frac{m}{y} = \frac{am}{y}$ .

So  $r \times \frac{x}{d} \times \frac{h+1}{3} = \frac{hrx+rx}{3d}$ . And  $a \times \frac{1}{b} = \frac{a}{b}$ . Hence,

159. A FRACTION IS MULTIPLIED INTO A QUANTITY EQUAL TO ITS DENOMINATOR, BY CANCELLING THE DENOMINATOR.

Thus  $\frac{a}{b} \times b = a$ . For  $\frac{a}{b} \times b = \frac{ab}{b}$ . But the letter  $b$ , being in both the numerator and denominator, may be set aside. (Art. 145.)

So  $\frac{3m}{a-y} \times (a-y) = 3m$ . And  $\frac{h+3d}{3+m} \times (3+m) = h+3d$ .

On the same principle, a fraction is multiplied into any factor in its denominator, by cancelling that factor.

Thus  $\frac{a}{by} \times y = \frac{ay}{by} = \frac{a}{b}$ . And  $\frac{h}{24} \times 6 = \frac{h}{4}$ .

160. From the definition of multiplication by a fraction, it follows that what is commonly called a *compound fraction*,\*

is the *product* of two or more fractions. Thus  $\frac{3}{4}$  of  $\frac{a}{b}$  is  $\frac{3}{4} \times \frac{a}{b}$ . For  $\frac{3}{4}$  of  $\frac{a}{b}$ , is  $\frac{1}{4}$  of  $\frac{a}{b}$  taken three times, that is,  $\frac{a}{4b} + \frac{a}{4b} + \frac{a}{4b}$ . But this is the same as  $\frac{a}{b}$  multiplied by  $\frac{3}{4}$ . (Art. 154.)

Hence, *reducing a compound fraction to a simple one, is the same, as multiplying fractions into each other.*

Ex. 1. Reduce  $\frac{2}{7}$  of  $\frac{a}{b+2}$ . Ans.  $\frac{2a}{7b+14}$ .

2. Reduce  $\frac{2}{3}$  of  $\frac{4}{5}$  of  $\frac{b+h}{2a-m}$ . Ans.  $\frac{8b+8h}{30a-15m}$ .

3. Reduce  $\frac{1}{7}$  of  $\frac{1}{3}$  of  $\frac{1}{3-d}$ . Ans.  $\frac{1}{168-21d}$ .

\* By a compound fraction is meant a fraction of a fraction, and not a fraction whose numerator or denominator is a compound quantity.

161. The expressions  $\frac{2}{3}a$ ,  $\frac{1}{5}b$ ,  $\frac{4}{7}y$ , &c. are equivalent to  $\frac{2a}{3}$ ,  $\frac{b}{5}$ ,  $\frac{4y}{7}$ . For  $\frac{2}{3}a$  is  $\frac{2}{3}$  of  $a$ , which is equal to  $\frac{2}{3} \times a = \frac{2a}{3}$ . (Art. 158.) So  $\frac{1}{5}b = \frac{1}{5} \times b = \frac{b}{5}$ .

### DIVISION OF FRACTIONS.

162. TO DIVIDE ONE FRACTION BY ANOTHER, INVERT THE DIVISOR, AND THEN PROCEED AS IN MULTIPLICATION. (Art. 155.)

Ex. 1. Divide  $\frac{a}{b}$  by  $\frac{c}{d}$ . Ans.  $\frac{a}{b} \times \frac{d}{c} = \frac{ad}{bc}$ .

To understand the reason of the rule, let it be premised, that the product of any fraction into the same fraction inverted is always a unit.

Thus  $\frac{a}{b} \times \frac{b}{a} = \frac{ab}{ab} = 1$ . And  $\frac{d}{h+y} \times \frac{h+y}{d} = 1$ . (Art. 128.)

But a quantity is not altered by multiplying it by a unit. Therefore if a dividend be multiplied, first into the divisor inverted, and then into the divisor itself, the last product will be equal to the dividend. Now, by the definition, art. 115, "division is finding a quotient, which multiplied into the divisor will produce the dividend." And as the dividend multiplied into the divisor inverted is such a quantity, the quotient is truly found by the rule.

This explanation will probably be best understood, by attending to the examples. In several which follow, the *proof* of the division will be given, by multiplying the quotient into the divisor. This will present, at one view, the dividend multiplied into the inverted divisor, and into the divisor itself.

2. Divide  $\frac{m}{2d}$  by  $\frac{3h}{y}$ . Ans.  $\frac{m}{2d} \times \frac{y}{3h} = \frac{my}{6dh}$ .

Proof.  $\frac{my}{6dh} \times \frac{3h}{y} = \frac{m}{2d}$  the dividend.

3. Divide  $\frac{x+d}{r}$  by  $\frac{5d}{y}$ . Ans.  $\frac{x+d}{r} \times \frac{y}{5d} = \frac{xy+dy}{5dr}$ .

Proof.  $\frac{xy+dy}{5dr} \times \frac{5d}{y} = \frac{x+d}{r}$ .



4. Divide  $\frac{4dh}{x}$  by  $\frac{4hr}{a}$ . Ans.  $\frac{4dh}{x} \times \frac{a}{4hr} = \frac{ad}{rx}$ .

Proof.  $\frac{ad}{rx} \times \frac{4hr}{a} = \frac{4dh}{x}$  the dividend.

5. Divide  $\frac{36d}{5}$  by  $\frac{18h}{10y}$ . Ans.  $\frac{36d}{5} \times \frac{10y}{18h} = \frac{4dy}{h}$ .

6. Divide  $\frac{ab+1}{3y}$  by  $\frac{ab-1}{x}$ . 7. Divide  $\frac{h-my}{4}$  by  $\frac{3}{a+1}$ .

163. When a fraction is divided by an *integer*, the *denominator* of the fraction is multiplied into the integer.

Thus the quotient of  $\frac{a}{b}$  divided by  $m$ , is  $\frac{a}{bm}$ .

For  $m = \frac{m}{1}$ ; and by the last article,  $\frac{a}{b} \div \frac{m}{1} = \frac{a}{b} \times \frac{1}{m} = \frac{a}{bm}$ .

So  $\frac{1}{a-b} \div h = \frac{1}{a-b} \times \frac{1}{h} = \frac{1}{ah-bh}$ . And  $\frac{3}{4} \div 6 = \frac{3}{24} = \frac{1}{8}$ .

In fractions, multiplication is made to perform the office of division; because division in the usual form often leaves a troublesome remainder: but there is no remainder in multiplication. In many cases, there are methods of shortening the operation. But these will be suggested by practice, without the aid of particular rules.

164. By the definition, art. 49, "the *reciprocal* of a quantity, is the quotient arising from dividing a unit by that quantity."

Therefore, the reciprocal of  $\frac{a}{b}$ , is  $1 \div \frac{a}{b} = 1 \times \frac{b}{a} = \frac{b}{a}$ . That is,

*The reciprocal of a fraction is the fraction inverted.*

Thus the reciprocal of  $\frac{b}{m+y}$  is  $\frac{m+y}{b}$ ; the reciprocal of  $\frac{1}{3y}$  is  $\frac{3y}{1}$  or  $3y$ ; the reciprocal of  $\frac{1}{4}$  is 4. Hence the reciprocal of a fraction whose numerator is 1, is the denominator of the fraction.

Thus the reciprocal of  $\frac{1}{a}$  is  $a$ ; of  $\frac{1}{a+b}$ , is  $a+b$ , &c.

165. A fraction sometimes occurs in the numerator or denominator of another fraction, as  $\frac{\frac{2}{3}a}{b}$ . It is often convenient, in the course of a calculation, to transfer such a fraction, from the numerator to the denominator of the principal fraction, or the contrary. That this may be done, without altering the value, if the fraction transferred be *inverted*, is evident, from the following principles :

First, *Dividing* by a fraction, is the same as *multiplying* by the fraction *inverted*. (Art. 162.)

Secondly, *Dividing the numerator* of a fraction has the same effect on the value, as *multiplying the denominator*; and multiplying the numerator has the same effect, as dividing the denominator. (Art. 139.)

Thus in the expression  $\frac{\frac{2}{3}a}{x}$  the numerator of  $\frac{a}{x}$  is multiplied into  $\frac{2}{3}$ . But the value will be the same, if, instead of multiplying the numerator, we divide the denominator by  $\frac{2}{3}$ , that is, multiply the denominator by  $\frac{3}{2}$ .

$$\text{Therefore } \frac{\frac{2}{3}a}{x} = \frac{a}{\frac{3}{2}x}. \quad \text{So } \frac{h}{\frac{7}{3}m} = \frac{\frac{3}{7}h}{m}.$$

$$\text{And } \frac{\frac{2}{3}d}{h+y} = \frac{d}{\frac{3}{2}(h+y)} = \frac{d}{\frac{3}{2}h + \frac{3}{2}y}. \quad \text{And } \frac{a-x}{\frac{7}{2}m} = \frac{\frac{7}{2}a - \frac{7}{2}x}{m}.$$

166. Multiplying the *numerator* is in effect multiplying the *value* of the fraction. (Art. 137.) On this principle, a fraction may be cleared of a fractional co-efficient which occurs in its numerator.

$$\text{Thus } \frac{\frac{3}{5}a}{b} = \frac{3}{5} \times \frac{a}{b} = \frac{3a}{5b}. \quad \text{And } \frac{\frac{1}{5}a}{y} = \frac{1}{5} \times \frac{a}{y} = \frac{a}{5y}.$$

$$\text{And } \frac{\frac{1}{3}h + \frac{1}{3}x}{m} = \frac{1}{3} \times \frac{h+x}{m} = \frac{h+x}{3m}. \quad \text{And } \frac{\frac{3}{5}x}{5a} = \frac{3x}{20a}.$$

$$\text{On the other hand, } \frac{3a}{7x} = \frac{3}{7} \times \frac{a}{x} = \frac{\frac{3}{7}a}{x}.$$

$$\text{And } \frac{a}{3y} = \frac{1}{3} \times \frac{a}{y} = \frac{\frac{1}{3}a}{y}. \quad \text{And } \frac{4a}{5d+5x} = \frac{\frac{4}{5}a}{d+x}.$$

167. But multiplying the *denominator*, by another fraction, is in effect *dividing* the value; (Art. 138.) that is, it is *multiplying* the value by the fraction *inverted*. The principal fraction may therefore be cleared of a fractional co-efficient, which occurs in its denominator.

$$\text{Thus } \frac{a}{\frac{3}{5}b} = \frac{a}{b} \div \frac{3}{5} = \frac{a}{b} \times \frac{5}{3} = \frac{5a}{3b}. \quad \text{And } \frac{a}{\frac{7}{2}x} = \frac{2a}{7x}.$$

$$\text{And } \frac{a+h}{\frac{3}{2}y} = \frac{9a+9h}{3y}. \quad \text{And } \frac{3h}{\frac{4}{7}m} = \frac{21h}{4m}.$$

$$\text{On the other hand, } \frac{7a}{3x} = \frac{a}{\frac{3}{7}x}.$$

$$\text{And } \frac{3y+3dx}{2m} = \frac{y+dx}{\frac{2}{3}m}. \quad \text{And } \frac{3x}{y} = \frac{x}{\frac{1}{3}y}.$$

167.b. The numerator or the denominator of a fraction, may be itself a fraction. The expression may be reduced to a more simple form, on the principles which have been applied in the preceding cases.

$$\text{Thus } \frac{\frac{a}{b}}{\frac{c}{d}} = \frac{a}{b} \div \frac{c}{d} = \frac{ad}{bc}.$$

$$\text{And } \frac{\frac{x}{y}}{h} = \frac{x}{hy}. \quad \text{And } \frac{\frac{r}{m}}{\frac{n}{m}} = \frac{nr}{m}.$$

---

SECTION VII.

---

SIMPLE EQUATIONS.

ART. 168. **T**HE subjects of the preceding sections are introductory to what may be considered the peculiar province of algebra, the investigation of the values of unknown quantities, by means of *equations*.

AN EQUATION IS A PROPOSITION, EXPRESSING IN ALGEBRAIC CHARACTERS, THE EQUALITY BETWEEN ONE QUANTITY OR SET

OF QUANTITIES AND ANOTHER, OR BETWEEN DIFFERENT EXPRESSIONS FOR THE SAME QUANTITY.\* Thus  $x+a=b+c$ , is an equation, in which the sum of  $x$  and  $a$ , is equal to the sum of  $b$  and  $c$ . The quantities on the two sides of the sign of equality, are sometimes called the *members* of the equation; the several terms on the *left* constituting the *first* member, and those on the *right*, the *second* member.

169. The object aimed at, in what is called the *resolution* or *reduction* of an equation, is to *find the value of the unknown quantity*. In the first statement of the conditions of a problem, the known and unknown quantities are frequently thrown promiscuously together. To find the value of that which is required, it is necessary to bring it to stand by itself, while all the others are on the opposite side of the equation. But, in doing this, care must be taken not to *destroy* the equation, by rendering the two members unequal. Many changes may be made in the arrangement of the terms, without affecting the equality of the sides.

170. THE REDUCTION OF AN EQUATION CONSISTS, THEN, IN BRINGING THE UNKNOWN QUANTITY BY ITSELF, ON ONE SIDE, AND ALL THE KNOWN QUANTITIES ON THE OTHER SIDE, WITHOUT DESTROYING THE EQUATION.

To effect this, it is evident that one of the members must be as much increased or diminished as the other. If a quantity be added to one, and not to the other, the equality will be destroyed. But the members will remain equal;

If the same or equal quantities be *added* to each. Ax. 1.

If the same or equal quantities be *subtracted* from each. Ax. 2.

If each be *multiplied* by the same or equal quantities. Ax. 3.

If each be *divided* by the same or equal quantities. Ax. 4.

171. It may be farther observed that, in general, if the unknown quantity is connected with others by addition, multiplication, division, &c. the reduction is made by a *contrary* process. If a known quantity is *added* to the unknown, the equation is reduced by subtraction. If one is *multiplied* by the other, the reduction is effected by *division*, &c. The reason of this will be seen, by attending to the several cases in the following articles. The *known* quantities may be expressed either by letters or figures. The *unknown* quantity is represented by one of the last letters of the alphabet, generally,  $x$ ,  $y$ , or  $z$ . (Art. 27.) The principal reductions to

\* See Note D.

be considered in this section, are those which are effected by *transposition*, *multiplication*, and *division*. These ought to be made perfectly familiar, as one or more of them will be necessary, in the resolution of almost every equation.

### TRANSPOSITION.

172. In the equation

$$x - 7 = 9,$$

the number 7 being connected with the unknown quantity  $x$  by the sign  $-$ , the one is *subtracted* from the other. To reduce the equation by a contrary process, let 7 be *added* to both sides. It then becomes

$$x - 7 + 7 = 9 + 7.$$

The equality of the members is preserved, because one is as much increased as the other. (Axiom 1.) But on one side, we have  $-7$  and  $+7$ . As these are equal, and have contrary signs, they *balance each other*, and may be cancelled. (Art. 77.) The equation will then be

$$x = 9 + 7.$$

Here the value of  $x$  is found. It is shown to be equal to  $9 + 7$ , that is to 16. The equation is therefore reduced. The unknown quantity is on one side by itself, and all the known quantities on the other side.

In the same manner, if

$$x - b = a$$

Adding  $b$  to both sides

$$x - b + b = a + b$$

And cancelling ( $-b + b$ )

$$x = a + b.$$

Here it will be seen that the last equation is the same as the first, except that  $b$  is on the opposite side, with a contrary sign.

Next suppose

$$y + c = d.$$

Here  $c$  is *added* to the unknown quantity  $y$ . To reduce the equation by a contrary process, let  $c$  be subtracted from both sides, that is, let  $-c$  be applied to both sides. We then have

$$y + c - c = d - c.$$

The equality of the members is not affected, because one is as much diminished as the other. When  $(+c - c)$  is cancelled, the equation is reduced, and is

$$y = d - c.$$

This is the same as  $y + c = d$ , except that  $c$  has been transposed, and has received a contrary sign. We hence obtain the following general rule:

173. WHEN KNOWN QUANTITIES ARE CONNECTED WITH THE UNKNOWN QUANTITY BY THE SIGN  $+$  OR  $-$ , THE EQUATION IS REDUCED BY TRANSPOSING THE KNOWN QUANTITIES TO THE OTHER SIDE, AND PREFIXING THE CONTRARY SIGN.

This is called reducing an equation by *addition* or *subtraction*, because it is, in effect, adding or subtracting certain quantities, to or from, each of the members.

Ex. 1. Reduce the equation  $x + 3b - m = h - d$   
 Transposing  $+3b$ , we have  $x - m = h - d - 3b$   
 And transposing  $-m$ ,  $x = h - d - 3b + m$

174. When several terms on the same side of an equation are *alike*, they may be united in one, by the rules for reduction in addition. (Art. 72 and 74.)

Ex. 2. Reduce the equation  $x + 5b - 4h = 7b$   
 Transposing  $5b - 4h$   $x = 7b - 5b + 4h$   
 Uniting  $7b - 5b$  in one term  $x = 2b + 4h$ .

175. The *unknown* quantity must also be transposed, whenever it is on both sides of the equation. It is not material on which side it is finally placed. For if  $x = 3$ ; it is evident that  $3 = x$ . It may be well however, to bring it on that side, where it will have the affirmative sign, when the equation is reduced.

Ex. 3. Reduce the equation  $2x + 2h = h + d + 3x$   
 By transposition  $2h - h - d = 3x - 2x$   
 And  $h - d = x$ .

176. When the *same term*, with the same sign, is on *opposite sides* of the equation, instead of transposing, we may *expunge* it from each. For this is only subtracting the same quantity from equal quantities. (Ax. 2.)

Ex. 4. Reduce the equation  $x + 3h + d = b + 3h + 7d$   
 Expunging  $3h$   $x + d = b + 7d$   
 And  $x = b + 6d$ .

177. As *all* the terms of an equation may be transposed, or supposed to be transposed; and it is immaterial which member is written first; it is evident that the *signs of all the terms may be changed*, without affecting the equality.

Thus, if we have  $x - b = d - a$   
 Then by transposition  $-d + a = -x + b$   
 Or, inverting the members  $-x + b = -d + a$ .

178. If all the terms on *one* side of an equation be transposed, each member will be equal to 0.

Thus, if  $x+b=d$ , then  $x+b-d=0$ .

It is frequently convenient to reduce an equation to this form, in which the positive and negative terms *balance* each other. In the example just given,  $x+b$  is balanced by  $-d$ . For in the first of the two equations,  $x+b$  is equal to  $d$ .

Ex. 5. Reduce  $a+2x-8=b-4+x+a$ .  $x = b + 4$

6. Reduce  $y+ab-hm=a+2y-ab+hm$ .  $2ab - 2y = hm - a$

7. Reduce  $h+30+7x=8-6h+6x-d+b$ .  $x = b - 28 - 7d$

8. Reduce  $bh+21-4x+d=12-3x+d-7bh$ .  $8bh + 9 = x$

### REDUCTION OF EQUATIONS BY MULTIPLICATION.

179. The unknown quantity, instead of being connected with a known quantity by the sign  $+$  or  $-$ , may be *divided* by it, as in the equation  $\frac{x}{a}=b$ .

Here the reduction can not be made, as in the preceding instances, by transposition. But if both members be *multiplied* by  $a$ , (Art. 170,) the equation will become

$$x=ab.$$

For a *fraction is multiplied into its denominator, by removing the denominator*. This has been proved from the properties of fractions. (Art. 159.) It is also evident from the sixth axiom.

Thus  $x = \frac{ax}{a} = \frac{3x}{3} = \frac{(a+b) \times x}{a+b} = \frac{dx+5x}{d+5}$ , &c. For in each of these instances,  $x$  is both multiplied and divided by the same quantity; and this makes no alteration in the value. Hence,

180. WHEN THE UNKNOWN QUANTITY IS DIVIDED BY A KNOWN QUANTITY, THE EQUATION IS REDUCED BY MULTIPLYING EACH SIDE BY THIS KNOWN QUANTITY.

The same transpositions are to be made in this case, as in the preceding examples. It must be observed also, that *every* term of the equation is to be multiplied. For the several terms in each member constitute a compound multiplicand, which is to be multiplied according to art. 98.

Ex. 1. Reduce the equation  $\frac{x}{c}+a=b+d$

Multiplying both sides by

$$c$$

The product is

$$x+ac=bc+cd$$

And

$$x=bc+cd-ac.$$

2. Reduce the equation  $\frac{x-4}{6} + 5 = 20$   
 Multiplying by 6  $x-4+30=120$   
 And  $x=120+4-30=94$ .

3. Reduce the equation  $\frac{x}{a+b} + d = h$   
 Multiplying by  $a+b$  (Art. 100.)  $x+ad+bd=ah+bh$   
 And  $x=ah+bh-ad-bd$ .

181. When the unknown quantity is in the *denominator* of a fraction, the reduction is made in a similar manner, by multiplying the equation by this denominator.

Ex. 4. Reduce the equation  $\frac{6}{10-x} + 7 = 8$   
 Multiplying by  $10-x$   $6+70-7x=80-8x$   
 And  $x=4$ .

182. Though it is not generally *necessary*, yet it is often convenient, to remove the denominator from a fraction consisting of *known* quantities only. This may be done, in the same manner, as the denominator is removed from a fraction which contains the unknown quantity.

Take for example  $\frac{x}{a} = \frac{d}{b} + \frac{h}{c}$   
 Multiplying by  $a$   $x = \frac{ad}{b} + \frac{ah}{c}$   
 Multiplying by  $b$   $bx = ad + \frac{abh}{c}$   
 Multiplying by  $c$   $bcx = acd + abh$ .

Or we may multiply by the product of *all* the denominators at once.

In the same equation  $\frac{x}{a} = \frac{d}{b} + \frac{h}{c}$   
 Multiplying by  $abc$   $\frac{abcx}{a} = \frac{abcd}{b} + \frac{abch}{c}$

Then by cancelling from each term, the letter which is common to its numerator and denominator, (Art. 145,) we have  $bcx = acd + abh$ , as before. Hence,

183. AN EQUATION MAY BE CLEARED OF FRACTIONS BY MULTIPLYING EACH SIDE INTO ALL THE DENOMINATORS.



Thus the equation

$$\frac{x}{a} = \frac{b}{d} + \frac{e}{g} - \frac{h}{m}$$

Is the same as

$$dgm x = abgm + adem - adgh.$$

And the equation

$$\frac{x}{2} = \frac{2}{3} + \frac{4}{5} + \frac{6}{2}$$

Is the same as

$$30x = 40 + 48 + 180.$$

In clearing an equation of fractions, it will be necessary to observe, that the sign  $-$  prefixed to any fraction, denotes that the whole value is to be subtracted, (Art. 142,) which is done by changing the signs of all the terms in the numerator.

The equation

$$\frac{a-d}{x} = c - \frac{3b + 2hm + 6n}{r}$$

Is the same as

$$ar - dr = crx - 3bx + 2hmx + 6nx.$$

### REDUCTION OF EQUATIONS BY DIVISION.

184. WHEN THE UNKNOWN QUANTITY IS MULTIPLIED INTO ANY KNOWN QUANTITY, THE EQUATION IS REDUCED BY DIVIDING BOTH SIDES BY THIS KNOWN QUANTITY. (Ax. 4.)

Ex. 1. Reduce the equation

$$ax + b - 3h = d$$

By transposition

$$ax = d + 3h - b$$

Dividing by  $a$

$$x = \frac{d + 3h - b}{a}.$$

2. Reduce the equation

$$2x = \frac{a}{c} - \frac{d}{h} + 4b$$

Clearing of fractions

$$2chx = ah - cd + 4bch$$

Dividing by  $2ch$

$$x = \frac{ah - cd + 4bch}{2ch}.$$

185. If the unknown quantity has co-efficients in *several terms*, the equation must be divided by *all* these co-efficients, connected by their signs, according to art. 121.

Ex. 3. Reduce the equation

$$3x - bx = a - d$$

That is, (Art. 120.)

$$(3-b) \times x = a - d$$

Dividing by  $3-b$

$$x = \frac{a-d}{3-b}$$

Ex. 4. Reduce the equation

$$ax + x = h - 4$$

Dividing by  $a+1$

$$x = \frac{h-4}{a+1}$$

Ex. 5. Reduce the equation

$$x - \frac{x-b}{h} = \frac{a+d}{4}$$

Clearing of fractions

$$4hx - 4x = ah + dh - 4b$$

Dividing by  $4h-4$

$$x = \frac{ah + dh - 4b}{4h - 4}$$

186. If any quantity, either known or unknown, is found as a factor in *every term*, the equation may be *divided* by it. On the other hand, if any quantity is a *divisor* in every term, the equation may be *multiplied* by it. In this way, the factor or divisor will be removed, so as to render the expression more simple.

Ex. 6. Reduce the equation

$$ax + 3ab = 6ad + a$$

Dividing by  $a$

$$x + 3b = 6d + 1$$

And

$$x = 6d + 1 - 3b$$

7. Reduce the equation

$$\frac{x+1}{x} - \frac{b}{x} = \frac{h-d}{x}$$

Multiplying by  $x$  (Art. 159.)

$$x+1-b = h-d$$

And

$$x = h-d+b-1.$$

8. Reduce the equation

$$x \times (a+b) - a - b = d \times (a+b)$$

Dividing by  $a+b$  (Art. 118.)

$$x-1 = d$$

And

$$x = d+1.$$



187. Sometimes the conditions of a problem are at first stated, not in an equation, but by means of a *proportion*. To show how this may be reduced to an equation, it will be necessary to anticipate the subject of a future section, so far as to admit the principle that "when four quantities are in geometrical proportion, the product of the two extremes is equal to the product of the two means:" a principle which is at the foundation of the Rule of Three in arithmetic. See Webber's Arithmetic.

Thus, if  $a : b :: c : d$ ,

Then  $ad = bc$

And if  $3 : 4 :: 6 : 8$ ;

And  $3 \times 8 = 4 \times 6$ . Hence,

188. A PROPORTION IS CONVERTED INTO AN EQUATION, BY MAKING THE PRODUCT OF THE EXTREMES, ONE SIDE OF THE EQUATION; AND THE PRODUCT OF THE MEANS, THE OTHER SIDE.

Ex. 1. Reduce to an equation  $ax : b :: ch : d.$   
 The product of the extremes is  $adx$   
 The product of the means is  $bch$   
 The equation is, therefore  $adx = bch.$

2. Reduce to an equation  $a + b : c :: h - m : y.$   
 The equation is  $ay + by = ch - cm.$

189. ON THE OTHER HAND, AN EQUATION MAY BE CONVERTED INTO A PROPORTION, BY RESOLVING ONE SIDE OF THE EQUATION INTO TWO FACTORS, FOR THE MIDDLE TERMS OF THE PROPORTION; AND THE OTHER SIDE INTO TWO FACTORS, FOR THE EXTREMES.

As a quantity may often be resolved into different pairs of factors; (Art. 42,) a variety of proportions may frequently be derived from the same equation.

Ex. 1. Reduce to a proportion  $abc = deh$   
 The side  $abc$  may be resolved into  $a \times bc$ , or  $ab \times c$ , or  $ac \times b.$   
 And  $deh$  may be resolved into  $d \times eh$ , or  $de \times h$  or  $dh \times e.$   
 Therefore  $a : d :: ch : bc$  And  $ac : dh :: e : b$   
 And  $ab : de :: h : c$  And  $ac : d :: eh : b \&c.$

For in each of these instances, the product of the extremes is  $abc$ , and the product of the means  $deh.$

2. Reduce to a proportion  $ax + bx = cd - ch$   
 The first member may be resolved into  $x \times (a + b)$   
 And the second into  $c \times (d - h)$   
 Therefore  $x : c :: d - h : a + b$  And  $d - h : x :: a + b : c, \&c.$

190. If for any term or terms in an equation, any other expression of the same value be *substituted*, it is manifest that the equality of the sides will not be affected.

Thus, instead of 16, we may write  $2 \times 8$  or  $\frac{64}{4}$ , or  $25 - 9$ , &c.

For these are only different forms of expression for the same quantity.

191. It will generally be well to have the several steps, in the reduction of equations, succeed each other in the following order.

First, Clear the equation of fractions. (Art. 183.)

Secondly, Transpose and unite the terms. (Arts. 173, 4, 5.)

Thirdly, Divide by the co-efficients of the unknown quantity. (Arts. 184, 5.)

*Examples.*

1. Reduce the equation

Clearing of fractions  
 Transp. and uniting terms  
 Dividing by 4

$$\frac{3x}{4} + 6 = \frac{5x}{8} + 7$$

$$24x + 192 = 20x + 224$$

$$4x = 32$$

$$x = 8.$$

2. Reduce the equation

Clearing of fractions

Dividing

$$\frac{x}{a} + h = \frac{x}{b} - \frac{x}{c} + d$$

$$bcx + abx - acx = abcd - abch$$

$$x = \frac{abcd - abch}{bc + ab - ac}$$

3. Reduce  $40 - 6x - 16 = 120 - 14x$ .Ans.  $x = 12$ .4. Reduce  $\frac{x-3}{2} + \frac{x}{3} = 20 - \frac{x-19}{2}$ Ans.  $x = \frac{93}{4}$ .5. Reduce  $\frac{x}{3} + \frac{x}{5} = 20 - \frac{x}{4}$ .6. Reduce  $\frac{1-a}{x} - 4 = 5$ .7. Reduce  $\frac{3}{x+4} - 2 = 8$ .8. Reduce  $\frac{6x}{x+4} = 1$ .9. Reduce  $x + \frac{x}{2} + \frac{x}{3} = 11$ .10. Reduce  $\frac{x}{2} + \frac{x}{3} - \frac{x}{4} = \frac{7}{10}$ .11. Reduce  $\frac{x-5}{4} + 6x = \frac{284-x}{5}$ .12. Reduce  $3x + \frac{2x+6}{5} = 5 + \frac{11x-37}{2}$ .13. Reduce  $\frac{6x-4}{3} - 2 = \frac{18-4x}{3} + x$ .14. Reduce  $21 + \frac{3x-11}{16} = \frac{5x-5}{8} + \frac{97-7x}{2}$ .15. Reduce  $3x - \frac{x-4}{4} - 4 = \frac{5x+14}{3} - \frac{1}{12}$ .16. Reduce  $\frac{7x+5}{3} - \frac{16+4x}{5} + 6 = \frac{3x+9}{2}$ .17. Reduce  $\frac{17-3x}{5} - \frac{4x+2}{3} = 5 - 6x + \frac{7x+14}{3}$ .18. Reduce  $x - \frac{3x-3}{5} + 4 = \frac{20-x}{2} - \frac{6x-8}{7} + \frac{4x-4}{5}$ .

19. Reduce  $\frac{6x+7}{9} + \frac{7x-13}{6x+3} = \frac{2x+4}{3}$ .

20. Reduce  $\frac{5x+4}{2} : \frac{18-x}{4} :: 7:4$ .

---

### SOLUTION OF PROBLEMS.

192. In the solution of problems, by means of equations, two things are necessary: First, to translate the statement of the question from common to algebraic language, in such a manner as to form an equation: Secondly, to reduce this equation to a state in which the unknown quantity will stand by itself, and its value be given in known terms, on the opposite side. The manner in which the latter is effected, has already been considered. The former will probably occasion more perplexity to a beginner; because the conditions of questions are so various in their nature, that the proper method of stating them cannot be easily learned, like the reduction of equations, by a system of definite rules. Practice however will soon remove a great part of the difficulty.

193. It is one of the principal peculiarities of an algebraic solution, that the *quantity sought* is itself introduced into the operation. This enables us to make a statement of the conditions, in the same form, as though the problem were already solved. Nothing then remains to be done, but to *reduce* the equation, and to find the aggregate value of the known quantities. (Art. 53.) As these are equal to the *unknown* quantity on the other side of the equation, the value of that also is determined, and therefore the problem is solved.

Problem 1. A man being asked how much he gave for his watch, replied; If you multiply the price by 4, and to the product add 70, and from this sum subtract 50, the remainder will be equal to 220 dollars.

To solve this, we must first translate the conditions of the problem, into such algebraic expressions, as will form an equation.

Let the price of the watch be represented by	$x$
This price is to be mult'd by 4, which makes	$4x$
To the product, 70 is to be added, making	$4x+70$
From this, 50 is to be subtracted, making	$4x+70-50$

Here we have a number of the conditions, expressed in algebraic terms; but have as yet no *equation*. We must observe then, that by the last condition of the problem, the preceding terms are said to be *equal* to 220.

We have, therefore, this equation  $4x + 70 - 50 = 220$ .  
Which reduced gives  $x = 50$ .

Here the value of  $x$  is found to be 50 dollars, which is the price of the watch.

194. To *prove* whether we have obtained the true value of the letter which represents the unknown quantity, we have only to substitute this value, for the letter itself, in the equation which contains the first statement of the conditions of the problem; and to see whether the sides are equal, after the substitution is made. For if the answer thus satisfies the conditions proposed, it is the quantity sought. Thus, in the preceding example,

The original equation is  $4x + 70 - 50 = 220$   
Substituting 50 for  $x$ , it becomes  $4 \times 50 + 70 - 50 = 220$   
That is,  $220 = 220$ .

Prob. 2. What number is that, to which, if its half be added, and from the sum 20 be subtracted, the remainder will be a fourth part of the number itself?

In stating questions of this kind, where fractions are concerned, it should be recollected, that  $\frac{1}{3}x$  is the same as  $\frac{x}{3}$ ; that  $\frac{2}{5}x = \frac{2x}{5}$ , &c. (Art. 161.)

In this problem, let  $x$  be put for the number required.

Then by the conditions proposed,  $x + \frac{x}{2} - 20 = \frac{x}{4}$

And reducing the equation  $x = 16$ .

Proof  $16 + \frac{16}{2} - 20 = \frac{16}{4}$ .

Prob. 3. A father divides his estate among his three sons, in such a manner, that,

The first has \$1000 less than half of the whole;

The second has 300 less than one third of the whole;

The third has 600 less than a fourth of the whole;

What is the value of the estate?

If the whole estate be represented by  $x$ , then the several shares will be  $\frac{x}{2} - 1000$ , and  $\frac{x}{3} - 300$ , and  $\frac{x}{4} - 600$ .

And as these constitute the whole estate, they are together equal to  $x$ .

We have then this equation  $\frac{x}{2} - 1000 + \frac{x}{3} - 800 + \frac{x}{4} - 600 = x$ .

Which reduced gives  $x = 28800$ .

Proof  $\frac{28800}{2} - 1000 + \frac{28800}{3} - 800 + \frac{28800}{4} - 600 = 28800$ .

195. To avoid an unnecessary introduction of unknown quantities into an equation, it may be well to observe, in this place, that when the *sum* or *difference* of two quantities is given, both of them may be expressed by means of the same letter. For if one of the two quantities be subtracted from their sum, it is evident the remainder will be equal to the other. And if the difference of two quantities be subtracted from the greater, the remainder will be the less.

Thus, if the sum of the two numbers be	20
And if one of them be represented by	$x$
The other will be equal to	$20 - x$

Prob. 4. Divide 48 into two such parts, that if the less be divided by 4, and the greater by 6, the sum of the quotients will be 9.

Here, if  $x$  be put for the smaller part, the greater will be  $48 - x$ .

By the conditions of the problem	$\frac{x}{4} + \frac{48 - x}{6} = 9$ .
----------------------------------	--

Therefore	$x = 12$ , the less.
-----------	----------------------

And	$48 - x = 36$ , the greater.
-----	------------------------------

196. Letters may be employed to express the *known* quantities in an equation, as well as the unknown. A particular value is assigned to the numbers, when they are introduced into the calculation: and at the close, the numbers are restored. (Art. 52.)

Prob. 5. If to a certain number, 720 be added, and the sum be divided by 125; the quotient will be equal to 7392 divided by 462. What is that number?

Let  $x =$  the number required.

$a = 720$	$d = 7392$
$b = 125$	$h = 462$

Then by the conditions of the problem

$$\frac{x+a}{b} = \frac{d}{h}.$$

Therefore

$$x = \frac{bd-ah}{h}$$

Restoring the numbers,  $x = \frac{(125 \times 7392) - (720 \times 462)}{462} = 1280.$

197. When the resolution of an equation brings out a *negative* answer, it shows that the value of the unknown quantity is *contrary* to the quantities which, in the statement of the question, are considered positive. See *Negative Quantities*. (Art. 54, &c.)

Prob. 6. A merchant gains or loses, in a bargain, a certain sum. In a second bargain, he gains 350 dollars, and, in a third, loses 60. In the end, he finds he has gained 200 dollars, by the three together. How much did he gain or lose by the first?

In this example, as the profit and loss are opposite in their nature, they must be distinguished by contrary signs. (Art. 57.) If the profit is marked +, the loss must be -.

Let  $x$  = the sum required.

Then according to the statement  $x + 350 - 60 = 200$

And

$$x = -90$$

The negative sign prefixed to the answer, shows that there was a *loss* in the first bargain; and therefore that the proper sign of  $x$  is negative also. But this being determined by the answer, the omission of it in the course of the calculation can lead to no mistake.

Prob. 7. A ship sails 4 degrees north, then 13 S. then 17 N. then 19 S. and has finally 11 degrees of south latitude. What was her latitude at starting?

Let  $x$  = the latitude sought.

Then marking the northings +, and the southings -;

By the statement

$$x + 4 - 13 + 17 - 19 = -11$$

And

$$x = 0.$$

The answer here shows that the place from which the ship started was on the equator, where the latitude is nothing.

Prob. 8. If a certain number is divided by 12, the quotient, dividend, and divisor added together, will amount to 64. What is the number?



Let  $x =$  the number sought.

Then

$$\frac{x}{12} + x + 12 = 64$$

And

$$x = \frac{624}{13} = 48.$$

Prob. 9. An estate is divided among four children, in such a manner, that

The first has 200 dollars more than  $\frac{1}{4}$  of the whole,

The second has 340 dollars more than  $\frac{1}{5}$  of the whole,

The third has 300 dollars more than  $\frac{1}{6}$  of the whole,

The fourth has 400 dollars more than  $\frac{1}{8}$  of the whole.

What is the value of the estate?      Ans. 4800 dollars.

Prob. 10. What is that number which is as much less than 500, as a fifth part of it is greater than 40?      Ans. 450.

Prob. 11. There are two numbers whose difference is 40, and which are to each other as 6 to 5. What are the numbers?      Ans. 240 and 200.

Prob. 12. Three persons,  $A$ ,  $B$ , and  $C$  draw prizes in a lottery.  $A$  draws 200 dollars;  $B$  draws as much as  $A$ , together with a third of what  $C$  draws; and  $C$  draws as much as  $A$  and  $B$  both. What is the amount of the three prizes?      Ans. 1200 dollars.

Prob. 13. What number is that, which is to 12 increased by three times the number, as 2 to 9?      Ans. 8.

Prob. 14. A ship and a boat are descending a river at the same time. The ship passes a certain fort, when the boat is 13 miles below. The ship descends five miles, while the boat descends three. At what distance below the fort, will they be together?      Ans.  $32\frac{1}{2}$  miles.

Prob. 15. What number is that, a sixth part of which exceeds an eighth part of it by 20?      Ans. 480.

Prob. 16. Divide a prize of 2000 dollars into two such parts, that one of them shall be to the other, as 9 : 7.

Ans. The parts are 1125, and 875.

Prob. 17. What sum of money is that, whose third part, fourth part, and fifth part, added together, amount to 94 dollars?      Ans. 120 dollars.

Prob. 18. Two travellers, *A* and *B*, 360 miles apart, travel towards each other till they meet. *A*'s progress is 10 miles an hour, and *B*'s 8. How far does each travel before they meet?      Ans. *A* goes 200 miles, and *B* 160.

Prob. 19. A man spent one third of his life in England, one fourth of it in Scotland, and the remainder of it, which was 20 years, in the United States. To what age did he live?      Ans. To the age of 48.

Prob. 20. What number is that,  $\frac{1}{4}$  of which is greater than  $\frac{1}{5}$  of it by 96?

Prob. 21. A post is  $\frac{1}{5}$  in the earth,  $\frac{3}{7}$  in the water, and 13 feet above the water. What is the length of the post?      Ans. 35 feet.

Prob. 22. What number is that, to which 10 being added,  $\frac{2}{5}$  of the sum will be 66?

Prob. 23. Of the trees in an orchard,  $\frac{3}{4}$  are apple trees,  $\frac{1}{10}$  pear trees, and the remainder peach trees, which are 20 more than  $\frac{1}{8}$  of the whole. What is the whole number in the orchard?      Ans. 800.

Prob. 24. A gentleman bought several gallons of wine for 94 dollars; and after using 7 gallons himself, sold  $\frac{1}{4}$  of the remainder for 20 dollars. How many gallons had he at first?      Ans. 47.

Prob. 25. *A* and *B* have the same income. *A* contracts an annual debt amounting to  $\frac{1}{7}$  of it; *B* lives upon  $\frac{4}{5}$  of it; at the end of 10 years, *B* lends to *A* enough to pay off his debts, and has 160 dollars to spare. What is the income of each?      Ans. 280 dollars.

Prob. 26. A gentleman lived single  $\frac{1}{4}$  of his whole life; and after having been married 5 years more than  $\frac{1}{4}$  of his life, he had a son who died 4 years before him, and who reached only half the age of his father. To what age did the father live?      Ans. 84.

Prob. 27. What number is that, to which, if  $\frac{1}{3}$ ,  $\frac{1}{4}$ , and  $\frac{2}{7}$  of it be added, the sum will be 73?

Prob. 28. A person, after spending 100 dollars more than  $\frac{1}{5}$  of his income, had remaining 35 dollars more than  $\frac{1}{2}$  of it. Required his income.

Prob. 29. In the composition of a quantity of gunpowder,  
 The *nitre* was 10 lb. more than  $\frac{2}{3}$  of the whole,  
 The *sulphur*  $4\frac{1}{2}$  lb. less than  $\frac{1}{6}$  of the whole,  
 The *charcoal* 2 lb. less than  $\frac{1}{7}$  of the nitre.  
 What was the amount of gunpowder? Ans. 69 lb.

Prob. 30. A cask which held 146 gallons, was filled with a mixture of brandy, wine, and water. There were 15 gallons of wine more than of brandy, and as much water as the brandy and wine together. What quantity was there of each?

Prob. 31. Four persons purchased a farm in company for 4755 dollars; of which *B* paid three times as much as *A*; *C* paid as much as *A* and *B*; and *D* paid as much as *C* and *B*. What did each pay? Ans. 317, 951, 1268, 2219.

Prob. 32. It is required to divide the number 99 into five such parts, that the first may exceed the second by 3, be less than the third by 10, greater than the fourth by 9, and less than the fifth by 16.

Let  $x =$  the first part.

Then  $x - 3 =$  the second,  $x - 9 =$  the fourth,

$x + 10 =$  the third,  $x + 16 =$  the fifth.

Therefore  $x + x - 3 + x + 10 + x - 9 + x + 16 = 99$ .

And  $x = 17$ .

Prob. 33. A father divided a small sum among four sons.  
 The third had 9 shillings more than the fourth;  
 The second had 12 shillings more than the third;  
 The first had 18 shillings more than the second;  
 And the whole sum was 6 shillings more than 7 times the sum which the youngest received.

What was the sum divided? Ans. 153.

Prob. 34. A farmer had two flocks of sheep, each containing the same number. Having sold from one of these 39, and from the other 93, he finds twice as many remaining in the one, as in the other. How many did each flock originally contain?

Prob. 35. An express, travelling at the rate of 60 miles a day, had been dispatched 5 days, when a second was sent after him, travelling 75 miles a day. In what time will the one overtake the other? Ans. 20 days.

Prob. 36. The age of *A* is double that of *B*, the age of *B* triple that of *C*, and the sum of all their ages 140. What is the age of each?

Prob. 37. Two pieces of cloth, of the same price by the yard, but of different lengths, were bought, the one for five pounds, the other for  $6\frac{1}{2}$ . If 10 be added to the length of each, the sums will be as 5 to 6. Required the length of each piece.

Prob. 38.  $A$  and  $B$  began trade with equal sums of money. The first year,  $A$  gained forty pounds, and  $B$  lost 40. The second year,  $A$  lost  $\frac{1}{3}$  of what he had at the end of the first, and  $B$  gained 40 pounds less than twice the sum which  $A$  had lost.  $B$  had then twice as much money as  $A$ . What sum did each begin with?      Ans. 320 pounds.

Prob. 39. What number is that, which being severally added to 36 and 52, will make the former sum to the latter, as 3 to 4?

Prob. 40. A gentleman bought a chaise, horse, and harness, for 360 dollars. The horse cost twice as much as the harness; and the chaise cost twice as much as the harness and horse together. What was the price of each?

Prob. 41. Out of a cask of wine, from which had leaked  $\frac{1}{3}$  part, 21 gallons were afterwards drawn; when the cask was found to be half full. How much did it hold?

Prob. 42. A man has 6 sons, each of whom is 4 years older than his next younger brother; and the eldest is three times as old as the youngest. What is the age of each?

Prob. 43. Divide the number 49 into two such parts, that the greater increased by 6, shall be to the less diminished by 11, as 9 to 2.

Prob. 44. What two numbers are as 2 to 3; to each of which, if 4 be added, the sums will be as 5 to 7?

Prob. 45. A person bought two casks of porter, one of which held just 3 times as much as the other; from each of these he drew 4 gallons, and then found that there were 4 times as many gallons remaining in the larger, as in the other. How many gallons were there in each?

Prob. 46. Divide the number 68 into two such parts, that the difference between the greater and 84, shall be equal to 3 times the difference between the less and 40.

Prob. 47. Four places are situated in the order of the letters  $A, B, C, D$ . The distance from  $A$  to  $D$ , is 34 miles. The distance from  $A$  to  $B$  is to the distance from  $C$  to  $D$  as 2 to 3. And  $\frac{1}{4}$  of the distance from  $A$  to  $B$ , added to half

the distance from *C* to *D*, is three times the distance from *B* to *C*. What are the respective distances ?

Ans. From *A* to *B*=12; from *B* to *C*=4; from *C* to *D*=18.

Prob. 48. Divide the number 36 into 3 such parts, that  $\frac{1}{2}$  of the first,  $\frac{1}{3}$  of the second, and  $\frac{1}{4}$  of the third, shall be equal to each other.

Prob. 49. A merchant supported himself 3 years for 50 pounds a year, and at the end of each year, added to that part of his stock which was not thus expended, a sum equal to one third of this part. At the end of the third year, his original stock was doubled. What was that stock ?

Ans. 740 pounds.

Prob. 50. A general having lost a battle, found that he had only half of his army, +3600 men, left fit for action;  $\frac{1}{8}$  of the army +600 being wounded; and the rest, who were  $\frac{1}{2}$  of the whole, either slain, taken prisoners, or missing. Of how many men did his army consist ? Ans. 24000.

For the solution of many algebraic problems, an acquaintance with the calculations of powers and radical quantities is required. It will therefore be necessary to attend to these, before finishing the subject of equations.



## SECTION VIII.

### INVOLUTION AND POWERS.

ART. 198. **W**HEN A QUANTITY IS MULTIPLIED INTO ITSELF, THE PRODUCT IS CALLED A POWER.

Thus  $2 \times 2 = 4$ , the square or second power of 2.  
 $2 \times 2 \times 2 = 8$ , the cube or third power.  
 $2 \times 2 \times 2 \times 2 = 16$ , the fourth power, &c.

So  $10 \times 10 = 100$ , the second power of 10.  
 $10 \times 10 \times 10 = 1000$ , the third power.  
 $10 \times 10 \times 10 \times 10 = 10000$ , the fourth power, &c.

And  $a \times a = aa$ , the second power of  $a$ .  
 $a \times a \times a = aaa$ , the third power.  
 $a \times a \times a \times a = aaaa$ , the fourth power, &c.

199. The original quantity itself, though not, like the powers proceeding from it, produced by multiplication, is nevertheless called the *first power*. It is also called the *root* of the other powers, because it is that from which they are all derived.

200. As it is inconvenient, especially in the case of high powers, to write down all the letters or factors of which the powers are composed, an abridged method of notation is generally adopted. The root is written only once; and then a number or letter is placed at the right hand, and a little elevated, to signify how many times the root is *employed as a factor*, to produce the power. This number or letter is called the *index* or *exponent* of the power. Thus  $a^2$  is put for  $a \times a$  or  $aa$ , because the root  $a$  is *twice* repeated as a factor, to produce the power  $aa$ . And  $a^3$  stands for  $aaa$ ; for here  $a$  is repeated *three times* as a factor.

The index of the *first power* is 1; but this is commonly omitted. Thus  $a^1$  is the same as  $a$ .

201. Exponents must not be confounded with *co-efficients*. A co-efficient shows how often a quantity is taken as a *part* of a whole. An exponent shows how often a quantity is taken as a *factor* in a product.

Thus  $4a = a + a + a + a$ .

But  $a^4 = a \times a \times a \times a$ .

202. The scheme of notation by exponents has the peculiar advantage of enabling us to express an *unknown* power. For this purpose the index is a *letter*, instead of a numerical figure. In the solution of a problem, a quantity may occur, which we know to be *some* power of another quantity. But it may not be yet ascertained whether it is a square, a cube, or some higher power. Thus in the expression  $a^x$ , the index  $x$  denotes that  $a$  is involved to *some* power, though it does not determine *what* power. So  $b^m$  and  $d^n$  are powers of  $b$  and  $d$ ; and are read the  $m$ th power of  $b$ , and the  $n$ th power of  $d$ . When the value of the index is found, a *number* is generally substituted for the letter. Thus, if  $m=3$ , then  $b^m = b^3$ ; but if  $m=5$ , then  $b^m = b^5$ .

203. The method of expressing powers by exponents is also of great advantage in the case of *compound* quantities.

Thus  $\overline{a+b+d}^3$  or  $\overline{a+b+d}^3$  or  $(a+b+d)^3$ , is  $(a+b+d) \times (a+b+d) \times (a+b+d)$  that is, the cube of  $(a+b+d)$ . But this involved at length would be  $a^3+3a^2b+3a^2d+3ab^2+6abd+3ad^2+b^3+3b^2d+3bd^2+d^3$ .

204. If we take a series\* of powers whose indices increase or decrease by 1, we shall find that the powers themselves increase by a *common multiplier*, or decrease by a *common divisor*; and that this multiplier or divisor is the original quantity from which the powers are raised.

Thus in the series  $aaaaa, aaaa, aaa, aa, a$ ;  
Or  $a^5, a^4, a^3, a^2, a^1$ ;

the indices counted from right to left are 1, 2, 3, 4, 5; and the common difference between them is a unit. If we begin on the *right*, and *multiply* by  $a$ , we produce the several powers, in succession, from right to left.

Thus  $a \times a = a^2$  the 2d term. And  $a^3 \times a = a^4$ .  
 $a^2 \times a = a^3$  the 3d term.  $a^4 \times a = a^5$ , &c.

If we begin on the *left*, and *divide* by  $a$ ,

We have  $a^5 \div a = a^4$ . And  $a^3 \div a = a^2$ .  
 $a^4 \div a = a^3$ .  $a^2 \div a = a^1$ .

205. But this division may be carried still farther; and we shall then obtain a new set of quantities.

Thus  $a \div a = \frac{a}{a} = 1$ . (Art. 128.) And  $\frac{1}{a} \div a = \frac{1}{aa}$ . (Art. 163.)  
 $1 \div a = \frac{1}{a}$ .  $\frac{1}{aa} \div a = \frac{1}{aaa}$ , &c.

The whole series then

Is  $aaaaa, aaaa, aaa, aa, a, 1, \frac{1}{a}, \frac{1}{aa}, \frac{1}{aaa}$ , &c.

Or  $a^5, a^4, a^3, a^2, a, 1, \frac{1}{a}, \frac{1}{a^2}, \frac{1}{a^3}$ , &c.

Here the quantities on the *right* of 1, are the *reciprocals* of those on the *left*. (Art. 49.) The former, therefore, may be properly called *reciprocal powers* of  $a$ ; while the latter may be termed, for distinction sake, *direct powers* of  $a$ . It may be added, that the powers on the left are also the reciprocals of those on the right.

\* NOTE. The term *series* is applied to a number of quantities succeeding each other, in some regular order. It is not confined to any particular law of increase or decrease.

For  $1 \div \frac{1}{a} = 1 \times \frac{a}{1} = a$ . (Art. 162.) And  $1 \div \frac{1}{a^3} = a^3$ .

$$1 \div \frac{1}{a^2} = 1 \times \frac{a^2}{1} = a^2. \qquad 1 \div \frac{1}{a^4} = a^4, \text{ \&c.}$$

206. The same plan of notation is applicable to *compound* quantities. Thus from  $a+b$ , we have the series,

$$(a+b)^3, (a+b)^2, (a+b), 1, \frac{1}{a+b}, \frac{1}{(a+b)^2}, \frac{1}{(a+b)^3}, \text{ \&c.}$$

207. For the convenience of calculation, another form of notation is given to reciprocal powers.

According to this,  $\frac{1}{a}$  or  $\frac{1}{a^1} = a^{-1}$ . And  $\frac{1}{aaa}$  or  $\frac{1}{a^3} = a^{-3}$ .

$$\frac{1}{aa} \text{ or } \frac{1}{a^2} = a^{-2}. \qquad \frac{1}{aaaa} \text{ or } \frac{1}{a^4} = a^{-4}, \text{ \&c.}$$

And to make the indices a complete series, with 1 for the common difference, the term  $\frac{a}{a}$  or 1, which is considered as *no* power, is written  $a^0$ .

The powers both direct and reciprocal\* then,

Instead of  $aaaa, aaa, aa, a, \frac{a}{a}, \frac{1}{a}, \frac{1}{aa}, \frac{1}{aaa}, \frac{1}{aaaa}, \text{ \&c.}$

Will be  $a^4, a^3, a^2, a^1, a^0, a^{-1}, a^{-2}, a^{-3}, a^{-4}, \text{ \&c.}$

Or  $a^{+4}, a^{+3}, a^{+2}, a^{+1}, a^0, a^{-1}, a^{-2}, a^{-3}, a^{-4}, \text{ \&c.}$

And the indices taken by themselves will be,

$$+4, +3, +2, +1, 0, -1, -2, -3, -4, \text{ \&c.}$$

208. The root of a power may be expressed by more letters than one.

Thus  $aa \times aa$ , or  $\overline{aa}^2$  is the second power of  $aa$ .

And  $aa \times aa \times aa$ , or  $\overline{aa}^3$  is the third power of  $aa$ , &c.

Hence a certain power of one quantity, may be a different power of another quantity. Thus  $a^4$  is the second power of  $a^2$ , and the fourth power of  $a$ .

209. All the powers of 1 are the same. For  $1 \times 1$ , or  $1 \times 1 \times 1$ , &c. is still 1.

\* See Note E.



## INVOLUTION.

210. Involution is finding any power of a quantity, by multiplying it into itself. The reason of the following general rule is manifest, from the nature of powers.

MULTIPLY THE QUANTITY INTO ITSELF, TILL IT IS TAKEN AS A FACTOR, AS MANY TIMES AS THERE ARE UNITS IN THE INDEX OF THE POWER TO WHICH THE QUANTITY IS TO BE RAISED.

This rule comprehends all the instances which can occur in involution. But it will be proper to give an explanation of the manner in which it is applied to particular cases.

211. A single letter is involved, by giving it the index of the proposed power; or by repeating it as many times, as there are units in that index.

The 4th power of  $a$ , is  $a^4$  or  $aaaa$ . (Art. 198.)

The 6th power of  $y$ , is  $y^6$  or  $yyyyyy$ .

The  $n$ th power of  $x$ , is  $x^n$  or  $xxx \dots n$  times repeated.

212. The method of involving a quantity which consists of several factors, depends on the principle, that *the power of the product of several factors is equal to the product of their powers.*

Thus  $(ay)^2 = a^2y^2$ . For by art. 210;  $(ay)^2 = ay \times ay$ .

But  $ay \times ay = ayy = a^2y^2$ .

So  $(bmx)^3 = bmx \times bmx \times bmx = bbbmmmmxxx = b^3m^3x^3$ .

And  $(ady)^n = ady \times ady \times ady \dots n$  times  $= a^n d^n y^n$ .

In finding the power of a product, therefore, we may either involve the whole at once; or we may involve each of the factors separately, and then multiply their several powers into each other.

Ex. 1. The 4th power of  $dhy$ , is  $(dhy)^4$ , or  $d^4h^4y^4$ .

2. The 3d power of  $4b$ , is  $(4b)^3$ , or  $4^3b^3$ , or  $64b^3$ .

3. The  $n$ th power of  $6ad$ , is  $(6ad)^n$ , or  $6^n a^n d^n$ .

4. The 3d power of  $3m \times 2y$ , is  $(3m \times 2y)^3$ , or  $27m^3 \times 8y^3$ .

213. A compound quantity, consisting of terms connected by  $+$  and  $-$ , is involved by an actual multiplication of its several parts. Thus,

$(a+b)^1 = a+b$ , the first power.

$$\begin{array}{r} a^2 + ab \\ + ab + b^2 \\ \hline \end{array}$$

$(a+b)^2 = a^2 + 2ab + b^2$ , the second power of  $(a+b)$ .

$$\begin{array}{r} a^3 + 2a^2b + ab^2 \\ + a^2b + 2ab^2 + b^3 \\ \hline \end{array}$$

$(a+b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$ , the third power.

$$\begin{array}{r} a^4 + 3a^3b + 3a^2b^2 + ab^3 \\ + a^3b + 3a^2b^2 + 3ab^3 + b^4 \\ \hline \end{array}$$

$(a+b)^4 = a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4$ , the 4th power, &c.

2. The square of  $a-b$ , is  $a^2 - 2ab + b^2$ .
3. The cube of  $a+1$ , is  $a^3 + 3a^2 + 3a + 1$ .
4. The square  $a+b+h$ , is  $a^2 + 2ab + 2ah + b^2 + 2bh + h^2$ .
5. Required the cube of  $a+2d+3$ .
6. Required the 4th power of  $b+2$ .
7. Required the 5th power of  $x+1$ .
8. Required the 6th power of  $1-b$ .

214. The squares of *binomial* and *residual* quantities occur so frequently in algebraic processes, that it is important to make them familiar.

If we multiply  $a+h$  into itself, and also  $a-h$ ,

$$\begin{array}{r} \text{We have } a+h \\ a+h \\ \hline \end{array}$$

$$\begin{array}{r} a^2 + ah \\ + ah + h^2 \\ \hline \end{array}$$

$$a^2 + 2ah + h^2.$$

$$\begin{array}{r} \text{And } a-h \\ a-h \\ \hline \end{array}$$

$$\begin{array}{r} a^2 - ah \\ - ah + h^2 \\ \hline \end{array}$$

$$a^2 - 2ah + h^2.$$

Here it will be seen that, in each case, the first and last terms are squares of  $a$  and  $h$ ; and that the middle term is twice the product of  $a$  into  $h$ . Hence the squares of bino-

mial and residual quantities, without multiplying each of the terms separately, may be found, by the following proposition.\*

THE SQUARE OF A BINOMIAL, THE TERMS OF WHICH ARE BOTH POSITIVE, IS EQUAL TO THE SQUARE OF THE FIRST TERM, + TWICE THE PRODUCT OF THE TWO TERMS; + THE SQUARE OF THE LAST TERM.

And the square of a *residual* quantity, is equal to the square of the first term, - twice the product of the two terms, + the square of the last term.

- Ex. 1. The square of  $2a+b$ , is  $4a^2 + 4ab + b^2$ .  
 2. The square of  $h+1$ , is  $h^2 + 2h + 1$ .  
 3. The square of  $ab+cd$ , is  $a^2b^2 + 2abcd + c^2d^2$ .  
 4. The square of  $6y+3$ , is  $36y^2 + 36y + 9$ .  
 5. The square of  $3d-h$ , is  $9d^2 - 6dh + h^2$ .  
 6. The square of  $a-1$ , is  $a^2 - 2a + 1$ .

For the method of finding the higher powers of binomials, see one of the succeeding sections.

215. For many purposes, it will be sufficient to express the powers of compound quantities by *exponents*, without an actual multiplication.

Thus the square of  $a+b$ , is  $\overline{a+b}^2$ , or  $(a+b)^2$ . Art. 203.

The  $n$ th power of  $bc+8+x$ , is  $(bc+8+x)^n$ .

In cases of this kind, the vinculum must be drawn over *all* the terms of which the compound quantity consists.

216. But if the root consists of several *factors*, the vinculum which is used in expressing the power, may either extend over the whole; or may be applied to each of the factors separately, as convenience may require.

Thus the square of  $\overline{a+b} \times \overline{c+d}$ , is either  
 $\overline{a+b \times c+d}^2$  or  $\overline{a+b}^2 \times \overline{c+d}^2$ .

For the first of these expressions is the square of the product of the two factors, and the last is the product of their squares. But one of these is equal to the other. (Art. 212.)

The cube of  $a \times \overline{b+d}$ , is  $(a \times \overline{b+d})^3$ , or  $a^3 \times (b+d)^3$ .

\* Euclid's Elements, Book II. Prop. 4.

217. When a quantity whose power has been expressed by a vinculum and an index, is afterwards involved by an actual multiplication of the terms, it is said to be *expanded*.

Thus  $(a+b)^2$ , when expanded, becomes  $a^2+2ab+b^2$ .

And  $(a+b+h)^2$ , becomes  $a^2+2ab+2ah+b^2+2bh+h^2$ .

218. With respect to the sign which is to be prefixed to quantities involved, it is important to observe, that WHEN THE ROOT IS POSITIVE, ALL ITS POWERS ARE POSITIVE ALSO; BUT WHEN THE ROOT IS NEGATIVE, THE ODD POWERS ARE NEGATIVE, WHILE THE EVEN POWERS ARE POSITIVE.

For the proof of this, see art. 109.

The 2d power of  $-a$  is  $+a^2$

The 3d power is  $-a^3$

The 4th power is  $+a^4$

The 5th power is  $-a^5$ , &c.

219. Hence any *odd* power has the same sign as its root. But an *even* power is positive, whether its root is positive or negative.

Thus  $+a \times +a = a^2$

And  $-a \times -a = a^2$ .

220. A QUANTITY WHICH IS ALREADY A POWER, IS INVOLVED BY MULTIPLYING ITS INDEX, INTO THE INDEX OF THE POWER TO WHICH IT IS TO BE RAISED.

1. The 3d power of  $a^2$ , is  $a^{2 \times 3} = a^6$ .

For  $a^2 = aa$ : and the cube of  $aa$  is  $aa \times aa \times aa = aaaaaa = a^6$ ; which is the 6th power of  $a$ , but the 3d power of  $a^2$ .

For a farther illustration of this rule, see arts. 233, 4.

2. The 4th power of  $a^3b^2$ , is  $a^{3 \times 4}b^{2 \times 4} = a^{12}b^8$ .

3. The 3d power of  $4a^2x$ , is  $64a^6x^3$ .

4. The 4th power of  $2a^3 \times 3x^2d$ , is  $16a^{12} \times 81x^8d^4$ ,

5. The 5th power of  $(a+b)^2$ , is  $(a+b)^{10}$ .

6. The  $n$ th power of  $a^3$ , is  $a^{3n}$ .

7. The  $n$ th power of  $(x-y)^m$ , is  $(x-y)^{mn}$ .

8.  $\overline{a^3+b^3}^2 = a^6 + 2a^3b^3 + b^6$ . (Art. 214.)

9.  $\overline{a^3 \times b^3}^2 = a^6 \times b^6$ .      10.  $\overline{(a^3b^2h^4)}^3 = a^9b^6h^{12}$ .

221. The rule is equally applicable to powers whose exponents are *negative*.

Ex. 1. The 3d power of  $a^{-2}$ , is  $a^{-2 \times 3} = a^{-6}$ .

For  $a^{-2} = \frac{1}{aa}$ . (Art. 207.) And the 3d power of this is

$$\frac{1}{aa} \times \frac{1}{aa} \times \frac{1}{aa} = \frac{1}{aaaaaa} = \frac{1}{a^6} = a^{-6}.$$

2. The 4th power  $a^2 b^{-3}$ , is  $a^8 b^{-12}$ , or  $\frac{a^8}{b^{12}}$ .

3. The cube of  $2x^m y^{-m}$ , is  $8x^{3m} y^{-3m}$ .

4. The square of  $b^3 x^{-1}$ , is  $b^6 x^{-2}$ .

5. The  $n$ th power of  $x^{-m}$ , is  $x^{-mn}$ , or  $\frac{1}{x^{mn}}$ .

222. It must be observed here, as in art. 218, that if the sign which is *prefixed* to the power be  $-$ , it must be changed to  $+$ , whenever the index becomes an even number.

Ex. 1. The square of  $-a^3$ , is  $+a^6$ . For the square of  $-a^3$ , is  $-a^3 \times -a^3$ , which, according to the rules for the signs in multiplication, is  $+a^6$ .

2. But the *cube* of  $-a^3$ , is  $-a^9$ . For  $-a^3 \times -a^3 \times -a^3 = -a^9$ .

3. The square of  $-x^n$ , is  $+x^{2n}$ .

4. The  $n$ th power of  $-a^3$ , is  $\pm a^{3n}$ .

Here the power will be positive or negative, according as the number which  $n$  represents is even or odd.

223. A FRACTION IS INVOLVED, BY INVOLVING BOTH THE NUMERATOR, AND THE DENOMINATOR.

1. The square of  $\frac{a}{b}$  is  $\frac{a^2}{b^2}$ . For, by the rule for the multiplication of fractions, (Art. 155.)

$$\frac{a}{b} \times \frac{a}{b} = \frac{aa}{bb} = \frac{a^2}{b^2}.$$

2. The 2d, 3d, and  $n$ th powers of  $\frac{1}{a}$ , are  $\frac{1}{a^2}$ ,  $\frac{1}{a^3}$  and  $\frac{1}{a^n}$ .

3. The cube of  $\frac{2xr^2}{3y}$ , is  $\frac{8x^3 r^6}{27y^3}$ .

4. The  $n$ th power of  $\frac{x^2 r}{ay^m}$ , is  $\frac{x^{2n} r^n}{a^n y^{mn}}$ .

5. The square of  $\frac{-a^3 \times (d+m)}{(x+1)^3}$ , is  $\frac{a^6 \times (d+m)^2}{(x+1)^6}$ .

6. The cube of  $\frac{-a^{-1}}{x^{-3}}$ , is  $\frac{-a^{-3}}{x^{-9}}$ . (Art. 221.)

224. Examples of *binomials*, in which one of the terms is a fraction.

1. Find the square of  $x + \frac{1}{2}$ , and  $x - \frac{1}{2}$ , as in art. 214.

$$\begin{array}{r} x + \frac{1}{2} \\ x + \frac{1}{2} \\ \hline \end{array}$$

$$\begin{array}{r} x^2 + \frac{1}{2}x \\ \quad + \frac{1}{2}x + \frac{1}{4} \\ \hline \end{array}$$

$$x^2 + x + \frac{1}{4}.$$

$$\begin{array}{r} x - \frac{1}{2} \\ x - \frac{1}{2} \\ \hline \end{array}$$

$$\begin{array}{r} x^2 - \frac{1}{2}x \\ \quad - \frac{1}{2}x + \frac{1}{4} \\ \hline \end{array}$$

$$x^2 - x + \frac{1}{4}.$$

2. The square of  $a + \frac{2}{3}$ , is  $a^2 + \frac{4a}{3} + \frac{4}{9}$ .

3. The square of  $x + \frac{b}{2}$ , is  $x^2 + bx + \frac{b^2}{4}$ .

4. The square of  $x - \frac{b}{m}$ , is  $x^2 - \frac{2bx}{m} + \frac{b^2}{m^2}$ .



225. It has been shown, (Art. 165,) that a *fractional coefficient* may be transferred, from the numerator to the denominator of a fraction, or from the denominator to the numerator. By recurring to the scheme of notation for reciprocal powers, (Art. 207,) it will be seen that *any factor* may also be transferred, *if the sign of its index be changed*.

1. Thus, in the fraction  $\frac{ax^{-2}}{y}$ , we may transfer  $x$  from the numerator to the denominator.

$$\text{For } \frac{ax^{-2}}{y} = \frac{a}{y} \times x^{-2} = \frac{a}{y} \times \frac{1}{x^2} = \frac{a}{yx^2}.$$

2. In the fraction  $\frac{a}{by^3}$ , we may transfer  $y$  from the denominator to the numerator.

$$\text{For } \frac{a}{by^3} = \frac{a}{b} \times \frac{1}{y^3} = \frac{a}{b} \times y^{-3} = \frac{ay^{-3}}{b}.$$

$$3. \frac{da^{-4}}{x^3} = \frac{d}{x^3 a^4}.$$

$$4. \frac{b}{ay^n} = \frac{by^{-n}}{a}.$$

226. In the same manner, we may transfer a factor which has a positive index in the numerator, or a negative index in the denominator.

1. Thus  $\frac{ax^3}{b} = \frac{a}{bx^{-3}}$ . For  $x^3$  is the reciprocal of  $x^{-3}$ , (Arts. 205, 207,) that is,  $x^3 = \frac{1}{x^{-3}}$ . Therefore  $\frac{ax^3}{b} = \frac{a}{bx^{-3}}$ .

$$2. \frac{h}{by^{-2}} = \frac{hy^2}{b}.$$

$$3. \frac{ad^2}{xy^{-3}} = \frac{ay^3}{xd^{-2}}.$$

227. Hence, the denominator of any fraction may be entirely removed, or the numerator may be reduced to a unit, without altering the value of the expression.

$$1. \text{ Thus } \frac{a}{b} = \frac{1}{ba^{-1}}, \text{ or } ab^{-1}.$$

$$2. \frac{x^{-2}}{b^{-n}} = \frac{1}{x^2 b^{-n}}, \text{ or } b^n x^{-2}.$$

$$3. \frac{x^4 a^{-m}}{b^n c^{-3}} = \frac{1}{b^n a^m x^{-4} c^{-3}}, \text{ or } c^3 x^4 a^{-m} b^{-n}.$$

### ADDITION AND SUBTRACTION OF POWERS.

228. It is obvious that powers may be added, like other quantities, *by writing them one after another, with their signs* (Art. 69.)

Thus the sum of  $a^3$  and  $b^2$ , is  $a^3 + b^2$ .

And the sum of  $a^2 - b^n$  and  $h^5 - d^4$ , is  $a^2 - b^n + h^5 - d^4$ .

229. The same powers of the same letters are like quantities; (Art. 45,) and their co-efficients may be added or subtracted, as in arts. 72 and 74.

Thus the sum of  $2a^2$  and  $3a^2$ , is  $5a^2$ .

It is as evident that twice the square of  $a$ , and three times the square of  $a$ , are five times the square of  $a$ , as that twice  $a$  and three times  $a$ , are five times  $a$ .

To	$-3x^6y^5$	$3b^m$	$3a^4y^n$	$-5a^3h^6$	$3(a+y)^n$
Add	$-2x^6y^5$	$6b^m$	$-7a^4y^n$	$6a^3h^6$	$4(a+y)^n$
Sum	$-5x^6y^5$	$9b^m$	$-4a^4y^n$	$1a^3h^6$	$7(a+y)^n$

230. But powers of *different letters*, and *different powers* of the *same letter*, must be added by writing them down with their signs.

The sum of  $a^2$  and  $a^3$ , is  $a^2 + a^3$ .

It is evident that the square of  $a$ , and the cube of  $a$ , are neither twice the square of  $a$ , nor twice the cube of  $a$ .

The sum of  $a^3b^n$  and  $3a^5b^6$ , is  $a^3b^n + 3a^5b^6$ .

231. *Subtraction* of powers is to be performed in the same manner as addition, except that the signs of the subtrahend are to be changed according to art. 82.

From	$2a^4$	$-3b^n$	$3h^2b^6$	$a^3b^n$	$5(a-h)^6$
Sub.	$-6a^4$	$4b^n$	$4h^2b^6$	$a^3b^n$	$2(a-h)^6$
Diff.	$+8a^4$	$-7b^n$	$-h^2b^6$	$-2a^3b^n$	$3(a-h)^6$

MULTIPLICATION OF POWERS.

232. Powers may be multiplied, like other quantities, by writing the factors one after another, either with, or without, the sign of multiplication between them. (Art. 93.)

Thus the product of  $a^3$  into  $b^2$ , is  $a^3b^2$ , or  $aaabb$ .

Mult.	$x^{-3}$	$h^2b^{-n}$	$3a^3y^2$	$dh^3x^{-n}$	$a^2b^3y^2$
Into	$a^n$	$a^4$	$-2x$	$4by^4$	$a^3b^2y$
Prod.	$a^m x^{-3}$	$a^4 h^2 b^{-n}$	$-6a^6 xy^2$	$4by^4 dh^3 x^{-n}$	$a^2 b^3 y^2 a^3 b^2 y$

The product in the last example, may be abridged, by bringing together the letters which are repeated.

It will then become  $a^5 b^5 y^3$ .

The reason of this will be evident, by recurring to the series of powers in art. 207, viz.

$a^{+4}, a^{+3}, a^{+2}, a^{+1}, a^0, a^{-1}, a^{-2}, a^{-3}, a^{-4}, \&c.$

Or, which is the same,

$aaaa, aaa, aa, a, 1, \frac{1}{a}, \frac{1}{aa}, \frac{1}{aaa}, \frac{1}{aaaa}, \&c.$



By comparing the several terms with each other, it will be seen that if any two or more of them be multiplied together, their product will be a power whose exponent is the *sum* of the exponents of the factors.

Thus  $a^2 \times a^3 = aa \times aaa = aaaaa = a^5$ .

Here 5, the exponent of the product, is equal to  $2+3$ , the sum of the exponents of the factors.

So  $a^n \times a^m = a^{n+m}$ .

For  $a^n$ , is  $a$  taken for a factor as many times, as there are units in  $n$  ;

And  $a^m$ , is  $a$  taken for a factor as many times, as there are units in  $m$  ;

Therefore the product must be  $a$  taken for a factor as many times as there are units in both  $m$  and  $n$ . Hence,

233. POWERS OF THE SAME ROOT MAY BE MULTIPLIED, BY ADDING THEIR EXPONENTS.

Thus  $a^2 \times a^6 = a^{2+6} = a^8$ . And  $x^3 \times x^2 \times x = x^{3+2+1} = x^6$ .

Mult.	$4a^n$	$3x^4$	$b^2y^3$	$a^2b^3y^2$	$(b+h-y)^n$
Into	$2a^n$	$2x^3$	$b^4y$	$a^3b^2y$	$b+h-y$
Prod.	$8a^{2n}$	$5x^7$	$b^6y^4$	$a^5b^5y^3$	$(b+h-y)^{n+1}$

Mult.  $x^3 + x^2y + xy^2 + y^3$  into  $x-y$ .      Ans.  $x^4 - y^4$ .

Mult.  $4x^2y + 3xy - 1$  into  $2x^2 - x$ .

Mult.  $x^3 + x - 5$  into  $2x^2 + x + 1$ .

234. The rule is equally applicable to powers whose exponents are *negative*.

1. Thus  $a^{-2} \times a^{-3} = a^{-5}$ .      That is  $\frac{1}{aa} \times \frac{1}{aaa} = \frac{1}{aaaaa}$

2.  $y^{-n} \times y^{-m} = y^{-n-m}$ .      That is  $\frac{1}{y^n} \times \frac{1}{y^m} = \frac{1}{y^n y^m}$

3.  $-a^{-2} \times a^{-3} = -a^{-5}$ .      4.  $a^{-2} \times a^3 = a^{3-2} = a^1$ .

5.  $a^{-n} \times a^m = a^{m-n}$ .      6.  $y^{-2} \times y^2 = y^0 = 1$ .

235. If  $a+b$  be multiplied into  $a-b$ , the product will be  $a^2 - b^2$ : (Art. 110,) that is,

THE PRODUCT OF THE SUM AND DIFFERENCE OF TWO QUANTITIES, IS EQUAL TO THE DIFFERENCE OF THEIR SQUARES.

This is another instance of the facility with which *general truths* are demonstrated in algebra. See arts. 23 and 77.

If the sum and difference of the *squares* be multiplied, the product will be equal to the difference of the *fourth powers*, &c.

$$\text{Thus } (a-y) \times (a+y) = a^2 - y^2.$$

$$(a^2 - y^2) \times (a^2 + y^2) = a^4 - y^4.$$

$$(a^4 - y^4) \times (a^4 + y^4) = a^8 - y^8, \text{ \&c.}$$

### DIVISION OF POWERS.

236. Powers may be divided, like other quantities, by rejecting from the dividend a factor equal to the divisor; or by placing the divisor under the dividend, in the form of a fraction.

Thus the quotient of  $a^3b^2$  divided by  $b^2$ , is  $a^3$ . (Art. 116.)

Divide	$9a^3y^4$	$12b^3x^n$	$a^2b + 3a^2y^4$	$d \times (a-h+y)^3$
By	$-3a^3$	$2b^3$	$a^2$	$(a-h+y)^3$
Quot.	$-3y^4$		$b + 3y^4$	$d$

The quotient of  $a^5$  divided by  $a^3$ , is  $\frac{a^5}{a^3}$ . But this is equal to  $a^2$ . For, in the series

$$a^{+4}, a^{+3}, a^{+2}, a^{+1}, a^0, a^{-1}, a^{-2}, a^{-3}, a^{-4}, \text{ \&c.}$$

if any term be divided by another, the index of the quotient will be equal to the *difference* between the index of the dividend, and that of the divisor.

$$\text{Thus } a^5 \div a^3 = \frac{aaaaa}{aaa} = a^2. \quad \text{And } a^m \div a^n = \frac{a^m}{a^n} = a^{m-n}.$$

Hence,

237. A POWER MAY BE DIVIDED BY ANOTHER POWER OF THE SAME ROOT, BY EXTRACTING THE INDEX OF THE DIVISOR FROM THAT OF THE DIVIDEND.

Thus  $y^3 \div y^2 = y^{3-2} = y^1$ . That is  $\frac{yyy}{yy} = y$ .

And  $a^{n+1} \div a = a^{n+1-1} = a^n$ . That is  $\frac{aa^n}{a} = a^n$ .

And  $x^n \div x^n = x^{n-n} = x^0 = 1$ . That is  $\frac{x^n}{x^n} = 1$ .

Divide	$y^{2m}$	$b^6$	$8a^{n+m}$	$a^{n+3}$	$12(b+y)^n$
By	$y^m$	$b^3$	$4a^m$	$a^2$	$3(b+y)^3$
Quot.	$y^2$	$b^3$	$2a^n$	$\frac{1}{a}$	$4(b+y)^{n-8}$

238. The rule is equally applicable to powers whose exponents are *negative*.

1. The quotient of  $a^{-5}$  by  $a^{-3}$ , is  $a^{-2}$ .

That is  $\frac{1}{aaaaa} \div \frac{1}{aaa} = \frac{1}{aaaaa} \times \frac{aaa}{1} = \frac{aaa}{aaaaa} = \frac{1}{aa}$ .

2.  $-x^{-5} \div x^{-3} = -x^{-2}$ . That is  $\frac{1}{-x^5} \div \frac{1}{x^3} = \frac{x^3}{-x^5} = \frac{1}{-x^2}$ .

3.  $h^2 \div h^{-1} = h^{2+1} = h^3$ . That is  $h^2 \div \frac{1}{h} = h^2 \times \frac{h}{1} = h^3$ .

4.  $6a^n \div 2a^{-3} = 3a^{n+3}$ .      5.  $ba^3 \div a = ba^2$ .

6.  $b^3 \div b^5 = b^{3-5} = b^{-2}$ .      7.  $a^4 \div a^7 = a^{-3}$ .

8.  $(a^3 + y^3)^m \div (a^3 + y^3)^n = (a^3 + y^3)^{m-n}$ .

9.  $(b+x)^n \div (b+x) = (b+x)^{n-1}$ .

The multiplication and division of powers by adding and subtracting their indices, should be made very familiar; as they have numerous and important applications, in the higher branches of algebra.

**EXAMPLES OF FRACTIONS CONTAINING POWERS.**

239. In the section on fractions, the following examples were omitted, for the sake of avoiding an anticipation of the subject of powers.

1. Reduce  $\frac{5a^4}{3a^2}$  to lower terms.      Ans.  $\frac{5a^2}{3}$ .

For  $\frac{5a^4}{3a^2} = \frac{5aaaa}{3aa} = \frac{5aa}{3}$ . (Art. 145.)

2. Reduce  $\frac{6x^6}{3x^5}$  to lower terms.      Ans.  $\frac{2x}{1}$  or  $2x$ .

3. Reduce  $\frac{3a^4 + 4a^6}{5a^3}$  to lower terms. Ans.  $\frac{3a + 4a^3}{5}$ .

4. Reduce  $\frac{8a^3y - 12a^2y^2 + 6ay^3}{6a^2y + 4ay^2}$  to lower terms.

Ans.  $\frac{4a^2 - 6ay + 3y^2}{3a + 2y}$  obtained by dividing each term by  $2ay$ .

5. Reduce  $\frac{a^4}{a^3}$  and  $\frac{a^{-3}}{a^{-4}}$ , to a common denominator.

$a^2 \times a^{-4}$  is  $a^{-2}$ , the first numerator. (Art. 146.)

$a^3 \times a^{-3}$  is  $a^0 = 1$ , the second numerator.

$a^3 \times a^{-4}$  is  $a^{-1}$ , the common denominator.

The fractions reduced are therefore  $\frac{a^{-2}}{a^{-1}}$  and  $\frac{1}{a^{-1}}$ .

6. Reduce  $\frac{2a^4}{5a^3}$  and  $\frac{a^2}{a^4}$ , to a common denominator.

Ans.  $\frac{2a^8}{5a^7}$  and  $\frac{5a^5}{5a^7}$ , or  $\frac{2a^3}{5a^2}$  and  $\frac{5}{5a^2}$  (Art. 145.)

7. Multiply  $\frac{3x^2}{4x^3}$  into  $\frac{dx}{2x^4}$ . Ans.  $\frac{3dx^3}{8x^7} = \frac{3d}{8x^4}$ .

8. Multiply  $\frac{a^3 + b}{b^4}$ , into  $\frac{a - b}{3}$ .

9. Multiply  $\frac{a^5 + 1}{x^2}$ , into  $\frac{b^2 - 1}{x + a}$ .

10. Multiply  $\frac{b^4}{a^{-2}}$ , into  $\frac{h^{-3}}{x}$ , and  $\frac{a^n}{y^{-3}}$ .

11. Divide  $\frac{a^4}{y^5}$  by  $\frac{a^3}{y^2}$ . Ans.  $\frac{a^4y^2}{a^3y^3} = \frac{a}{y}$ .

12. Divide  $\frac{a^3 - x^4}{a^2}$ , by  $\frac{x^2 - a^{-2}}{a}$ .

13. Divide  $\frac{b - y^{-1}}{y}$  by  $\frac{a^3 + b^{-4}}{y^3}$ .

14. Divide  $\frac{h^3 - 1}{d^4}$ , by  $\frac{d^n + 1}{h}$ .

## SECTION IX.

## EVOLUTION AND RADICAL QUANTITIES.\*

ART. 240. **I**F a quantity is multiplied into itself, the product is a *power*. On the contrary, if a quantity is resolved into any number of *equal factors*, each of these is a *root* of that quantity.

Thus  $b$  is a root of  $bbb$ ; because  $bbb$  may be resolved into the three equal factors  $b$ , and  $b$ , and  $b$ .

In subtraction, a quantity is resolved into *two parts*.

In division, a quantity is resolved into *two factors*.

In evolution, a quantity is resolved into *equal factors*.

241. A ROOT OF A QUANTITY, THEN, IS A FACTOR, WHICH MULTIPLIED INTO ITSELF A CERTAIN NUMBER OF TIMES, WILL PRODUCE THAT QUANTITY.

The number of times the root must be taken as a factor, to produce the given quantity, is denoted by the name of the root.

Thus 2 is the 4th root of 16; because  $2 \times 2 \times 2 \times 2 = 16$ , where 2 is taken *four* times as a factor, to produce 16.

So  $a^3$  is the square root of  $a^6$ ; for  $a^3 \times a^3 = a^6$ . (Art. 233.)

And  $a^2$  is the cube root of  $a^6$ ; for  $a^2 \times a^2 \times a^2 = a^6$ .

And  $a$  is the 6th root of  $a^6$ ; for  $a \times a \times a \times a \times a \times a = a^6$ .

Powers and roots are correlative terms. If one quantity is a power of another, the latter is a root of the former. As  $b^3$  is the cube of  $b$ ;  $b$  is the cube root of  $b^3$ .

242. There are two methods in use, for expressing the roots of quantities, one by means of the radical sign  $\sqrt{\quad}$ , and the other by a fractional index. The latter is generally to be preferred. But the former has its uses on particular occasions.

\* Newton's Arithmetic, Maclaurin, Emerson, Euler, Saunderson, and Simpson.

When a root is expressed by the radical sign, the sign is placed over the given quantity, in this manner,  $\sqrt{a}$ .

Thus  $\sqrt[2]{a}$  is the 2d or square root of  $a$ .

$\sqrt[3]{a}$  is the 3d or cube root.

$\sqrt[n]{a}$  is the  $n$ th root.

And  $\sqrt[n]{a+y}$  is the  $n$ th root of  $a+y$ .

243. The figure placed over the radical sign, denotes the number of factors into which the given quantity is resolved; in other words, the number of times the root must be taken as a factor, to produce the given quantity.

So that  $\sqrt[2]{a} \times \sqrt[2]{a} = a$ .

And  $\sqrt[3]{a} \times \sqrt[3]{a} \times \sqrt[3]{a} = a$ .

And  $\sqrt[n]{a} \times \sqrt[n]{a} \dots n \text{ times} = a$ .

The figure for the *square* root is commonly omitted;  $\sqrt{a}$  being put for  $\sqrt[2]{a}$ . Whenever, therefore, the radical sign is used without a figure, the square root is to be understood.

244. When a figure or letter is *prefixed* to the radical sign, without any character between them; the two quantities are to be considered as *multiplied* together.

Thus  $2\sqrt{a}$ , is  $2 \times \sqrt{a}$ , that is, 2 multiplied into the root of  $a$ , or, which is the same thing, *twice* the root of  $a$ .

And  $x\sqrt{b}$ , is  $x \times \sqrt{b}$ , or  $x$  times the root of  $b$ .

When no co-efficient is prefixed to the radical sign, 1 is always to be understood;  $\sqrt{a}$  being the same as  $1\sqrt{a}$ , that is, *once* the root of  $a$ .

245. The method of expressing roots by radical signs, has no very apparent connection with the other parts of the scheme of algebraic notation. But the plan of indicating them by *fractional indices*, is derived directly from the mode of expressing *powers* by *integral indices*. To explain this, let  $a^6$  be a given quantity. If the index be divided into any number of equal parts, each of these will be the index of a root of  $a^6$ .

Thus the *square* root of  $a^6$ , is  $a^3$ . For, according to the definition, (Art. 241,) the square root of  $a^6$  is a factor, which multiplied into itself will produce  $a^6$ . But  $a^3 \times a^3 = a^6$ . (Art. 233.) Therefore,  $a^3$  is the square root of  $a^6$ . The index of the given quantity  $a^6$ , is here divided into the two equal parts 3 and 3. Of course, the quantity itself is resolved into the two equal factors  $a^3$  and  $a^3$ .

The *cube* root of  $a^6$  is  $a^2$ . For  $a^2 \times a^2 \times a^2 = a^6$ .

Here the index is divided into *three* equal parts, and the quantity itself resolved into three equal factors.

The square root of  $a^2$  is  $a^1$  or  $a$ . For  $a \times a = a^2$ .

By extending the same plan of notation, *fractional indices* are obtained.

Thus, in taking the square root of  $a^1$  or  $a$ , the index 1 is divided into the two equal parts  $\frac{1}{2}$  and  $\frac{1}{2}$ ; and the root is  $a^{\frac{1}{2}}$ .

On the same principle,

The cube root of  $a$ , is  $a^{\frac{1}{3}} = \sqrt[3]{a}$ .

The  $n$ th root, is  $a^{\frac{1}{n}} = \sqrt[n]{a}$ , &c.

And the  $n$ th root of  $a+x$ , is  $(a+x)^{\frac{1}{n}} = \sqrt[n]{a+x}$ .

246. In all these cases, the denominator of the fractional index, expresses the number of factors into which the given quantity is resolved.

So that  $a^{\frac{1}{3}} \times a^{\frac{1}{3}} \times a^{\frac{1}{3}} = a$ . And  $a^{\frac{1}{n}} \times a^{\frac{1}{n}} \dots n$  times  $= a$ .

247. It follows from this plan of notation, that

$$a^{\frac{1}{2}} \times a^{\frac{1}{2}} = a^{\frac{1}{2} + \frac{1}{2}}. \text{ For } a^{\frac{1}{2} + \frac{1}{2}} = a^1 \text{ or } a.$$

$$a^{\frac{1}{3}} \times a^{\frac{1}{3}} \times a^{\frac{1}{3}} = a^{\frac{1}{3} + \frac{1}{3} + \frac{1}{3}} = a^1, \text{ \&c.}$$

where the multiplication is performed in the same manner, as the multiplication of powers, (Art. 233,) that is, by *adding the indices*.

248. Every root as well as every power of 1 is 1. (Art. 209.) For a root is a factor which multiplied into itself will produce the given quantity. But no factor except 1 can produce 1, by being multiplied into itself.

So that  $1^n$ ,  $1$ ,  $\sqrt{1}$ ,  $\sqrt[n]{1}$ , &c. are all equal.

249. *Negative* indices are used in the notation of roots, as well as of powers. See art. 207.

$$\text{Thus } \frac{1}{a^{\frac{1}{2}}} = a^{-\frac{1}{2}} \quad \frac{1}{a^{\frac{1}{3}}} = a^{-\frac{1}{3}} \quad \frac{1}{a^n} = a^{-\frac{1}{n}}.$$

## POWERS OF ROOTS.

250. It has been shown in what manner any power or root may be expressed by means of an index. The index of a power is a whole number. That of a root is a fraction whose numerator is 1. There is also another class of quantities, which may be considered, either as powers of roots, or roots of powers.

Suppose  $a^{\frac{1}{2}}$  is multiplied into itself, so as to be repeated three times as a factor.

The product  $a^{\frac{1}{2} + \frac{1}{2} + \frac{1}{2}}$  or  $a^{\frac{3}{2}}$  (Art. 247,) is evidently the cube of  $a^{\frac{1}{2}}$ , that is, the cube of the square root of  $a$ . This fractional index denotes, therefore, a *power of a root*. The denominator expresses the root, and the numerator the power. The denominator shows into how many equal factors or roots the given quantity is resolved; and the numerator shows how many of these roots are to be multiplied together.

Thus  $a^{\frac{4}{3}}$  is the 4th power of the cube root of  $a$ .

The denominator shows that  $a$  is resolved into the three factors or roots  $a^{\frac{1}{3}}$ , and  $a^{\frac{1}{3}}$ , and  $a^{\frac{1}{3}}$ . And the numerator shows that four of these are to be multiplied together; which will produce the fourth power of  $a^{\frac{1}{3}}$ ; that is,

$$a^{\frac{1}{3}} \times a^{\frac{1}{3}} \times a^{\frac{1}{3}} \times a^{\frac{1}{3}} = a^{\frac{4}{3}}.$$

251. As  $a^{\frac{3}{2}}$  is a power of a root, so it is a *root of a power*. Let  $a$  be raised to the third power  $a^3$ . The square root of this is  $a^{\frac{3}{2}}$ . For the root of  $a^3$  is a quantity which multiplied into itself will produce  $a^3$ .

But according to art. 247,  $a^{\frac{3}{2}} = a^{\frac{1}{2}} \times a^{\frac{1}{2}} \times a^{\frac{1}{2}}$ ; and this multiplied into itself, (Art. 103,) is

$$a^{\frac{1}{2}} \times a^{\frac{1}{2}} \times a^{\frac{1}{2}} \times a^{\frac{1}{2}} \times a^{\frac{1}{2}} \times a^{\frac{1}{2}} = a^3.$$

Therefore  $a^{\frac{3}{2}}$  is the square root of the cube of  $a$ .

In the same manner, it may be shown that  $a^{\frac{m}{n}}$  is the  $m$ th power of the  $n$ th root of  $a$ ; or the  $n$ th root of the  $m$ th pow-



er: that is, a root of a power is equal to the same power of the same root. For instance, the fourth power of the cube root of  $a$ , is the same, as the cube root of the fourth power of  $a$ .

252. Roots, as well as powers, of the same letter, may be multiplied by *adding their exponents*. (Art. 247.) It will be easy to see, that the same principle may be extended to powers of roots, when the exponents have a common denominator.

$$\text{Thus } a^{\frac{2}{7}} \times a^{\frac{3}{7}} = a^{\frac{2}{7} + \frac{3}{7}} = a^{\frac{5}{7}}.$$

For the first numerator shows how often  $a^{\frac{1}{7}}$  is taken as a factor to produce  $a^{\frac{2}{7}}$ . (Art. 250.)

And the second numerator shows how often  $a^{\frac{1}{7}}$  is taken as a factor to produce  $a^{\frac{3}{7}}$ .

The *sum* of the numerators, therefore, shows how often the root must be taken, for the *product*. (Art. 103.)

$$\text{Or thus, } a^{\frac{2}{7}} = a^{\frac{1}{7}} \times a^{\frac{1}{7}}.$$

$$\text{And } a^{\frac{3}{7}} = a^{\frac{1}{7}} \times a^{\frac{1}{7}} \times a^{\frac{1}{7}}.$$

$$\text{Therefore } a^{\frac{2}{7}} \times a^{\frac{3}{7}} = a^{\frac{1}{7}} \times a^{\frac{1}{7}} \times a^{\frac{1}{7}} \times a^{\frac{1}{7}} \times a^{\frac{1}{7}} = a^{\frac{5}{7}}.$$

253. The value of a quantity is not altered, by applying to it a fractional index whose numerator and denominator are equal.

Thus  $a = a^{\frac{2}{2}} = a^{\frac{3}{3}} = a^{\frac{n}{n}}$ . For the denominator shows that  $a$  is resolved into a certain number of factors; and the numerator shows that all these factors are included in  $a^{\frac{n}{n}}$ .

$$\text{Thus } a^{\frac{3}{3}} = a^{\frac{1}{3}} \times a^{\frac{1}{3}} \times a^{\frac{1}{3}}, \text{ which is equal to } a.$$

$$\text{And } a^{\frac{n}{n}} = a^{\frac{1}{n}} \times a^{\frac{1}{n}} \times a^{\frac{1}{n}} \dots n \text{ times.}$$

On the other hand, when the numerator of a fractional index becomes equal to the denominator, the expression may be rendered more simple by *rejecting* the index.

Instead of  $a^{\frac{n}{n}}$ , we may write  $a$ .

254. The index of a power or root may be exchanged, for any other index of the same value.

$$\text{Instead of } a^{\frac{2}{3}}, \text{ we may put } a^{\frac{4}{6}}.$$

For, in the latter of these expressions,  $a$  is supposed to be resolved into *twice* as many factors as in the former; and the numerator shows that *twice* as many of these factors are to be multiplied together. So that the whole value is not altered.

Thus  $x^{\frac{2}{3}} = x^{\frac{4}{6}} = x^{\frac{6}{9}}$ , &c. that is, the square of the cube root is the same, as the fourth power of the sixth root, the sixth power of the 9th root, &c.

So  $a^2 = a^{\frac{4}{2}} = a^{\frac{6}{3}} = a^{\frac{2n}{n}}$ . For the value of each of these indices is 2. (Art. 135.)

255. From the preceding article, it will be easily seen, that a fractional index may be expressed in *decimals*.

1. Thus  $a^{\frac{1}{2}} = a^{\frac{5}{10}}$  or  $a^{0.5}$ ; that is, the square root is equal to the 5th power of the tenth root.

2.  $a^{\frac{1}{4}} = a^{\frac{25}{100}}$  or  $a^{0.25}$ ; that is, the fourth root is equal to the 25th power of the 100th root.

$$3. a^{\frac{2}{5}} = a^{0.4}$$

$$5. a^{\frac{9}{5}} = a^{1.8}$$

$$4. a^{\frac{7}{2}} = a^{3.5}$$

$$6. a^{\frac{11}{4}} = a^{2.75}$$

In many cases however, the decimal can be only an *approximation* to the true index.

Thus  $a^{\frac{1}{3}} = a^{0.3}$  nearly.

$a^{\frac{1}{3}} = a^{0.33333}$  very nearly.

In this manner, the approximation may be carried to any degree of exactness which is required.

Thus  $a^{\frac{5}{3}} = a^{1.66666}$ .

$a^{\frac{11}{7}} = a^{1.57142}$ .

These decimal indices form a very important class of numbers, called *logarithms*.

It is frequently convenient to vary the notation of powers of roots, by making use of a vinculum, or the radical sign  $\sqrt{\quad}$ . In doing this, we must keep in mind, that the power of a root is the same, as the root of a power; (Art. 251,) and also, that the *denominator* of a fractional exponent expresses a *root*, and the *numerator*, a *power*. (Art. 250.)

Instead, therefore, of  $a^{\frac{2}{3}}$ , we may write  $(a^{\frac{1}{3}})^2$ , or  $(a^2)^{\frac{1}{3}}$ , or  $\sqrt[3]{a^2}$ .

The first of these three forms, denotes the square of the cube root of  $a$ ; and each of the two last, the cube root of the square of  $a$ .

$$\text{So } a^{\frac{m}{n}} = a^{\frac{1}{n}|^m} = a^{\frac{m}{n}} = \sqrt[n]{a^m}$$

$$\text{And } (bx)^{\frac{3}{4}} = (b^3x^3)^{\frac{1}{4}} = \sqrt[4]{b^3x^3}.$$

$$\text{And } \sqrt{a+y^3} = \sqrt{a+y^3}|^{\frac{1}{5}} = \sqrt[5]{a+y^3}.$$

## EVOLUTION.

257. Evolution is the opposite of involution. One is finding a *power* of a quantity, by multiplying it into itself. The other is finding a *root*, by resolving a quantity into equal factors. A quantity is resolved into any number of equal factors, by dividing its *index* into as many *equal parts*, (Art. 245.)

Evolution may be performed, then, by the following general rule;

DIVIDE THE INDEX OF THE QUANTITY, BY THE NUMBER EXPRESSING THE ROOT TO BE FOUND.

Or, place over the quantity the radical sign belonging to the required root.

1. Thus the cube root of  $a^6$  is  $a^2$ . For  $a^2 \times a^2 \times a^2 = a^6$ .

Here 6, the index of the given quantity, is divided by 3, the number expressing the cube root.

2. The cube root of  $a$  or  $a^1$ , is  $a^{\frac{1}{3}}$  or  $\sqrt[3]{a}$ .

For  $a^{\frac{1}{3}} \times a^{\frac{1}{3}} \times a^{\frac{1}{3}}$ , or  $\sqrt[3]{a} \times \sqrt[3]{a} \times \sqrt[3]{a} = a$ . (Arts. 243, 246)

3. The 5th root of  $ab$ , is  $(ab)^{\frac{1}{5}}$  or  $\sqrt[5]{ab}$ .

4. The  $n$ th root of  $a^2$ , is  $a^{\frac{2}{n}}$  or  $\sqrt[n]{a^2}$ .

5. The 7th root of  $2d-x$ , is  $(2d-x)^{\frac{1}{7}}$  or  $\sqrt[7]{2d-x}$ .

6. The 5th root of  $a^{-x}|^3$ , is  $a^{-x}|^{\frac{3}{5}}$  or  $\sqrt[5]{a^{-x}|^3}$ .

7. The cube root of  $a^{\frac{1}{2}}$ , is  $a^{\frac{1}{6}}$ . (Art. 163.)

8. The 4th root of  $a^{-1}$  is  $a^{-\frac{1}{4}}$ .

9. The cube root of  $a^{\frac{2}{3}}$  is  $a^{\frac{2}{9}}$ .

10. The  $n$ th root of  $x^m$ , is  $x^{\frac{m}{n}}$ .

258. According to the rule just given, the cube root of the square root is found, by dividing the index  $\frac{1}{2}$  by 3, as in example 7th. But instead of dividing by 3, we may multiply by  $\frac{1}{3}$ . For  $\frac{1}{2} \div 3 = \frac{1}{2} \div \frac{3}{1} = \frac{1}{2} \times \frac{1}{3}$ . (Art. 162.)

So  $\frac{1}{m} \div n = \frac{1}{m} \times \frac{1}{n}$ . Therefore the  $m$ th root of the  $n$ th root of  $a$  is equal to  $a^{\frac{1}{n} \times \frac{1}{m}}$ .

That is,  $a^{\frac{1}{n} \times \frac{1}{m}} = a^{\frac{1}{n}} \times a^{\frac{1}{m}} = a^{\frac{1}{nm}}$ .

Here the two fractional indices are reduced to one by multiplication.

It is sometimes necessary to reverse this process; to resolve an index into two factors.

Thus  $x^{\frac{1}{8}} = x^{\frac{1}{4}} \times \frac{1}{2} = x^{\frac{1}{4} \times \frac{1}{2}}$ . That is, the 8th root of  $x$  is equal to the square root of the 4th root.

So  $a+b \Big|^{m \times n} = a+b \Big|^{m} \times \frac{1}{n} = a+b \Big|^{m \times \frac{1}{n}}$ .

It may be necessary to observe, that resolving the index into factors, is not the same as resolving the quantity into factors. The latter is effected, by dividing the index into parts.

259. The rule in art. 257, may be applied to every case in evolution. But when the quantity whose root is to be found, is composed of several factors, there will frequently be an advantage in taking the root of each of the factors separately.

This is done upon the principle, that the root of the product of several factors, is equal to the product of their roots.

Thus  $\sqrt{ab} = \sqrt{a} \times \sqrt{b}$ . For each member of the equation, if involved, will give the same power.

The square of  $\sqrt{ab}$  is  $ab$ . (Art. 241.)

The square of  $\sqrt{a} \times \sqrt{b}$ , is  $\sqrt{a} \times \sqrt{a} \times \sqrt{b} \times \sqrt{b}$ . (Art. 102.)

But  $\sqrt{a} \times \sqrt{a} = a$ . (Art. 241.) And  $\sqrt{b} \times \sqrt{b} = b$ .

Therefore the square of  $\sqrt{a} \times \sqrt{b} = \sqrt{a} \times \sqrt{a} \times \sqrt{b} \times \sqrt{b} = ab$ , which is also the square of  $\sqrt{ab}$ .

On the same principle,  $(ab)^{\frac{1}{n}} = a^{\frac{1}{n}} b^{\frac{1}{n}}$ .

When, therefore, a quantity consists of several factors, we may either extract the root of the whole together; or we may find the root of the factors separately, and then multiply them into each other.

- Ex. 1. The cube root of  $xy$ , is either  $(xy)^{\frac{1}{3}}$  or  $x^{\frac{1}{3}} y^{\frac{1}{3}}$ .
2. The 5th root of  $3y$ , is  $\sqrt[5]{3y}$  or  $\sqrt[5]{3} \times \sqrt[5]{y}$ .
3. The 6th root of  $abh$ , is  $(abh)^{\frac{1}{6}}$ , or  $a^{\frac{1}{6}} b^{\frac{1}{6}} h^{\frac{1}{6}}$ .
4. The cube root of  $8b$ , is  $(8b)^{\frac{1}{3}}$ , or  $2b^{\frac{1}{3}}$ .
5. The  $n$ th root of  $x^ny$ , is,  $(x^ny)^{\frac{1}{n}}$  or  $xy^{\frac{1}{n}}$ .

260. THE ROOT OF A FRACTION IS EQUAL TO THE ROOT OF THE NUMERATOR DIVIDED BY THE ROOT OF THE DENOMINATOR.

1. Thus the square root of  $\frac{a}{b} = \frac{a^{\frac{1}{2}}}{b^{\frac{1}{2}}}$ . For  $\frac{a^{\frac{1}{2}}}{b^{\frac{1}{2}}} \times \frac{a^{\frac{1}{2}}}{b^{\frac{1}{2}}} = \frac{a}{b}$ .
2. So the  $n$ th root of  $\frac{a}{b} = \frac{a^{\frac{1}{n}}}{b^{\frac{1}{n}}}$ . For  $\frac{a^{\frac{1}{n}}}{b^{\frac{1}{n}}} \times \frac{a^{\frac{1}{n}}}{b^{\frac{1}{n}}} \cdots n \text{ times} = \frac{a}{b}$ .
3. The square root of  $\frac{x}{ay}$ , is  $\frac{\sqrt{x}}{\sqrt{ay}}$ . 4.  $\sqrt{\frac{ah}{xy}} = \frac{\sqrt{ah}}{\sqrt{xy}}$ .

261. For determining what *sign* to prefix to a root, it is important to observe, that

AN ODD ROOT OF ANY QUANTITY HAS THE SAME SIGN AS THE QUANTITY ITSELF;

AN EVEN ROOT OF AN AFFIRMATIVE QUANTITY IS AMBIGUOUS;

AN EVEN ROOT OF A NEGATIVE QUANTITY IS IMPOSSIBLE.

That the 3d, 5th, 7th, or any other *odd* root of a quantity must have the same sign as the quantity itself, is evident from art. 219.

262. But an *even* root of an *affirmative* quantity, may be either affirmative or negative. For the quantity may be produced from the one, as well as from the other. (Art. 219.)

Thus the square root of  $a^2$  is  $+a$  or  $-a$ .

An even root of an affirmative quantity is, therefore, said to be *ambiguous*, and is marked with both  $+$  and  $-$ .

Thus the square root of  $3b$ , is  $\pm \sqrt{3b}$ .

The 4th root of  $x$ , is  $\pm x^{\frac{1}{4}}$ .

The ambiguity does not exist, however, when, from the nature of the case, or a previous multiplication, it is known whether the power has actually been produced from a positive, or from a negative quantity. See art. 299.

263. But no even root of a *negative* quantity can be found.

The square root of  $-a^2$  is neither  $+a$  nor  $-a$ .

For  $+a \times +a = +a^2$ . And  $-a \times -a = +a^2$  also.

An even root of a negative quantity is, therefore, said to be *impossible* or *imaginary*.

There are purposes to be answered, however, by applying the radical sign to negative quantities. The expression  $\sqrt{-a}$  is often to be found in algebraic processes. For, although we are unable to assign it a rank, among either positive or negative quantities; yet we know that when multiplied into itself, its product is  $-a$ , because  $\sqrt{-a}$  is by notation a *root* of  $-a$ , that is, a quantity which multiplied into itself produces  $-a$ .

This may, at first view, seem to be an exception to the general rule that the product of two negatives is affirmative. But it is to be considered, that  $\sqrt{-a}$  is not itself a negative quantity, but the *root* of a negative quantity.

The mark of subtraction here, must not be confounded with that which is *prefixed* to the radical sign. The expression  $\sqrt{-a}$  is not equivalent to  $-\sqrt{a}$ . The former is a root of  $-a$ ; but the latter is a root of  $+a$ :

$$\text{For } -\sqrt{a} \times -\sqrt{a} = +\sqrt{aa} = a.$$

The root of  $-a$ , however, may be *ambiguous*. It may be either  $+\sqrt{-a}$ , or  $-\sqrt{-a}$ .

One of the uses of imaginary expressions, is to indicate an impossible or absurd supposition, in the statement of a problem. Suppose it be required to divide the number 14 into two such parts, that their product shall be 60. If one of the parts be  $x$ , the other will be  $14-x$ . And by the supposition

$$x \times (14-x) = 60, \text{ or } 14x - x^2 = 60.$$

This reduced, by the rules in the following section, will give

$$x = 7 + \sqrt{-11}.$$

As the value of  $x$  is here found to contain an imaginary expression, we infer that there is an inconsistency in the statement of the problem: that the number 14 cannot be divided into any two parts whose product shall be 60.\*

264. The methods of extracting the roots of *compound* quantities are to be considered in a future section. But there is one class of these, the squares of *binomial* and *residual* quantities, which it will be proper to attend to in this place. It has been shown, (Art. 214,) that the square of a binomial quantity consists of *three terms*, two of which are complete powers, and the other is a double product of the roots of these powers. The square of  $a+b$ , for instance, is

$$a^2 + 2ab + b^2,$$

two terms of which,  $a^2$  and  $b^2$ , are complete powers, and  $2ab$  is twice the product of  $a$  into  $b$ , that is, of the root of  $a^2$  into the root of  $b^2$ .

Whenever, therefore, we meet with a quantity of this description, we may know that its square root is a binomial; and this may be found, by taking the root of the two terms which are complete powers, and connecting them by the sign  $+$ . The other term disappears in the root. Thus, to find the square root of

$$x^2 + 2xy + y^2,$$

take the root of  $x^2$ , and the root of  $y^2$ , and connect them by the sign  $+$ . The binomial root will then be  $x+y$ .

In a *residual* quantity, the double product has the sign  $-$  prefixed, instead of  $+$ . The square of  $a-b$ , for instance, is  $a^2 - 2ab + b^2$ . (Art. 214.) And to obtain the root of a quantity of this description, we have only to take the roots of the two complete powers, and connect them by the sign  $-$ . Thus the square root of  $x^2 - 2xy + y^2$  is  $x-y$ . Hence,

265. TO EXTRACT A BINOMIAL OR RESIDUAL SQUARE ROOT, TAKE THE ROOTS OF THE TWO TERMS WHICH ARE COMPLETE POWERS, AND CONNECT THEM BY THE SIGN, WHICH IS PREFIXED TO THE OTHER TERM.

Ex. 1. Find the root of  $x^2 + 2x + 1$ .

The two terms which are complete powers are  $x^2$  and 1.

Their roots are  $x$  and 1. (Art. 248.)

The binomial root is, therefore,  $x+1$ .

\* See Note F.

2. The square root of  $x^2 - 2x + 1$ , is  $x - 1$ . (Art. 214.)
3. The square root of  $a^2 + a + \frac{1}{4}$ , is  $a + \frac{1}{2}$ . (Art. 224.)
4. The square root of  $a^2 + \frac{4}{3}a + \frac{4}{9}$ , is  $a + \frac{2}{3}$ .
5. The square root of  $a^2 + ab + \frac{b^2}{4}$ , is  $a + \frac{b}{2}$ .
6. The square root of  $a^2 + \frac{2ab}{c} + \frac{b^2}{c^2}$ , is  $a + \frac{b}{c}$ .

266. A ROOT WHOSE VALUE CANNOT BE EXACTLY EXPRESSED IN NUMBERS, IS CALLED A SURD.

Thus  $\sqrt{2}$  is a surd, because the square root of 2 cannot be expressed in numbers, with perfect exactness.

In decimals, it is 1.41421356 nearly.

But though we are unable to assign the value of such a quantity *when taken alone*, yet by multiplying it into itself, or by combining it with other quantities, we may produce expressions whose value can be determined. There is therefore a system of rules generally appropriated to surds. But as all quantities whatever, when under the same radical sign, or having the same index, may be treated in nearly the same manner; it will be most convenient to consider them together, under the general name of *Radical Quantities*; understanding by this term, every quantity which is found under a radical sign, or which has a fractional index.

267. Every quantity which is not a surd, is said to be *rational*. But for the purpose of distinguishing between radicals and other quantities, the term rational will be applied, in this section, to those only which do not appear under a radical sign, and which have not a fractional index.

### REDUCTION OF RADICAL QUANTITIES.

268. Before entering on the consideration of the rules for the addition, subtraction, multiplication, and division of radical quantities, it will be necessary to attend to the methods of reducing them from one form to another.

*First*, to reduce a *rational* quantity to the form of a radical;

RAISE THE QUANTITY TO A POWER OF THE SAME NAME AS THE GIVEN ROOT, AND THEN APPLY THE CORRESPONDING RADICAL SIGN OR INDEX.



Ex. 1. Reduce  $a$  to the form of the  $n$ th root.

The  $n$ th power of  $a$  is  $a^n$ . (Art. 211.)

Over this place the radical sign, and it becomes  $\sqrt[n]{a^n}$ .

It is thus reduced to the form of a radical quantity, without any alteration of its value. For  $\sqrt[n]{a^n} = a^{\frac{n}{n}} = a$ .

2. Reduce 4 to the form of the cube root.

$$\text{Ans. } \sqrt[3]{64} \text{ or } (64)^{\frac{1}{3}}.$$

3. Reduce  $3a$  to the form of the 4th root.

$$\text{Ans. } \sqrt[4]{81a^4}.$$

4. Reduce  $\frac{1}{3}ab$  to the form of the square root.

$$\text{Ans. } \left(\frac{1}{9}a^2b^2\right)^{\frac{1}{2}}.$$

5. Reduce  $3 \times \sqrt{a-x}$  to the form of the cube root.

$$\text{Ans. } \sqrt[3]{27 \times \sqrt{a-x}^3}. \text{ See art. 212.}$$

6. Reduce  $a^2$  to the form of the cube root.

The cube of  $a^2$  is  $a^6$ . (Art. 220.)

And the cube root of  $a^6$  is  $\sqrt[3]{a^6} = a^{\frac{6}{3}}$ .

In cases of this kind, where a *power* is to be reduced to the form of the  $n$ th root, it must be raised to the  $n$ th power, not of the *given letter*, but of the *power* of the letter.

Thus in the example,  $a^6$  is the cube; not of  $a$ , but of  $a^2$ .

7. Reduce  $a^3b^4$  to the form of the square root.

8. Reduce  $a^m$  to the form of the  $n$ th root.

269. *Secondly*, to reduce quantities which have different indices, to others of the same value having a *common index*;

1. Reduce the indices to a common denominator;

2. Involve each quantity, to the power expressed by the numerator of its reduced index.

3. Take the root denoted by the common denominator.

Ex. 1. Reduce  $a^{\frac{1}{4}}$  and  $b^{\frac{1}{6}}$  to a common index.

1st. The indices  $\frac{1}{4}$  and  $\frac{1}{6}$  reduced to a common denominator, are  $\frac{3}{12}$  and  $\frac{2}{12}$ . (Art. 146.)

2d. The quantities  $a$  and  $b$  involved to the powers expressed by the two numerators, are  $a^3$  and  $b^2$ .

3d. The root denoted by the common denominator is  $\sqrt[12]{\phantom{x}}$ .  
The answer, then, is  $\overline{a^3|12}^{\frac{1}{2}}$  and  $\overline{b^2|12}^{\frac{1}{2}}$ .

The two quantities are thus reduced to a common index, without any alteration in their values.

For by art. 254,  $a^{\frac{1}{4}} = a^{\frac{3}{12}}$ , which by art. 258,  $= \overline{a^3|12}^{\frac{1}{2}}$ .

And universally  $a^{\frac{1}{n}} = a^{\frac{m}{mn}} = \overline{a^m|mn}^{\frac{1}{n}}$ .

2. Reduce  $a^{\frac{1}{2}}$  and  $bx^{\frac{2}{3}}$  to a common index.

The indices reduced to a common denominator are  $\frac{3}{6}$  and  $\frac{4}{6}$ .

The quantities, then, are  $a^{\frac{3}{6}}$  and  $(bx)^{\frac{4}{6}}$ , or  $\overline{a^3|6}$  and  $\overline{b^4x^4|6}$ .

3. Reduce  $a^2$  and  $b^{\frac{1}{n}}$ . Ans.  $\overline{a^{2n}|n}$  and  $b^{\frac{1}{n}}$ .

4. Reduce  $x^{\frac{1}{n}}$  and  $y^{\frac{1}{m}}$ . Ans.  $\overline{x^m|mn}$  and  $\overline{y^n|mn}$ .

5. Reduce  $2^{\frac{1}{2}}$  and  $3^{\frac{1}{3}}$ . Ans.  $8^{\frac{1}{6}}$  and  $9^{\frac{1}{6}}$ .

6. Reduce  $(a+b)^2$  and  $(x-y)^{\frac{2}{3}}$ . Ans.  $\overline{a+b|3}^{\frac{2}{3}}$  and  $\overline{x-y|3}^{\frac{2}{3}}$ .

7. Reduce  $a^{\frac{1}{3}}$  and  $b^{\frac{1}{5}}$ .      8. Reduce  $x^{\frac{2}{3}}$  and  $5^{\frac{1}{2}}$ .

270. When it is required to reduce a quantity to a *given* index;

Divide the index of the quantity by the given index, place the quotient over the quantity, and set the given index over the whole.

This is merely resolving the original index into two factors, according to art 258.

Ex. 1. Reduce  $a^{\frac{1}{6}}$  to the index  $\frac{1}{2}$ .

By art. 162,  $\frac{1}{6} \div \frac{1}{2} = \frac{1}{6} \times \frac{2}{1} = \frac{2}{6} = \frac{1}{3}$ .

This is the index to be placed over  $a$ , which then becomes  $a^{\frac{1}{3}}$ ; and the given index set over this makes it  $\overline{a^{\frac{1}{3}}|2}^{\frac{1}{2}}$ , the answer:

2. Reduce  $a^2$  and  $x^{\frac{3}{2}}$ , to the common index  $\frac{1}{3}$ .

$2 \div \frac{1}{3} = 2 \times 3 = 6$ , the first index      }  
 $\frac{3}{2} \div \frac{1}{3} = \frac{3}{2} \times 3 = \frac{9}{2}$ , the second index      }

Therefore  $(a^6)^{\frac{1}{3}}$  and  $(x^{\frac{9}{2}})^{\frac{1}{3}}$  are the quantities required.

3. Reduce  $4^{\frac{1}{2}}$  and  $3^{\frac{1}{3}}$ , to the common index  $\frac{1}{6}$ .

Answer.  $(4^3)^{\frac{1}{6}}$  and  $(3^2)^{\frac{1}{6}}$ .

271. *Thirdly*, to remove a part of a root from under the radical sign;

If the quantity can be resolved into two factors, one of which is an exact power of the same name with the root; FIND THE ROOT OF THIS POWER, AND PREFIX IT TO THE OTHER FACTOR, WITH THE RADICAL SIGN BETWEEN THEM.

This rule is founded on the principle, that the root of the *product* of two factors is equal to the product of their roots. (Art. 259.)

It will generally be best to resolve the radical quantity into such factors, that one of them shall be the *greatest* power which will divide the quantity without a remainder. If there is no exact power which will divide the quantity, the reduction cannot be made.

Ex. 1. Remove a factor from  $\sqrt{8}$ .

The greatest square which will divide 8 is 4.

We may then resolve 8 into the factors 4 and 2. For  $4 \times 2 = 8$ .

The root of this product is equal to the product of the roots of its factors; that is,  $\sqrt{8} = \sqrt{4} \times \sqrt{2}$ .

But  $\sqrt{4} = 2$ . Instead of  $\sqrt{4}$ , therefore, we may substitute its equal 2. We then have  $2 \times \sqrt{2}$  or  $2\sqrt{2}$ .

This is commonly called reducing a radical quantity to its *most simple terms*. But the learner may not perhaps at once perceive, that  $2\sqrt{2}$  is a more simple expression than  $\sqrt{8}$ .

2. Reduce  $\sqrt{a^2x}$ . Ans.  $\sqrt{a^2} \times \sqrt{x} = a \times \sqrt{x} = a\sqrt{x}$ .

3. Reduce  $\sqrt{18}$ . Ans.  $\sqrt{9 \times 2} = \sqrt{9} \times \sqrt{2} = 3\sqrt{2}$ .

4. Reduce  $\sqrt[3]{64b^3c}$ . Ans.  $\sqrt[3]{64b^3} \times \sqrt[3]{c} = 4b\sqrt[3]{c}$ .

5. Reduce  $\sqrt[4]{\frac{a^4b}{c^5d}}$  Ans.  $\frac{a}{c} \sqrt[4]{\frac{b}{cd}}$  (Art. 260.)

6. Reduce  $\sqrt[n]{a^nb}$ . Ans.  $a \sqrt[n]{b}$ , or  $ab^{\frac{1}{n}}$ .

7. Reduce  $(a^3 - a^2b)^{\frac{1}{2}}$ . Ans.  $a(a-b)^{\frac{1}{2}}$ .

8. Reduce  $(54a^3b)^{\frac{1}{3}}$ . Ans.  $3a^2(2b)^{\frac{1}{3}}$ .

9. Reduce  $\sqrt{98a^2x}$ . 10. Reduce  $\sqrt[3]{a^3 + a^2b^2}$ .

272. By a contrary process, the co-efficient of a radical quantity may be introduced under the radical sign.

1. Thus  $a \sqrt[n]{b} = \sqrt[n]{a^n b}$ .

For  $a = \sqrt[n]{a^n}$  or  $a^{\frac{n}{n}}$ . Art. 253.) And  $\sqrt[n]{a^n} \times \sqrt[n]{b} = \sqrt[n]{a^n b}$ .

Here the co-efficient  $a$  is first raised to a power of the same name as the radical part, and is then introduced as a factor under the radical sign.

2.  $a(x-b)^{\frac{1}{3}} = (a^3 \times \overline{x-b})^{\frac{1}{3}} = (a^3 x - a^3 b)^{\frac{1}{3}}$ .

3.  $2ab(2ab^2)^{\frac{1}{3}} = (16a^4 b^5)^{\frac{1}{3}}$

4.  $\frac{a}{b} \left( \frac{b^2 c}{a^2 + b^2} \right)^{\frac{1}{2}} = \left( \frac{a^2 b^2 c}{a^2 b^2 + b^4} \right)^{\frac{1}{2}}$ .

#### ADDITION AND SUBTRACTION OF RADICAL QUANTITIES.

273. Radical quantities may be added like rational quantities, by writing them one after another with their signs. (Art. 69.)

Thus the sum of  $\sqrt{a}$  and  $\sqrt{b}$ , is  $\sqrt{a} + \sqrt{b}$ .

And the sum of  $a^{\frac{1}{2}} - h^{\frac{1}{3}}$  and  $x^{\frac{1}{4}} - y^{\frac{1}{n}}$ , is  $a^{\frac{1}{2}} - h^{\frac{1}{3}} + x^{\frac{1}{4}} - y^{\frac{1}{n}}$ .

But in many cases, several terms may be reduced to one; as in arts. 72 and 74.

The sum of  $2\sqrt{a}$  and  $3\sqrt{a}$  is  $2\sqrt{a} + 3\sqrt{a} = 5\sqrt{a}$ .

For it is evident that twice the root of  $a$ , and three times the root of  $a$ , are five times the root of  $a$ . Hence,

274. When the quantities to be added have the same radical part, under the same radical sign or index; *add the rational parts, and to the sum annex the RADICAL PARTS.*

If no rational quantity is prefixed to the radical sign, 1 is always to be understood. (Art. 244.)

To	$2\sqrt{ay}$	$5\sqrt{a}$	$3(x+h)^{\frac{1}{7}}$	$5bh^{\frac{1}{6}}$	$a\sqrt{b-h}$
Add	$\sqrt{ay}$	$-2\sqrt{a}$	$4(x+h)^{\frac{1}{7}}$	$7bh^{\frac{1}{6}}$	$y\sqrt{b-h}$
Sum	$3\sqrt{ay}$		$7(x+h)^{\frac{1}{7}}$		$(a+y)\sqrt{b-h}$

275. If the radical parts are originally different, they may sometimes be made alike, by the reductions in the preceding articles.

1. Add  $\sqrt{8}$  to  $\sqrt{50}$ . Here the radical parts are not the same. But by the reduction in art. 271,  $\sqrt{8}=2\sqrt{2}$ , and  $\sqrt{50}=5\sqrt{2}$ . The sum then is  $7\sqrt{2}$ .

2. Add  $\sqrt{16b}$  to  $\sqrt{4b}$ . Ans.  $4\sqrt{b}+2\sqrt{b}=6\sqrt{b}$ .

3. Add  $\sqrt{a^2x}$  to  $\sqrt{b^2x}$ . Ans.  $a\sqrt{x}+b\sqrt{x}=(a+b)\sqrt{x}$ .

4. Add  $(36a^2y)^{\frac{1}{2}}$  to  $(25y)^{\frac{1}{2}}$ . Ans.  $(6a+5)\sqrt{y}$ .

5. Add  $\sqrt{18a}$  to  $3\sqrt{2a}$ .

276. But if the radical parts, after reduction, are *different*, or have different *exponents*, they cannot be united in the same term; and must be added by writing them one after the other.

The sum of  $3\sqrt{b}$  and  $2\sqrt{a}$ , is  $3\sqrt{b}+2\sqrt{a}$ .

It is manifest that three times the root of  $b$ , and twice the root of  $a$ , are neither five times the root of  $b$ , nor five times the root of  $a$ , unless  $b$  and  $a$  are equal.

The sum of  $\sqrt[2]{a}$  and  $\sqrt[3]{a}$ , is  $\sqrt[2]{a}+\sqrt[3]{a}$ .

The *square* root of  $a$ , and the *cube* root of  $a$ , are neither twice the square root, nor twice the cube root of  $a$ .

277. *Subtraction* of radical quantities is to be performed in the same manner as addition, except that the signs in the subtrahend are to be changed according to art. 82.

From	$\sqrt{ay}$	$4\sqrt[n]{a+x}$	$3h^{\frac{1}{3}}$	$a(x+y)$	$-a^{-\frac{1}{n}}$
Sub.	$3\sqrt{ay}$	$3\sqrt[n]{a+x}$	$-5h^{\frac{1}{3}}$	$b(x+y)$	$-2a^{-\frac{1}{n}}$
Diff.	$-2\sqrt{ay}$		$8h^{\frac{1}{3}}$		$a^{-\frac{1}{n}}$

From  $\sqrt{50}$ , subtract  $\sqrt{8}$ . Ans.  $5\sqrt{2}-2\sqrt{2}=3\sqrt{2}$ . (Art. 275.)

From  $\sqrt[3]{b^4y}$ , subtract  $\sqrt[3]{by^4}$ . Ans.  $(b-y)\sqrt[3]{by}$ .

From  $\sqrt{x}$ , subtract  $\sqrt[5]{x}$ .

### MULTIPLICATION OF RADICAL QUANTITIES.

278. Radical quantities may be multiplied, like other

quantities, by writing the factors one after another, either with or without the sign of multiplication between them. (Art. 93.)

Thus the product of  $\sqrt{a}$  into  $\sqrt{b}$ , is  $\sqrt{a \times b}$ .

The product of  $h^{\frac{1}{3}}$  into  $y^{\frac{1}{2}}$  is  $h^{\frac{1}{3}}y^{\frac{1}{2}}$ .

But it is often expedient to bring the factors under the same radical sign. This may be done, if they are first reduced to a common index.

Thus  $\sqrt[n]{x} \times \sqrt{y} = \sqrt[n]{xy}$ . For the root of the product of several factors is equal to the product of their roots. (Art. 259.) Hence,

279. QUANTITIES UNDER THE SAME RADICAL SIGN OR INDEX, MAY BE MULTIPLIED TOGETHER LIKE RATIONAL QUANTITIES, THE PRODUCT BEING PLACED UNDER THE COMMON RADICAL SIGN OR INDEX.\*

Multiply  $\sqrt[3]{x}$  into  $\sqrt[3]{y}$ , that is,  $x^{\frac{1}{3}}$  into  $y^{\frac{1}{3}}$ .

The quantities reduced to the same index, (Art. 269,) are  $(x^3)^{\frac{1}{6}}$ , and  $(y^2)^{\frac{1}{6}}$ , and their product is  $(x^3y^2)^{\frac{1}{6}} = \sqrt[6]{x^3y^2}$ .

Mult.	$\sqrt{a + m}$	$\sqrt{dx}$	$a^{\frac{3}{2}}$	$(a+y)^{\frac{1}{n}}$	$a^{\frac{1}{m}}$
Into	$\sqrt{a - m}$	$\sqrt{hy}$	$x^{\frac{1}{2}}$	$(b+h)^{\frac{1}{n}}$	$x^{\frac{1}{n}}$
Prod.	$\sqrt{a^2 - m^2}$		$(a^3x)^{\frac{1}{2}}$		$(a^nx^m)^{\frac{1}{mn}}$

Multiply  $\sqrt{8xb}$  into  $\sqrt{2xb}$ . Prod.  $\sqrt{16x^2b^2} = 4xb$ .

In this manner the product of radical quantities often becomes *rational*.

Thus the product of  $\sqrt{2}$  into  $\sqrt{18} = \sqrt{36} = 6$ .

And the product of  $(a^2y^3)^{\frac{1}{4}}$  into  $(a^2y)^{\frac{1}{4}} = (a^4y^4)^{\frac{1}{4}} = ay$ .

280. ROOTS OF THE SAME LETTER OR QUANTITY MAY BE MULTIPLIED, BY ADDING THEIR FRACTIONAL EXPONENTS.

The exponents, like all other fractions, must be reduced to a common denominator, before they can be united in one term. (Art. 148.)

\* The case of an *imaginary* root of a negative quantity may be considered an exception. (Art. 263.)

Thus  $a^{\frac{1}{2}} \times a^{\frac{1}{3}} = a^{\frac{1}{2} + \frac{1}{3}} = a^{\frac{3}{6} + \frac{2}{6}} = a^{\frac{5}{6}}$ .

The values of the roots are not altered, by reducing their indices to a common denominator. (Art. 254.)

Therefore the first factor  $a^{\frac{1}{2}} = a^{\frac{3}{6}}$   
 And the second  $a^{\frac{1}{3}} = a^{\frac{2}{6}}$

But  $a^{\frac{3}{6}} = a^{\frac{1}{6}} \times a^{\frac{1}{6}} \times a^{\frac{1}{6}}$ . (Art. 250.)

And  $a^{\frac{2}{6}} = a^{\frac{1}{6}} \times a^{\frac{1}{6}}$ .

The product therefore is  $a^{\frac{1}{6}} \times a^{\frac{1}{6}} \times a^{\frac{1}{6}} \times a^{\frac{1}{6}} \times a^{\frac{1}{6}} = a^{\frac{5}{6}}$ .

And in all instances of this nature, the common denominator of the indices denotes a certain root; and the sum of the numerators shows how often this is to be repeated as a factor to produce the required product.

Thus  $a^{\frac{1}{n}} \times a^{\frac{1}{m}} = a^{\frac{m}{mn}} \times a^{\frac{n}{mn}} = a^{\frac{m+n}{mn}}$ .

Mult.	$3y^{\frac{1}{4}}$	$a^{\frac{1}{2}} \times a^{\frac{1}{3}}$	$(a+b)^{\frac{1}{2}}$	$(a-y)^{\frac{1}{n}}$	$x^{-\frac{1}{4}}$
Into	$y^{\frac{2}{3}}$	$a^{\frac{1}{4}}$	$(a+b)^{\frac{1}{4}}$	$(a-y)^{\frac{1}{m}}$	$x^{-\frac{1}{3}}$
Prod.	$3y^{\frac{11}{12}}$		$(a+b)^{\frac{3}{4}}$		$x^{-\frac{7}{12}}$

The product of  $y^{\frac{1}{2}}$  into  $y^{-\frac{1}{3}}$  is  $y^{\frac{3}{6} - \frac{2}{6}} = y^{\frac{1}{6}}$ .

The product of  $a^{\frac{1}{n}}$  into  $a^{-\frac{1}{n}}$ , is  $a^{\frac{1}{n} - \frac{1}{n}} = a^0 = 1$ .

And  $x^{n - \frac{1}{2}} \times x^{\frac{1}{2} - n} = x^{n - n + \frac{1}{2} - \frac{1}{2}} = x^0 = 1$ .

The product of  $a^2$  into  $a^{\frac{1}{3}} = a^{\frac{6}{3}} \times a^{\frac{1}{3}} = a^{\frac{7}{3}}$ .

281. From the last example, it will be seen, that *powers* and *roots* may be multiplied by a common rule. This is one of the many advantages derived from the notation by fractional indices. Any quantities whatever may be reduced to the form of radicals, (Art. 268,) and may then be subjected to the same modes of operation,

Thus  $y^3 \times y^{\frac{1}{6}} = y^{3 + \frac{1}{6}} = y^{\frac{19}{6}}$ .

And  $x \times x^{\frac{1}{n}} = x^{1 + \frac{1}{n}} = x^{\frac{n+1}{n}}$ .

The product will become rational, whenever the numerator of the index can be exactly divided by the denominator.

$$\text{Thus } a^3 \times a^{\frac{1}{3}} \times a^{\frac{2}{3}} = a^{\frac{1^2}{3}} = a^4.$$

$$\text{And } (a+b)^{\frac{4}{3}} \times (a+b)^{-\frac{1}{3}} = (a+b)^{\frac{3}{3}} = a+b.$$

$$\text{And } a^{\frac{3}{5}} \times a^{\frac{2}{5}} = a^{\frac{5}{5}} = a.$$

282. When radical quantities which are reduced to the same index, have RATIONAL CO-EFFICIENTS, THE RATIONAL PARTS MAY BE MULTIPLIED TOGETHER, AND THEIR PRODUCT PREFIXED TO THE PRODUCT OF THE RADICAL PARTS.

1. Multiply  $a\sqrt{b}$  into  $c\sqrt{d}$ .

The product of the rational parts is  $ac$ .

The product of the radical parts is  $\sqrt{bd}$ .

And the whole product is  $ac\sqrt{bd}$ .

For  $a\sqrt{b}$  is  $a \times \sqrt{b}$ . (Art. 244.) And  $c\sqrt{d}$  is  $c \times \sqrt{d}$ .

By art. 102,  $a \times \sqrt{b}$  into  $c \times \sqrt{d}$ , is  $a \times \sqrt{b} \times c \times \sqrt{d}$ ; or by changing the order of the factors,

$$a \times c \times \sqrt{b} \times \sqrt{d} = ac \times \sqrt{bd} = ac\sqrt{bd}.$$

2. Multiply  $ax^{\frac{1}{2}}$  into  $bd^{\frac{1}{3}}$ .

When the radical parts are reduced to a common index, the factors become  $a(x^3)^{\frac{1}{6}}$  and  $b(d^2)^{\frac{1}{6}}$ .

The product then is  $ab(x^3d^2)^{\frac{1}{6}}$ .

But in cases of this nature, we may save the trouble of reducing to a common index, by multiplying as in art. 278.

Thus  $ax^{\frac{1}{2}}$  into  $bd^{\frac{1}{3}}$  is  $ax^{\frac{1}{2}}bd^{\frac{1}{3}}$ .

Mult.	$a(b+x)^{\frac{1}{2}}$	$a\sqrt{y^2}$	$a\sqrt{x}$	$ax^{-\frac{1}{2}}$	$x^{\frac{3}{3}}$	$\sqrt[3]{3}$
Into	$y(b-x)^{\frac{1}{2}}$	$b\sqrt{hy}$	$b\sqrt{x}$	$by^{\frac{1}{2}}$	$y^{\frac{3}{3}}$	$\sqrt[3]{9}$
Prod.	$ay(b^2-x^2)^{\frac{1}{2}}$	$ab\sqrt{x^2}$	$= abx$			$3xy$

283. If the rational quantities, instead of being *co-efficients* to the radical quantities, are connected with them by the signs  $+$  and  $-$ , each term in the multiplier must be multiplied into each in the multiplicand, as in art. 100.



$$\begin{array}{r}
 \text{Multiply } a + \sqrt{b} \\
 \text{Into } c + \sqrt{d} \\
 \hline
 ac + c\sqrt{b} \\
 a\sqrt{d} + \sqrt{bd} \\
 \hline
 ac + c\sqrt{b} + a\sqrt{d} + \sqrt{bd}.
 \end{array}$$

The product of  $a + \sqrt{y}$  into  $1 + r\sqrt{y}$  is

$$a + \sqrt{y} + ar\sqrt{y} + ry.$$

1. Multiply  $\sqrt{a}$  into  $\sqrt[3]{b}$ . Ans.  $\sqrt[6]{a^3b^2}$ .
2. Multiply  $5\sqrt{5}$  into  $3\sqrt{8}$ . Ans.  $30\sqrt{10}$ .
3. Multiply  $2\sqrt{3}$  into  $3\sqrt[3]{4}$ . Ans.  $6\sqrt[6]{432}$ .
4. Multiply  $\sqrt{d}$  into  $\sqrt[3]{ab}$ . Ans.  $\sqrt[6]{a^2b^2d^3}$ .
5. Multiply  $\sqrt{\frac{2ab}{3c}}$  into  $\sqrt{\frac{9ad}{2b}}$ . Ans.  $\sqrt{\frac{3a^2d}{c}}$ .
6. Multiply  $a(a-x)^{\frac{1}{2}}$  into  $(c-d) \times (ax)^{\frac{1}{2}}$ .  
Ans.  $(ac-ad) \times (a^2x-ax^2)^{\frac{1}{2}}$ .

### DIVISION OF RADICAL QUANTITIES.

284. The division of radical quantities may be expressed, by writing the divisor under the dividend, in the form of a fraction.

Thus the quotient of  $\sqrt[3]{a}$  divided by  $\sqrt{b}$ , is  $\frac{\sqrt[3]{a}}{\sqrt{b}}$ .

And  $(a+h)^{\frac{1}{8}}$  divided by  $(b+x)^{\frac{1}{n}}$  is  $\frac{(a+h)^{\frac{1}{8}}}{(b+x)^{\frac{1}{n}}}$ .

In these instances, the radical sign or index is *separately* applied to the numerator and the denominator. But if the divisor and dividend are reduced to the *same* index or radical sign, this may be applied to the *whole* quotient.

Thus  $\sqrt[n]{a} \div \sqrt[n]{b} = \frac{\sqrt[n]{a}}{\sqrt[n]{b}} = \sqrt[n]{\frac{a}{b}}$ . For the root of a fraction is equal to the root of the numerator divided by the root of the denominator. (Art. 260.)

Again,  $\sqrt[n]{ab} \div \sqrt[n]{b} = \sqrt[n]{a}$ . For the product of this quotient into the divisor is equal to the dividend, that is,

$$\sqrt[n]{a} \times \sqrt[n]{b} = \sqrt[n]{ab}. \text{ Hence,}$$

285. QUANTITIES UNDER THE SAME RADICAL SIGN OR INDEX, MAY BE DIVIDED LIKE RATIONAL QUANTITIES, THE QUOTIENT BEING PLACED UNDER THE COMMON RADICAL SIGN OR INDEX.

Divide  $(x^3y^2)^{\frac{1}{6}}$  by  $y^{\frac{1}{3}}$ .

These reduced to the same index are  $(x^3y^2)^{\frac{1}{6}}$  and  $(y^2)^{\frac{1}{6}}$ .

And the quotient is  $(x^3)^{\frac{1}{6}} = x^{\frac{3}{6}} = x^{\frac{1}{2}}$ .

Divide	$\sqrt{6a^3x}$	$\sqrt{dhx^2}$	$(a^3+ax)^{\frac{1}{9}}$	$(a^3h)^{\frac{1}{m}}$	$(a^2y^2)^{\frac{1}{4}}$
By	$\sqrt{3x}$	$\sqrt{dx}$	$a^{\frac{1}{9}}$	$(ax)^{\frac{1}{m}}$	$(ay)^{\frac{1}{4}}$
Quot.	$\sqrt{2a^3}$		$(a^2+x)^{\frac{1}{9}}$		$(ay)^{\frac{1}{4}}$

286. A ROOT IS DIVIDED BY ANOTHER ROOT OF THE SAME LETTER OR QUANTITY, BY SUBTRACTING THE INDEX OF THE DIVISOR FROM THAT OF THE DIVIDEND.

$$\text{Thus } a^{\frac{1}{2}} \div a^{\frac{1}{6}} = a^{\frac{1}{2} - \frac{1}{6}} = a^{\frac{3}{6} - \frac{1}{6}} = a^{\frac{2}{6}} = a^{\frac{1}{3}}.$$

For  $a^{\frac{1}{2}} = a^{\frac{3}{6}} = a^{\frac{1}{6}} \times a^{\frac{1}{6}} \times a^{\frac{1}{6}}$  and this divided by  $a^{\frac{1}{6}}$  is

$$\frac{a^{\frac{1}{6}} \times a^{\frac{1}{6}} \times a^{\frac{1}{6}}}{a^{\frac{1}{6}}} = a^{\frac{1}{6}} \times a^{\frac{1}{6}} = a^{\frac{2}{6}} = a^{\frac{1}{3}}.$$

In the same manner, it may be shown that  $a^{\frac{1}{m}} \div a^{\frac{1}{n}} = a^{\frac{1}{m} - \frac{1}{n}}$ .

Divide	$(3a)^{\frac{11}{2}}$	$(ax)^{\frac{2}{3}}$	$a^{\frac{m+n}{nm}}$	$(b+y)^{\frac{2}{n}}$	$(r^2y^3)^{\frac{1}{7}}$
By	$a^{\frac{2}{3}}$	$(ax)^{\frac{1}{3}}$	$a^{\frac{1}{m}}$	$(b+y)^{\frac{1}{n}}$	$(r^2y^3)^{\frac{3}{7}}$
Quot.	$(3a)^{\frac{1}{4}}$		$a^{\frac{1}{n}}$		$(r^2y^3)^{-\frac{2}{7}}$

*Powers and roots may be brought promiscuously together, and divided according to the same rule. See art. 281.*

Thus  $a^2 \div a^{\frac{1}{3}} = a^{2 - \frac{1}{3}} = a^{\frac{5}{3}}$ . For  $a^{\frac{5}{3}} \times a^{\frac{1}{3}} = a^{\frac{6}{3}} = a^2$ .

So  $y^n \div y^{\frac{1}{m}} = y^{n - \frac{1}{m}}$ .

287. When radical quantities which are reduced to the same index have RATIONAL CO-EFFICIENTS, THE RATIONAL PARTS MAY BE DIVIDED SEPARATELY, AND THEIR QUOTIENT PREFIXED TO THE QUOTIENT OF THE RADICAL PARTS.

Thus  $ac\sqrt{bd} \div a\sqrt{b} = c\sqrt{d}$ . For this quotient multiplied into the divisor is equal to the dividend.

Divide	$24x\sqrt{ay}$	$18dh\sqrt{bx}$	$by(a^3x^2)^{\frac{1}{n}}$	$16\sqrt{32}$	$b\sqrt{xy}$
By	$6\sqrt{a}$	$2h\sqrt{x}$	$y(ax)^{\frac{1}{n}}$	$8\sqrt{4}$	$\sqrt{y}$
	$4x\sqrt{y}$		$b(a^2x)^{\frac{1}{n}}$		$b\sqrt{x}$

Divide  $ab(x^2b)^{\frac{1}{4}}$  by  $a(x)^{\frac{1}{2}}$ .

These reduced to the same index are  $ab(x^2b)^{\frac{1}{4}}$  and  $a(x^2)^{\frac{1}{4}}$ .

The quotient then is  $b(b)^{\frac{1}{4}} = (b^5)^{\frac{1}{4}}$ . (Art. 272.)

To save the trouble of reducing to a common index, the division may be expressed in the form of a fraction.

The quotient will then be  $\frac{ab(x^2b)^{\frac{1}{4}}}{a(x^{\frac{1}{2}})}$ .

1. Divide  $2^3\sqrt{bc}$  by  $3\sqrt{ac}$ .

Ans.  $\frac{2}{3} \sqrt[6]{\frac{b^2}{a^3c}}$ .

2. Divide  $10^3\sqrt{108}$  by  $5^3\sqrt{4}$ .

Ans.  $2^3\sqrt{27} = 6$ .

3. Divide  $10\sqrt{27}$  by  $2\sqrt{3}$ .

Ans. 15.

4. Divide  $8\sqrt{108}$  by  $2\sqrt{6}$ .

Ans.  $12\sqrt{2}$ .

5. Divide  $(a^2b^2d^3)^{\frac{1}{5}}$  by  $d^{\frac{1}{2}}$ .

Ans.  $(ab)^{\frac{1}{5}}$ .

6. Divide  $(16a^3 - 12a^2x)^{\frac{1}{2}}$  by  $2a$ .

Ans.  $(4a - 3x)^{\frac{1}{2}}$ .

### INVOLUTION OF RADICAL QUANTITIES.

288. RADICAL QUANTITIES, LIKE POWERS, ARE INVOLVED BY MULTIPLYING THE INDEX OF THE ROOT INTO THE INDEX OF THE REQUIRED POWER.

1. The square of  $a^{\frac{1}{3}} = a^{\frac{1}{3}} \times 2 = a^{\frac{2}{3}}$ . For  $a^{\frac{1}{3}} \times a^{\frac{1}{3}} = a^{\frac{2}{3}}$ .

2. The cube of  $a^{\frac{1}{4}} = a^{\frac{1}{4}} \times 3 = a^{\frac{3}{4}}$ . For  $a^{\frac{1}{4}} \times a^{\frac{1}{4}} \times a^{\frac{1}{4}} = a^{\frac{3}{4}}$ .

3. And universally, the  $n$ th power of  $a^{\frac{1}{m}} = a^{\frac{1}{m}} \times n = a^{\frac{n}{m}}$ .

For the  $n$ th power of  $a^{\frac{1}{m}} = a^{\frac{1}{m}} \times a^{\frac{1}{m}} \dots n$  times, and the sum of the indices will then be  $\frac{n}{m}$ .

4. The 5th power of  $a^{\frac{1}{2}} y^{\frac{1}{3}}$ , is  $a^{\frac{5}{2}} y^{\frac{5}{3}}$ . Or, by reducing the roots to a common index,

$$(a^3 y^2)^{\frac{1}{6}} \times 5 = (a^3 y^2)^{\frac{5}{6}}.$$

5. The cube of  $a^{\frac{1}{n}} x^{\frac{1}{m}}$  is  $a^{\frac{3}{n}} x^{\frac{3}{m}}$  or  $(a^m x^n)^{\frac{3}{nm}}$ .

6. The square of  $a^{\frac{2}{3}} x^{\frac{3}{4}}$ , is  $a^{\frac{4}{3}} x^{\frac{6}{4}}$ .

The cube of  $a^{\frac{1}{3}}$  is  $a^{\frac{1}{3}} \times 3 = a^{\frac{3}{3}} = a$ .

And the  $n$ th power of  $a^{\frac{1}{n}}$ , is  $a^{\frac{n}{n}} = a$ . That is,

289. A ROOT IS RAISED TO A POWER OF THE SAME NAME, BY REMOVING THE INDEX OR RADICAL SIGN.

Thus the cube of  $\sqrt[3]{b+x}$ , is  $b+x$ .

And the  $n$ th power of  $(a-y)^{\frac{1}{n}}$ , is  $a-y$ .

290. When the radical quantities have *rational co-efficients*, these must be also involved.

1. The square  $a^2 \sqrt{x}$ , is  $a^{2n} \sqrt{x^2}$ .

$$\text{For } a^2 \sqrt{x} \times a^n \sqrt{x} = a^{2n} \sqrt{x^2}.$$

2. The  $n$ th power of  $a^m x^{\frac{1}{m}}$ , is  $a^{nm} x^{\frac{n}{m}}$ .

3. The square of  $a \sqrt{x-y}$ , is  $a^2 \times (x-y)$ .

5. The cube of  $3a^3 \sqrt{y}$ , is  $27a^3 y$ .

291. But if the radical quantities are connected with others by the signs  $+$  and  $-$ , they must be involved by a multiplication of the several terms, as in art. 213.

Ex. 1. Required the squares of  $a + \sqrt{y}$  and  $a - \sqrt{y}$ .

$$\begin{array}{r} a + \sqrt{y} \\ a + \sqrt{y} \\ \hline a^2 + a\sqrt{y} \\ \quad a\sqrt{y} + y \\ \hline a^2 + 2a\sqrt{y} + y \end{array} \qquad \begin{array}{r} a - \sqrt{y} \\ a - \sqrt{y} \\ \hline a^2 - a\sqrt{y} \\ \quad - a\sqrt{y} + y \\ \hline a^2 - 2a\sqrt{y} + y \end{array}$$

2. Required the cube of  $a - \sqrt{b}$ .
3. Required the cube of  $2d + \sqrt{x}$ .



292. It is unnecessary to give a separate rule for the *evolution* of radical quantities, that is, for finding the root of a quantity which is already a root. The operation is the same as in other cases of evolution. The fractional index of the radical quantity is to be divided, by the number expressing the root to be found. Or, the radical sign belonging to the required root, may be placed over the given quantity. (Art. 257.) If there are rational co-efficients, the roots of these must also be extracted.

Thus, the square root of  $a^{\frac{1}{3}}$ , is  $a^{\frac{1}{3} \div 2} = a^{\frac{1}{6}}$ .

The cube root of  $a(xy)^{\frac{1}{2}}$ , is  $a^{\frac{1}{3}}(xy)^{\frac{1}{6}}$ .

The  $n$ th root of  $a^6\sqrt{by}$ , is  ${}^n\sqrt{a^6\sqrt{by}}$ .

293. It may be proper to observe, that dividing the *fractional* index of a root is the same in effect, as *multiplying* the number which is placed over the radical sign. For this number corresponds with the *denominator* of the fractional index; and a fraction is divided, by *multiplying* its denominator.

$$\begin{array}{ll} \text{Thus } \sqrt[2]{a} = a^{\frac{1}{2}}. & \sqrt[6]{a} = a^{\frac{1}{6}} \\ \sqrt[4]{a} = a^{\frac{1}{4}}. & \sqrt[2n]{a} = a^{\frac{1}{2n}}. \end{array}$$

On the other hand, *multiplying* the fractional index is equivalent to *dividing* the number which is placed over the radical sign.

Thus the square of  $\sqrt[6]{a}$  or  $a^{\frac{1}{6}}$ , is  $\sqrt[3]{a}$  or  $a^{\frac{1}{6} \times 3} = a^{\frac{1}{2}}$ .

293.b. In algebraic calculations, we have sometimes occasion to seek for a factor, which multiplied into a given radical quantity, will render the product *rational*. In the case of a *simple* radical, such a factor is easily found. For if the  $n$ th root of any quantity, be multiplied by the same root raised to a power whose index is  $n-1$ , the product will be the given quantity.

$$\text{Thus } \sqrt[n]{x} \times \sqrt[n]{x^{n-1}} \text{ or } x^{\frac{1}{n}} \times x^{\frac{n-1}{n}} = x^{\frac{n}{n}} = x.$$

$$\text{And } (x+y)^{\frac{1}{n}} \times (x+y)^{\frac{n-1}{n}} = x+y.$$

$$\text{So } \sqrt{a} \times \sqrt{a} = a. \quad \text{And } \sqrt[3]{a} \times \sqrt[3]{a^2} = \sqrt[3]{a^3} = a.$$

$$\text{And } \sqrt[4]{a} \times \sqrt[4]{a^3} = a, \text{ \&c.} \quad \text{And } (a+b)^{\frac{1}{3}} \times (a+b)^{\frac{2}{3}} = a+b.$$

$$\text{And } (x+y)^{\frac{1}{4}} \times (x+y)^{\frac{3}{4}} = x+y.$$

293.c. A factor which will produce a rational product, when multiplied into a *binomial surd* containing only the *square root*, may be found, by applying the principle, that the product of the sum and difference of two quantities, is equal to the difference of their squares. (Art. 235.) The binomial itself, after the sign which connects the terms is changed from  $+$  to  $-$ , or from  $-$  to  $+$ , will be the factor required.

Thus  $(\sqrt{a} + \sqrt{b}) \times (\sqrt{a} - \sqrt{b}) = \sqrt{a^2} - \sqrt{b^2} = a - b$ , which is free from radicals.

$$\text{So } (1 + \sqrt{2}) \times (1 - \sqrt{2}) = 1 - 2 = -1.$$

$$\text{And } (3 - 2\sqrt{2}) \times (3 + 2\sqrt{2}) = 1.$$

When the compound surd consists of *more than two* terms, it may be reduced, by successive multiplications, first to a binomial surd, and then to a rational quantity.

Thus  $(\sqrt{10} - \sqrt{2} - \sqrt{3}) \times (\sqrt{10} + \sqrt{2} + \sqrt{3}) = 5 - 2\sqrt{6}$ , a binomial surd.

$$\text{And } (5 - 2\sqrt{6}) \times (5 + 2\sqrt{6}) = 1.$$

Therefore  $(\sqrt{10} - \sqrt{2} - \sqrt{3})$  multiplied into  $(\sqrt{10} + \sqrt{2} + \sqrt{3}) \times (5 + 2\sqrt{6}) = 1$ .

293.d. It is sometimes desirable to clear from radical signs the numerator or denominator of a *fraction*. This may be effected, without altering the value of the fraction, if the

numerator and denominator be both multiplied by a factor which will render either of them rational, as the case may require.

1. If both parts of the fraction  $\frac{\sqrt{a}}{\sqrt{x}}$  be multiplied by  $\sqrt{a}$ , it will become  $\frac{\sqrt{a} \times \sqrt{a}}{\sqrt{x} \times \sqrt{a}} = \frac{a}{\sqrt{ax}}$ , in which the *numerator* is a rational quantity.

Or if both parts of the given fraction be multiplied by  $\sqrt{x}$ , it will become  $\frac{\sqrt{ax}}{x}$ , in which the *denominator* is rational.

2. The fraction 
$$\frac{b^{\frac{1}{2}}}{(a+x)^{\frac{1}{3}}} = \frac{b^{\frac{1}{2}} \times (a+x)^{\frac{2}{3}}}{(a+x)^{\frac{1}{3} + \frac{2}{3}}} = \frac{b^{\frac{1}{2}} \times (a+x)^{\frac{2}{3}}}{a+x}.$$

3. The fraction 
$$\frac{\sqrt[3]{y+x}}{a} = \frac{(y+x)^{\frac{1}{3} + \frac{2}{3}}}{a(y+x)^{\frac{2}{3}}} = \frac{y+x}{a(y+x)^{\frac{2}{3}}}.$$

4. The fraction 
$$\frac{a}{x^n} = \frac{ax^{\frac{n-1}{n}}}{x^n \times x^{\frac{n-1}{n}}} = \frac{a\sqrt[n]{x^{n-1}}}{x}.$$

5. The fraction 
$$\frac{\sqrt{2}}{3-\sqrt{2}} = \frac{\sqrt{2} \times (3+\sqrt{2})}{(3-\sqrt{2})(3+\sqrt{2})} = \frac{2+3\sqrt{2}}{7}.$$

6. The fraction 
$$\frac{3}{\sqrt{5}-\sqrt{2}} = \frac{3(\sqrt{5}+\sqrt{2})}{(\sqrt{5}-\sqrt{2})(\sqrt{5}+\sqrt{2})} = \sqrt{5}+\sqrt{2}.$$

7. The fraction 
$$\frac{6}{5^{\frac{1}{4}}} = \frac{6 \times 5^{\frac{3}{4}}}{5^{\frac{1}{4} + \frac{3}{4}}} = \frac{6}{5} \sqrt[4]{125}.$$

8. The fraction 
$$\frac{8}{\sqrt{3}+\sqrt{2}+1} = \frac{8 \times (\sqrt{3}-\sqrt{2}-1)(-\sqrt{2})}{(\sqrt{3}+\sqrt{2}+1)(\sqrt{3}-\sqrt{2}-1)(-\sqrt{2})} = 4-2\sqrt{6}+2\sqrt{2}.$$

9. Reduce  $\frac{2}{\sqrt{3}}$  to a fraction having a rational denominator.

10. Reduce  $\frac{a-\sqrt{b}}{a+\sqrt{b}}$  to a fraction having a rational denominator.

293.e. The arithmetical operation of finding the proximate value of a fractional surd, may be shortened, by ren-

dering either the numerator or the denominator rational. The root of a fraction is equal to the root of the numerator divided by the root of the denominator. (Art. 260.)

Thus  $\frac{\sqrt[n]{a}}{b} = \frac{\sqrt[n]{a}}{\sqrt[n]{b}}$ . But this may be reduced to  $\frac{a}{\sqrt[n]{b} \times \sqrt[n]{a^{n-1}}}$   
or  $\frac{\sqrt[n]{a} \times \sqrt[n]{b^{n-1}}}{b}$ . (Art. 293.d.)

The square root of  $\frac{a}{b}$  is  $\frac{\sqrt{a}}{\sqrt{b}}$ , or  $\frac{a}{\sqrt{ab}}$ , or  $\frac{\sqrt{ab}}{b}$ .

When the fraction is thrown into this form, the process of extracting the root arithmetically, will be confined either to the numerator, or to the denominator.

Thus the square root of  $\frac{3}{7} = \frac{\sqrt{3}}{\sqrt{7}} = \frac{\sqrt{3} \times \sqrt{7}}{\sqrt{7} \times \sqrt{7}} = \frac{\sqrt{21}}{7}$ .

#### Examples for Practice.

1. Find the 4th root of  $81a^2$ .
2. Find the 6th root of  $(a+b)^{-3}$ .
3. Find the  $n$ th root of  $(x-y)^{\frac{1}{6}}$ .
4. Find the cube root of  $-125 a^3 x^6$ .
5. Find the square root of  $\frac{4a^4}{9x^2y^2}$ .
6. Find the 5th root of  $\frac{32a^5x^{10}}{243}$ .
7. Find the square root of  $x^2 - 6bx + b^2$ .
8. Find the square root of  $a^2 + ay + \frac{y^2}{4}$ .
9. Reduce  $ax^2$  to the form of the 6th root.
10. Reduce  $-3y$  to the form of the cube root.
11. Reduce  $a^2$  and  $a^{\frac{1}{3}}$  to a common index.
12. Reduce  $4^{\frac{1}{3}}$  and  $5^{\frac{1}{4}}$  to a common index.
13. Reduce  $a^{\frac{1}{2}}$  and  $b^{\frac{1}{3}}$  to the common index  $\frac{1}{6}$ .
14. Reduce  $2^{\frac{1}{2}}$  and  $4^{\frac{1}{3}}$  to the common index  $\frac{1}{6}$ .



15. Remove a factor from  $\sqrt{294}$ .
16. Remove a factor from  $\sqrt{x^3 - a^2x^2}$ .
17. Find the sum and difference of  $\sqrt{16a^2x}$  and  $\sqrt{4a^2x}$ .
18. Find the sum and difference of  $\sqrt[3]{192}$  and  $\sqrt[3]{24}$ .
19. Multiply  $7\sqrt[3]{18}$  into  $5\sqrt[3]{4}$ .
20. Multiply  $4 + 2\sqrt{2}$  into  $2 - \sqrt{2}$ .
21. Multiply  $a(a + \sqrt{c})^{\frac{1}{2}}$  into  $b(a - \sqrt{c})^{\frac{1}{2}}$ .
22. Multiply  $2(a + b)^{\frac{1}{n}}$  into  $3(a + b)^{\frac{1}{m}}$ .
23. Divide  $6\sqrt{54}$  by  $3\sqrt{2}$ .
24. Divide  $4\sqrt[3]{72}$  by  $2\sqrt[3]{18}$ .
25. Divide  $\sqrt{7}$  by  $\sqrt[3]{7}$ .
26. Divide  $8\sqrt[3]{512}$  by  $4\sqrt[3]{2}$ .
27. Find the cube of  $17\sqrt{21}$ .
28. Find the square of  $5 + \sqrt{2}$ .
29. Find the 4th power of  $\frac{1}{6}\sqrt{6}$ .
30. Find the cube of  $\sqrt{x} - \sqrt{b}$ .
31. Find a factor which will make  $\sqrt[3]{y}$  rational.
32. Find a factor which will make  $\sqrt{5} - \sqrt{x}$  rational.
33. Reduce  $\frac{\sqrt{a}}{\sqrt{x}}$  to a fraction having a rational numerator.
34. Reduce  $\frac{\sqrt{6}}{\sqrt{7} \times \sqrt{3}}$  to a fraction having a rational denominator.

*A gentleman lived simple  
 and a lady lived simple  
 which was most to be  
 Let S = gentle man, he might have  
 5000 lbs, and the lady  
 5000 lbs, and the gentleman  
 5000 lbs, and the lady*

## SECTION X.

REDUCTION OF EQUATIONS BY INVOLUTION  
AND EVOLUTION.

ART. 294. **I**N an equation, the letter which expresses the unknown quantity is sometimes found under a *radical sign*. We may have  $\sqrt{x} = a$ .

To clear this of the radical sign, let each member of the equation be squared, that is, multiplied into itself. We shall then have

$$\sqrt{x} \times \sqrt{x} = aa, \quad \text{Or, (Art. 289,) } x = a^2.$$

The equality of the sides is not affected by this operation, because each is only multiplied into itself, that is, equal quantities are multiplied into equal quantities-

The same principle is applicable to any root whatever. If  $\sqrt[n]{x} = a$ ; then  $x = a^n$ . For by art. 289, a root is raised to a power of the same name, by removing the index or radical sign. Hence,

295. **WHEN THE UNKNOWN QUANTITY IS UNDER A RADICAL SIGN, THE EQUATION IS REDUCED BY INVOLVING BOTH SIDES, to a power of the same name, as the root expressed by the radical sign.**

It will generally be expedient to make the necessary transpositions, *before* involving the quantities; so that all those which are not under the radical sign, may stand on one side of the equation.

Ex. 1. Reduce the equation

Transposing +4

Involving both sides

$$\sqrt{x} + 4 = 9$$

$$\sqrt{x} = 9 - 4 = 5$$

$$x = 5^2 = 25.$$

2. Reduce the equation

By transposition,

By involution,

$$a + \sqrt[n]{x} - b = d$$

$$\sqrt[n]{x} = d + b - a$$

$$x = (d + b - a)^n.$$

*If a lower radical be taken and a cross, it is considered as a root.*

3. Reduce the equation  $\sqrt[3]{x+1}=4$   
 Involving both sides,  $x+1=4^3=64$   
 And  $x=63$ .

4. Reduce the equation  $4+3\sqrt{x-4}=6+\frac{1}{2}$   
 Clearing of fractions,  $8+6\sqrt{x-4}=13$   
 And  $\sqrt{x-4}=\frac{5}{6}$   
 Involving both sides,  $x-4=\frac{25}{36}$   
 And  $x=\frac{25}{36}+4$ .

5. Reduce the equation  $\sqrt{a^2+\sqrt{x}}=\frac{3+d}{\sqrt{a^2+\sqrt{x}}}$   
 Multiplying by  $\sqrt{a^2+\sqrt{x}}$ ,  $a^2+\sqrt{x}=3+d$   
 And  $\sqrt{x}=3+d-a^2$   
 Involving both sides,  $x=(3+d-a^2)^2$ .

In the first step in this example, multiplying the first member into  $\sqrt{a^2+\sqrt{x}}$ , that is, into itself, is the same as squaring it, which is done by taking away its radical sign. The other member being a fraction, is multiplied into a quantity equal to its denominator, by cancelling the denominator. (Art. 159.) There remains a radical sign over  $x$ , which must be removed by involving both sides of the equation.

6. Reduce  $3+2\sqrt{x-\frac{4}{5}}=6$ .      Ans.  $x=\frac{36}{10}\frac{1}{10}$ .

7. Reduce  $4\sqrt{\frac{x}{5}}=8$ .      Ans.  $x=20$ .

8. Reduce  $(2x+3)^{\frac{1}{3}}+4=7$ .      Ans.  $x=12$ .

9. Reduce  $\sqrt{12+x}=2+\sqrt{x}$ .      Ans.  $x=4$ .

10. Reduce  $\sqrt{x-a}=\sqrt{x}-\frac{1}{2}\sqrt{a}$ .      Ans.  $x=\frac{25a}{16}$ .

11. Reduce  $\sqrt{5}\times\sqrt{x+2}=2+\sqrt{5x}$ .      Ans.  $x=\frac{9}{20}$ .

12. Reduce  $\frac{x-ax}{\sqrt{x}}=\frac{\sqrt{x}}{x}$ .      Ans.  $x=\frac{1}{1-a}$ .

13. Reduce  $\frac{\sqrt{x+28}}{\sqrt{x+4}}=\frac{\sqrt{x+38}}{\sqrt{x+6}}$ .      Ans.  $x=4$ .

14. Reduce  $\sqrt{x} + \sqrt{a+x} = \frac{2a}{\sqrt{a+x}}$ .      Ans.  $x = \frac{1}{3}a$ .

15. Reduce  $x + \sqrt{a^2 + x^2} = \frac{2a^2}{\sqrt{a^2 + x^2}}$ .      Ans.  $x = a\sqrt{\frac{1}{3}}$ .

16. Reduce  $x + a = \sqrt{a^2 + x\sqrt{b^2 + x^2}}$ .      Ans.  $x = \frac{b^2 - 4a^2}{4a}$ .

17. Reduce  $\sqrt{2+x} + \sqrt{x} = \frac{4}{\sqrt{2+x}}$ .      Ans.  $x = \frac{2}{3}$ .

18. Reduce  $\sqrt{x-32} = 16 - \sqrt{x}$ .      Ans.  $x = 81$ .

19. Reduce  $\sqrt{4x+17} = 2\sqrt{x+1}$ .      Ans.  $x = 16$ .

20. Reduce  $\frac{\sqrt{6x-2}}{\sqrt{6x+2}} = \frac{4\sqrt{6x-9}}{4\sqrt{6x+6}}$ .      Ans.  $x = 6$ .

### REDUCTION OF EQUATIONS BY EVOLUTION.

296. In many equations, the letter which expresses the unknown quantity is involved to some power. Thus in the equation

$$x^2 = 16$$

we have the value of the *square* of  $x$ , but not of  $x$  itself. If the square root of both sides be extracted, we shall have

$$x = 4.$$

The equality of the members is not affected by this reduction. For if two quantities or sets of quantities are equal, their roots are also equal.

If  $(x+a)^n = b+h$ , then  $x+a = \sqrt[n]{b+h}$ . Hence,

297. WHEN THE EXPRESSION CONTAINING THE UNKNOWN QUANTITY IS A POWER, THE EQUATION IS REDUCED BY EXTRACTING THE ROOT OF BOTH SIDES, A ROOT OF THE SAME NAME AS THE POWER.

Ex. 1. Reduce the equation

$$6 + x^2 - 8 = 7$$

By transposition

$$x^2 = 7 + 8 - 6 = 9$$

By evolution

$$x = \pm\sqrt{9} = \pm 3.$$

The signs  $+$  and  $-$  are both placed before  $\sqrt{9}$ , because an even root of an affirmative quantity is *ambiguous*. (Art. 261.)

2. Reduce the equation  $5x^2 - 30 = x^2 + 34$   
 Transposing, &c.  $x^2 = 16$   
 By evolution,  $x = \pm 4$ .

3. Reduce the equation  $a + \frac{x^2}{b} = h - \frac{x^2}{d}$   
 Clearing of fractions, &c.  $x^2 = \frac{bdh - abd}{b + d}$   
 By evolution,  $x = \pm \left( \frac{bdh - abd}{b + d} \right)^{\frac{1}{2}}$

4. Reduce the equation  $a + dx^n = 10 - x^n$   
 Transposing, &c.  $x^n = \frac{10 - a}{d + 1}$   
 By evolution,  $x = \left( \frac{10 - a}{d + 1} \right)^{\frac{1}{n}}$ .

298. From the preceding articles, it will be easy to see in what manner an equation is to be reduced, when the expression containing the unknown quantity is a power, and at the same time under a radical sign; that is, when it is a root of a power. Both involution and evolution will be necessary in this case.

Ex. 1. Reduce the equation  $\sqrt[3]{x^2} = 4$   
 By involution  $x^2 = 4^3 = 64$   
 By evolution  $x = \pm \sqrt[3]{64} = \pm 4$ .

2. Reduce the equation  $\sqrt{x^m - a} = h - d$   
 By involution  $x^m - a = h^2 - 2hd + d^2$   
 And  $x^m = h^2 - 2hd + d^2 + a$   
 By evolution  $x = \sqrt[m]{h^2 - 2hd + d^2 + a}$ .

3. Reduce the equation  $(x + a)^{\frac{1}{2}} = \frac{a + b}{(x - a)^{\frac{1}{2}}}$   
 Multiplying by  $(x - a)^{\frac{1}{2}}$  (Art. 279.)  $(x^2 - a^2)^{\frac{1}{2}} = a + b$   
 By involution  $x^2 - a^2 = a^2 + 2ab + b^2$   
 Trans. and uniting terms  $x^2 = 2a^2 + 2ab + b^2$   
 By evolution  $x = (2a^2 + 2ab + b^2)^{\frac{1}{2}}$ .

*Problems.*

Prob. 1. A gentleman being asked his age, replied ; " If you add to it ten years, and extract the square root of the sum, and from this root subtract 2, the remainder will be 6." What was his age?

By the conditions of the problem	$\sqrt{x+10}-2=6$
By transposition,	$\sqrt{x+10}=6+2=8$
By involution,	$x+10=8^2=64.$
And	$x=64-10=54.$

Proof (Art. 194.)  $\sqrt{54+10}-2=6.$

Prob. 2. If to a certain number 22577 be added, and the square root of the sum be extracted, and from this 163 be subtracted, the remainder will be 237. What is the number?

Let $x$ =the number sought.	$b=163$
$a=22577$	$c=237.$

By the conditions proposed	$\sqrt{x+a}-b=c$
By transposition,	$\sqrt{x+a}=c+b$
By involution,	$x+a=(c+b)^2$
And	$x=(c+b)^2-a$
Restoring the numbers, (Art. 52.)	$x=(237+163)^2-22577$
That is,	$x=160000-22577=137423.$
Proof	$\sqrt{137423+22577}-163=237.$

299. When an equation is reduced by extracting an even root of a quantity, the solution does not determine whether the answer is positive or negative. (Art. 297.) But what is thus left ambiguous by the algebraic process, is frequently settled by the statement of the problem.

Prob. 3. A merchant gains in trade a sum, to which 320 dollars bears the same proportion, as five times this sum does to 2500. What is the amount gained?

Let $x$ =the sum required.
$a=320$
$b=2500.$

By the supposition  $a : x :: 5x : b$

Multiplying extremes and means  $5x^2 = ab$

And  $x = \left(\frac{ab}{5}\right)^{\frac{1}{2}}$

Restoring the numbers,  $x = \left(\frac{320 \times 2500}{5}\right)^{\frac{1}{2}} = 400.$

Here the answer is not marked as ambiguous, because by the statement of the problem it is *gain*, and not loss. It must therefore be positive. This might be determined, in the present instance, even from the algebraic process. Whenever the root of  $x^2$  is ambiguous, it is because we are ignorant whether the power has been produced by the multiplication of  $+x$ , or of  $-x$ , into itself. (Art. 262.) But here we have the multiplication actually performed. By turning back to the two first steps of the equation, we find that  $5x^2$  was produced by multiplying  $5x$  into  $x$ , that is  $+5x$  into  $+x$ .

Prob. 4. The distance to a certain place is such, that if 96 be subtracted from the square of the number of miles, the remainder will be 48. What is the distance?

Let  $x$  = the distance required.

By the supposition,  $x^2 - 96 = 48$

Therefore  $x = \sqrt{144} = 12.$

Prob. 5. If three times the square of a certain number be divided by four, and if the quotient be diminished by 12, the remainder will be 180. What is the number?

By the supposition  $\frac{3x^2}{4} - 12 = 180.$

Therefore  $x = \sqrt{256} = 16.$

Prob. 6. What number is that, the fourth part of whose square being subtracted from 8, leaves a remainder equal to four?  
Ans. 4.

Prob. 7. What two numbers are those, whose sum is to the greater as 10 to 7; and whose sum multiplied into the less produces 270?

Let  $10x$  = their sum.

Then  $7x$  = the greater, and  $3x$  = the less.

Therefore  $x = 3$ , and the numbers required are 21 and 9.

Prob. 8. What two numbers are those, whose difference is to the greater as 2 : 9 ; and the difference of whose squares is 128 ?  
 Ans. 18 and 14.

Prob. 9. It is required to divide the number 18 into two such parts, that the squares of those parts may be to each other as 25 to 16.

Let $x$ = the greater part.	Then $18 - x$ = the less.
By the condition proposed	$x^2 : (18 - x)^2 :: 25 : 16.$
Therefore	$16x^2 = 25 \times (18 - x)^2.$
By evolution	$4x = 5 \times (18 - x).$
And	$x = 10.$

Prob. 10. It is required to divide the number 14 into two such parts, that the quotient of the greater divided by the less, may be to the quotient of the less divided by the greater, as 16 : 9.  
 Ans. The parts are 8 and 6.

Prob. 11. What two numbers are as 5 to 4, the sum of whose cubes is 5103 ?

Let  $5x$  and  $4x$  = the two numbers.

Then  $x = 3$ , and the numbers are 15 and 12.

Prob. 12. Two travellers  $A$  and  $B$  set out to meet each other,  $A$  leaving the town  $C$ , at the same time that  $B$  left  $D$ . They travelled the direct road between  $C$  and  $D$ ; and on meeting, it appeared that  $A$  had travelled 18 miles more than  $B$ , and that  $A$  could have gone  $B$ 's distance in  $15\frac{3}{4}$  days, but  $B$  would have been 28 days in going  $A$ 's distance. Required the distance between  $C$  and  $D$ .

Let  $x$  = the number of miles  $A$  travelled.

Then  $x - 18$  = the number  $B$  travelled.

$$\frac{x - 18}{15\frac{3}{4}} = A\text{'s daily progress.}$$

$$\frac{x}{28} = B\text{'s daily progress.}$$

$$\text{Therefore } x : x - 18 :: \frac{x - 18}{15\frac{3}{4}} : \frac{x}{28}.$$

This reduced gives  $x = 72$ ,  $A$ 's distance.

The whole distance, therefore, from  $C$  to  $D = 126$  miles.



Prob. 13. Find two numbers, which are to each other as 8 to 5, and whose product is 360.      Ans. 24 and 15.

Prob. 14. A gentleman bought two pieces of silk, which together measured 36 yards. Each of them cost as many shillings by the yard, as there were yards in the piece, and their whole prices were as 4 to 1. What were the lengths of the pieces?      Ans. 24 and 12 yards.

Prob. 15. Find two numbers, which are to each other as 3 to 2; and the difference of whose fourth powers, is to the sum of their cubes, as 26 to 7.

Ans. The numbers are 6 and 4.

Prob. 16. Several gentlemen made an excursion, each taking the same sum of money. Each had as many servants attending him as there were gentlemen, the number of dollars which each had was double the number of all the servants, and the whole sum of money taken out was 3456 dollars. How many gentlemen were there?      Ans. 12.

Prob. 17. A detachment of soldiers from a regiment being ordered to march on a particular service, each company furnished four times as many men, as there were companies in the whole regiment; but these being found insufficient, each company furnished 3 men more; when their number was found to be increased in the ratio of 17 to 16. How many companies were there in the regiment?      Ans. 12.

### AFFECTED QUADRATIC EQUATIONS.

300. Equations are divided into classes, which are distinguished from each other, by the power of the letter that expresses the unknown quantity. Those which contain only the *first* power of the unknown quantity, are called equations of *one dimension*, or equations of the *first degree*. Those in which the highest power of the unknown quantity is a *square*, are called *quadratic*, or equations of the *second degree*; those in which the highest power is a *cube*, equations of the *third degree*, &c.

Thus  $x = a + b$ , is an equation of the *first degree*.

$x^2 = c$ , and  $x^2 + ax = d$ , are *quadratic* equations, or equations of the *second degree*.

$x^3 = h$ , and  $x^3 + ax^2 + bx = d$ , are *cubic* equations, or equations of the *third degree*.

301. Equations are also divided into *pure* and *affected* equations. A pure equation contains only *one power* of the unknown quantity. This may be the first, second, third, or any other power. An affected equation contains *different powers* of the unknown quantity, Thus,

$$\begin{cases} x^2 = d - b, & \text{is a pure quadratic equation.} \\ x^2 + bx = d, & \text{an affected quadratic equation.} \\ x^3 = b - c, & \text{a pure cubic equation.} \\ x^3 + ax^2 + bx = h, & \text{an affected cubic equation.} \end{cases}$$

A pure equation is also called a *simple* equation. But this term has been applied in too vague a manner. By some writers, it is extended to pure equations of every degree : by others, it is confined to those of the first degree.

In a *pure* equation, all the terms which contain the unknown quantity may be united in one, (Art. 185,) and the equation, however complicated in other respects, may be reduced by the rules which have already been given. But in an *affected* equation, as the unknown quantity is raised to *different powers*, the terms containing these powers can not be united. (Art. 230.) There are particular rules for the reduction of quadratic, cubic, and biquadratic equations. Of these, only the first will be considered at present.

302. AN AFFECTED QUADRATIC EQUATION IS ONE WHICH CONTAINS THE UNKNOWN QUANTITY IN ONE TERM, AND THE SQUARE OF THAT QUANTITY IN ANOTHER TERM.

The unknown quantity may be originally in *several* terms of the equation. But all these may be reduced to two, one containing the unknown quantity, and the other its square.

303. It has already been shown that a *pure* quadratic is solved by *extracting the root of both sides of the equation*. An *affected* quadratic may be solved in the same way, if the member which contains the unknown quantity is an *exact square*. Thus the equation

$$x^2 + 2ax + a^2 = b + h$$

may be reduced by evolution. For the first member is the square of a *binomial* quantity. (Art. 264.) And its root is  $x + a$ . Therefore,

$$\begin{aligned} x + a &= \sqrt{b + h}, \text{ and by transposing } a, \\ x &= \sqrt{b + h} - a. \end{aligned}$$

304. But it is not often the case, that a member of an affected quadratic equation is an exact square, till an additional term is applied, for the purpose of making the required reduction. In the equation

$$x^2 + 2ax = b$$

the side containing the unknown quantity is not a complete square. The two terms of which it is composed are indeed such, as might belong to the square of a binomial quantity. (Art. 214.) But one term is *wanting*. We have then to inquire, in what way this may be supplied. From having *two* terms of the square of a binomial given, how shall we find the *third*?

Of the three terms, two are complete powers, and the other is twice the product of the roots of these powers; (Art. 214,) or, which is the same thing, the product of one of the roots into twice the other. In the expression

$$x^2 + 2ax,$$

the term  $2ax$  consists of the factors  $2a$  and  $x$ . The latter is the unknown quantity. The other factor  $2a$  may be considered the *co-efficient* of the unknown quantity; a co-efficient being another name for a factor. (Art. 41.) As  $x$  is the root of the first term  $x^2$ ; the other factor  $2a$  is *twice* the root of the third term, which is wanted to complete the square. Therefore *half*  $2a$  is the root of the deficient term, and  $a^2$  is the term itself. The square completed is

$$x^2 + 2ax + a^2,$$

where it will be seen that the last term  $a^2$  is the square of half  $2a$ , and  $2a$  is the co-efficient of  $x$  the root of the first term.

In the same manner, it may be proved, that the last term of the square of any binomial quantity, is equal to the square of half the co-efficient of the root of the first term. From this principle, is derived the following rule :

305. TO COMPLETE THE SQUARE, in an affected quadratic equation; TAKE THE SQUARE OF HALF THE CO-EFFICIENT OF THE FIRST POWER OF THE UNKNOWN QUANTITY, AND ADD IT TO BOTH SIDES OF THE EQUATION.

Before completing the square, the known and unknown quantities must be brought on opposite sides of the equation by transposition; and the highest power of the unknown

quantity must have the affirmative sign, and be cleared of fractions, co-efficients, &c. See arts. 308, 9, 10, 11.

After the square is completed, the equation is reduced, by extracting the square root of both sides, and transposing the known part of the binomial root. (Art. 303.)

The quantity which is added to one side of the equation, to complete the square, must be added to the other side also, to preserve the equality of the two members. (Ax. 1.)

306. It will be important for the learner to distinguish between what is *peculiar* in the reduction of quadratic equations, and what is common to this and the other kinds which have already been considered. The peculiar part, in the resolution of affected quadratics, is the *completing of the square*. The other steps are similar to those by which pure equations are reduced.

For the purpose of rendering the completing of the square familiar, there will be an advantage in beginning with examples in which the equation is already prepared for this step.

Ex. 1. Reduce the equation	$x^2 + 6ax = b$
Completing the square	$x^2 + 6ax + 9a^2 = 9a^2 + b$
Extracting both sides (Art. 303.)	$x + 3a = \pm\sqrt{9a^2 + b}$
And	$x = -3a \pm\sqrt{9a^2 + b}$

Here the co-efficient of  $x$ , in the first step, is  $6a$ ;

The square of half this is  $9a^2$ , which being added to both sides completes the square. The equation is then reduced by extracting the root of each member, in the same manner as in art. 297, excepting that the square here being that of a *binomial*, its root is found by the rule in art. 265.

2. Reduce the equation	$x^2 - 8bx = h$
Completing the square,	$x^2 - 8bx + 16b^2 = 16b^2 + h$
Extracting both sides	$x - 4b = \pm\sqrt{16b^2 + h}$
And	$x = 4b \pm\sqrt{16b^2 + h}$

In this example, half the co-efficient of  $x$  is  $4b$ , the square of which  $16b^2$  is to be added to both sides of the equation.

3. Reduce the equation  $x^2 + ax = b + h$

Completing the square  $x^2 + ax + \frac{a^2}{4} = \frac{a^2}{4} + b + h$

By evolution  $x + \frac{a}{2} = \pm \left( \frac{a^2}{4} + b + h \right)^{\frac{1}{2}}$

And  $x = -\frac{a}{2} \pm \left( \frac{a^2}{4} + b + h \right)^{\frac{1}{2}}$ .

4. Reduce the equation  $x^2 - x = h - d$

Completing the square,  $x^2 - x + \frac{1}{4} = \frac{1}{4} + h - d$

And  $x = \frac{1}{2} \pm \left( \frac{1}{4} + h - d \right)^{\frac{1}{2}}$ .

Here the co-efficient of  $x$  is 1, the square of half which is  $\frac{1}{4}$ .

5. Reduce the equation  $x^2 + 3x = d + 6$

Completing the square,  $x^2 + 3x + \frac{9}{4} = \frac{9}{4} + d + 6$

And  $x = -\frac{3}{2} \pm \left( \frac{9}{4} + d + 6 \right)^{\frac{1}{2}}$ .

6. Reduce the equation  $x^2 - abx = ab - cd$

Completing the square,  $x^2 - abx + \frac{a^2b^2}{4} = \frac{a^2b^2}{4} + ab - cd$

And  $x = \frac{ab}{2} \pm \left( \frac{a^2b^2}{4} + ab - cd \right)^{\frac{1}{2}}$

7. Reduce the equation  $x^2 + \frac{ax}{b} = h$

Completing the square,  $x^2 + \frac{ax}{b} + \frac{a^2}{4b^2} = \frac{a^2}{4b^2} + h$

And  $x = -\frac{a}{2b} \pm \left( \frac{a^2}{4b^2} + h \right)^{\frac{1}{2}}$ .

By art. 158,  $\frac{ax}{b} = \frac{a}{b} \times x$ . The co-efficient of  $x$ , therefore,

is  $\frac{a}{b}$ . Half of this is  $\frac{a}{2b}$ , (Art. 163.) the square of which is  $\frac{a^2}{4b^2}$ .

8. Reduce the equation  $x^2 - \frac{x}{b} = 7h$ .

Completing the square,  $x^2 - \frac{x}{b} + \frac{1}{4b^2} = \frac{1}{4b^2} + 7h$

And  $x = \frac{1}{2b} + \left(\frac{1}{4b^2} + 7h\right)^{\frac{1}{2}}$ .

Here the fraction  $\frac{x}{b} = \frac{1}{b} \times x$ . (Art. 158.) Therefore the co-efficient of  $x$  is  $\frac{1}{b}$ .

307. In these and similar instances, the root of the third term of the completed square is easily found, because, this root is the same half co-efficient from which the term has just been derived. (Art. 304.) Thus in the last example, half the co-efficient of  $x$  is  $\frac{1}{2b}$ , and this is the root of the third term  $\frac{1}{4b^2}$ .

308. When the first power of the unknown quantity is in several terms, these should be united in one, if they can be by the rules for reduction in addition. But if there are *literal* co-efficients, these may be considered as constituting, together, a *compound* co-efficient or factor, into which the unknown quantity is multiplied.

Thus  $ax + bx + dx = (a + b + d) \times x$ . (Art. 120.) The square of half this compound co-efficient is to be added to both sides of the equation.

1. Reduce the equation  $x^2 + 3x + 2x + x = d$   
 Uniting terms,  $x^2 + 6x = d$   
 Completing the square,  $x^2 + 6x + 9 = 9 + d$   
 And  $x = -3 \pm \sqrt{9 + d}$

2. Reduce the equation  $x^2 + ax + bx = h$   
 By art. 120,  $x^2 + (a + b) \times x = h$

Therefore  $x^2 + (a + b) \times x + \left(\frac{a + b}{2}\right)^2 = \left(\frac{a + b}{2}\right)^2 + h$

By evolution,  $x + \frac{a + b}{2} = \pm \sqrt{\left(\frac{a + b}{2}\right)^2 + h}$

And  $x = -\frac{a + b}{2} \pm \sqrt{\left(\frac{a + b}{2}\right)^2 + h}$

3. Reduce the equation  $x^2 + ax - x = b$

By art. 120  $x^2 + (a-1)x = b$

Therefore  $x^2 + (a-1)x + \left(\frac{a-1}{2}\right)^2 = \left(\frac{a-1}{2}\right)^2 + b$

And  $x = -\frac{a-1}{2} \pm \sqrt{\left(\frac{a-1}{2}\right)^2 + b}$

309. After becoming familiar with the method of completing the square, in affected quadratic equations, it will be proper to attend to the steps which are preparatory to this. Here, however, little more is necessary, than an application of rules already given. The known and unknown quantities must be brought on opposite sides of the equation by transposition. And it will generally be expedient to make the square of the unknown quantity the first or leading term, as in the preceding examples. This indeed is not essential. But it will show, to the best advantage, the arrangement of the terms in the completed square.

1. Reduce the equation  $a + 5x - 3b = 3x - x^2$   
 Transp. and uniting terms  $x^2 + 2x = 3b - a$   
 Completing the square  $x^2 + 2x + 1 = 1 + 3b - a$   
 And  $x = -1 \pm \sqrt{1 + 3b - a}$

2. Reduce the equation  $\frac{x}{2} = \frac{36}{x+2} - 4$   
 Clearing of fractions, &c.  $x^2 + 10x = 56$   
 Completing the square  $x^2 + 10x + 25 = 25 + 56 = 81$   
 And  $x = -5 \pm \sqrt{81} = -5 \pm 9$

310. If the highest power of the unknown quantity has any co-efficient or divisor, it must, before the square is completed, by the rule in art. 305, be freed from these, by multiplication or division, as in arts. 180 and 184.

1. Reduce the equation  $x^2 + 24a - 6h = 12x - 5x^2$   
 Transp. and uniting terms;  $6x^2 - 12x = 6h - 24a$   
 Dividing by 6,  $x^2 - 2x = h - 4a$   
 Completing the square,  $x^2 - 2x + 1 = 1 + h - 4a$   
 Extracting and transp.  $x = 1 \pm \sqrt{1 + h - 4a}$

2. Reduce the equation  $h + 2x = d - \frac{bx^2}{a}$

Clearing of fractions  $bx^2 + 2ax = ad - ah$

Dividing by  $b$ ,  $x^2 + \frac{2ax}{b} = \frac{ad - ah}{b}$

Therefore  $x^2 + \frac{2ax}{b} + \frac{a^2}{b^2} = \frac{a^2}{b^2} + \frac{ad - ah}{b}$

And  $x = -\frac{a}{b} + \left(\frac{a^2}{b^2} + \frac{ad - ah}{b}\right)^{\frac{1}{2}}$

311. If the square of the unknown quantity is in *several terms*, the equation must be divided by *all* the co-efficients of this square, as in art. 185.

1. Reduce the equation  $bx^2 + dx^2 - 4x = b - h$

Dividing by  $b + d$ , (Art. 121.)  $x^2 - \frac{4x}{b+d} = \frac{b-h}{b+d}$

Therefore  $x = \frac{2}{b+d} + \sqrt{\left(\frac{2}{b+d}\right)^2 + \frac{b-h}{b+d}}$

2. Reduce the equation  $ax^2 + x = h + 3x - x^2$

Transp. and uniting terms  $ax^2 + x^2 - 2x = h$

Dividing by  $a + 1$ ,  $x^2 - \frac{2x}{a+1} = \frac{h}{a+1}$

Completing the square  $x^2 - \frac{2x}{a+1} + \left(\frac{1}{a+1}\right)^2 = \left(\frac{1}{a+1}\right)^2 + \frac{h}{a+1}$

Extracting and transp.  $x = \frac{1}{a+1} + \sqrt{\left(\frac{1}{a+1}\right)^2 + \frac{h}{a+1}}$

There is another method of completing the square, which, in many cases, particularly those in which the highest power of the unknown quantity has a co-efficient, is more simple in its application, than that given in art. 305.

Let  $ax^2 + bx = d$ .

If the equation be multiplied by  $4a$ , and if  $b^2$  be added to both sides, it will become

$$4a^2x^2 + 4abx + b^2 = 4ad + b^2;$$

the first member of which is a complete power of  $2ax + b$ .

Hence,

311.b. In a quadratic equation, the square may be completed, by multiplying the equation into 4 times the co-effi-



cient of the highest power of the unknown quantity, and adding to both sides, the square of the co-efficient of the lowest power.

The advantage of this method is, that it avoids the introduction of *fractions*, in completing the square.

This will be seen, by solving an equation by both methods.

$$\text{Let } ax^2 + dx = h.$$

Completing the square, by the rule just given ;

$$4a^2x^2 + 4adx + d^2 = 4ah + d^2$$

Extracting the root  $2ax + d = \pm \sqrt{4ah + d^2}$

And 
$$x = \frac{-d \pm \sqrt{4ah + d^2}}{2a}$$

Completing the square of the given equation by arts. 305 and 310 ;

$$x^2 + \frac{dx}{a} + \frac{d^2}{4a^2} = \frac{h}{a} + \frac{d^2}{4a^2}.$$

Extracting the root  $x + \frac{d}{2a} = \pm \sqrt{\frac{h}{a} + \frac{d^2}{4a^2}}$

And 
$$x = -\frac{d}{2a} \pm \sqrt{\frac{h}{a} + \frac{d^2}{4a^2}}.$$

If  $a=1$ , the rule will be reduced to this: "Multiply the equation by 4, and add to both sides, the square of the co-efficient of  $x$ ."

$$\text{Let } x^2 + dx = h$$

Completing the square  $4x^2 + 4dx + d^2 = 4h + d^2$

Extracting the root  $2x + d = \pm \sqrt{4h + d^2}$

And 
$$x = \frac{-d \pm \sqrt{4h + d^2}}{2}.$$

1. Reduce the equation  $3x^2 + 5x = 42$

Completing the square  $36x^2 + 60x + 25 = 529$

Therefore  $x = 3.$

2. Reduce the equation  $x^2 - 15x = -54$

Completing the square  $4x^2 - 60x + 225 = 9$

Therefore  $2x = 15 \pm 3 = 18 \text{ or } 12.$

312. In the square of a binomial, the first and last terms are always *positive*. For each is the square of one of the

terms of the root. (Art. 214.) But every square is positive. (Art. 218.) If then  $-x^2$  occurs in an equation, it can not, with this sign, form a part of the square of a binomial. But if *all* the signs in the equation be changed, the equality of the sides will be preserved, (Art. 177,) the term  $-x^2$  will become positive, and the square may be completed.

1. Reduce the equation	$-x^2 + 2x = d - h$
Changing all the signs	$x^2 - 2x = h - d$
Therefore	$x = 1 \pm \sqrt{1 + h - d}$ .
2. Reduce the equation	$4x - x^2 = -12$
Answer	$x = 2 \pm \sqrt{16}$ .

313. In a quadratic equation, the first term  $x^2$  is the square of a single letter. But a binomial quantity may consist of terms, one or both of which are already powers.

Thus  $x^3 + a$  is a binomial, and its square is

$$x^6 + 2ax^3 + a^2,$$

where the index of  $x$  in the first term is twice as great as in the second. When the third term is deficient, the square may be completed in the same manner as that of any other binomial. For the middle term is twice the product of the roots of the two others.

So the square of  $x^n + a$ , is  $x^{2n} + 2ax^n + a^2$ .

And the square of  $x^{\frac{1}{n}} + a$ , is  $x^{\frac{2}{n}} + 2ax^{\frac{1}{n}} + a^2$ .

Therefore,

314. ANY EQUATION WHICH CONTAINS ONLY TWO DIFFERENT POWERS OR ROOTS OF THE UNKNOWN QUANTITY, THE INDEX OF ONE OF WHICH IS TWICE THAT OF THE OTHER, MAY BE RESOLVED IN THE SAME MANNER AS A QUADRATIC EQUATION, BY COMPLETING THE SQUARE.

It must be observed, however, that in the binomial root, the letter expressing the unknown quantity may still have a fractional or integral index, so that a farther extraction, according to art. 297, may be necessary.

1. Reduce the equation	$x^4 - x^2 = b - a$
Completing the square	$x^4 - x^2 + \frac{1}{4} = \frac{1}{4} + b - a$
Extracting and transp.	$x^2 = \frac{1}{2} \pm \sqrt{\frac{1}{4} + b - a}$
Extracting again, (Art. 297,)	$x = \pm \sqrt{\frac{1}{2} \pm \sqrt{\frac{1}{4} + b - a}}$ .

2. Reduce the equation  $x^{2n} - 4bx^n = a$   
 Answer  $x = \pm \sqrt[n]{2b \pm \sqrt{4b^2 + a}}$
3. Reduce the equation  $x + 4\sqrt{x} = h - n$   
 Completing the square  $x + 4\sqrt{x} + 4 = h - n + 4$   
 Extracting and transp.  $\sqrt{x} = -2 \pm \sqrt{h - n + 4}$   
 Involving  $x = (-2 \pm \sqrt{h - n + 4})^2$
4. Reduce the equation  $\frac{x}{x^n} + 8x^{\frac{1}{n}} = a + b$   
 Completing the square  $\frac{x}{x^n} + 8x^{\frac{1}{n}} + 16 = a + b + 16$   
 Extracting and transp.  $\frac{1}{x^n} = -4 \pm \sqrt{a + b + 16}$   
 Involving  $x = (-4 \pm \sqrt{a + b + 16})^n$

315. The solution of a quadratic equation, whether pure or affected, gives two results. For after the equation is reduced, it contains an ambiguous root. In a *pure* quadratic, this root is the *whole* value of the unknown quantity. (Art. 297.)

Thus the equation  $x^2 = 64$   
 Becomes, when reduced,  $x = \pm \sqrt{64}$

That is, the value of  $x$  is either  $+8$  or  $-8$ , for each of these is a root of  $64$ . Here both the values of  $x$  are the same, except that they have contrary signs. This will be the case in every pure quadratic equation, because the whole of the second member is under the radical sign. The two values of the unknown quantity will be alike, except that one will be positive, and the other negative.

316. But in *affected* quadratics, a *part* only of one side of the reduced equation is under the radical sign. When this part is added to, or subtracted from, that which is without the radical sign; the two results will differ in quantity, and will have their signs in some cases alike, and in others unlike.

1. The equation  $x^2 + 8x = 20$   
 Becomes, when reduced  $x = -4 \pm \sqrt{16 + 20}$   
 That is  $x = -4 \pm 6$

Here the first value of  $x$  is,  $-4 + 6 = +2$  } one positive, and  
 And the second is,  $-4 - 6 = -10$  } the other negative.

$$\begin{array}{ll}
 2. \text{ The equation} & x^2 - 8x = -15 \\
 \text{Becomes, when reduced} & x = 4 \pm \sqrt{16 - 15} \\
 \text{That is} & x = 4 \pm 1
 \end{array}$$

Here the first value of  $x$  is  $4 + 1 = +5$  } both positive.  
 And the second is  $4 - 1 = +3$  }

That these two values of  $x$  are correctly found, may be proved, by substituting first one, and then the other, for  $x$  itself, in the original equation. (Art. 194.)

$$\text{Thus } 5^2 - 8 \times 5 = 25 - 40 = -15$$

$$\text{And } 3^2 - 8 \times 3 = 9 - 24 = -15.$$

317. In the reduction of an affected quadratic equation, the value of the unknown quantity is frequently found to be *imaginary*.

$$\begin{array}{ll}
 \text{Thus the equation} & x^2 - 3x = -20 \\
 \text{Becomes, when reduced,} & x = 4 \pm \sqrt{16 - 20} \\
 \text{That is,} & x = 4 \pm \sqrt{-4}.
 \end{array}$$

Here the root of the negative quantity  $-4$  can not be assigned, (Art. 263,) and therefore the value of  $x$  can not be found. There will be the same impossibility, in every instance in which the negative part of the quantities under the radical sign is greater than the positive part.\*

318. Whenever *one* of the values of the unknown quantity, in a quadratic equation, is imaginary, the *other* is so also. For both are equally affected by the imaginary root.

Thus, in the example above,

$$\begin{array}{ll}
 \text{The first value of } x \text{ is} & 4 + \sqrt{-4}, \\
 \text{And the second is} & 4 - \sqrt{-4}; \text{ each of which} \\
 \text{contains the imaginary quantity} & \sqrt{-4}.
 \end{array}$$

319. An equation which when reduced contains an imaginary root, is often of use, to enable us to determine whether a proposed question admits of an answer, or involves an absurdity.

Suppose it is required to divide 8 into two such parts, that the product will be 20.

\* See Note C.

If  $x$  is one of the parts, the other will be  $8-x$ . (Art. 195.)

By the conditions proposed  $(8-x) \times x = 20$

This becomes, when reduced,  $x = 4 \pm \sqrt{-4}$ .

Here the imaginary expression  $\sqrt{-4}$  shows that an answer is impossible; and that there is an absurdity in supposing that 8 may be divided into two such parts, that their product shall be 20.

320. Although a quadratic equation has two solutions, yet both these may not always be applicable to the subject proposed. The quantity under the radical sign may be produced either from a positive or a negative root. But both these roots may not, in every instance, belong to the problem to be solved. See art. 299.

Divide the number 30 into two such parts, that their product may be equal to 8 times their difference.

If  $x =$  the lesser part, then  $30-x =$  the greater.

By the supposition,  $x \times (30-x) = 8 \times (30-2x)$

This reduced, gives  $x = 23 \pm 17 = 40$  or  $6 =$  the lesser part.

But as 40 can not be a part of 30, the problem can have but one real solution, making the lesser part 6, and the greater part 24.

*Examples of Quadratic Equations.*

1. Reduce  $3x^2 - 9x - 4 = 80$ .      Ans.  $x = 7$ , or  $-4$ .

2. Reduce  $4x - \frac{36-x}{x} = 46$ .      Ans.  $x = 12$ , or  $-\frac{3}{4}$ .

3. Reduce  $4x - \frac{14-x}{x+1} = 14$ .      Ans.  $x = 4$ , or  $-\frac{7}{4}$ .

4. Reduce  $5x - \frac{3x-3}{x-3} = 2x + \frac{3x-6}{2}$ .      Ans.  $x = 4$ , or  $-1$ .

5. Reduce  $\frac{16}{x} - \frac{100-9x}{4x^2} = 3$ .      Ans.  $x = 4$ , or  $2\frac{1}{2}$ .

6. Reduce  $\frac{3x-4}{x-4} + 1 = 10 - \frac{x-2}{2}$ .      Ans.  $x = 12$ , or  $6$ .

7. Reduce  $\frac{x+4}{3} - \frac{7-x}{x-3} = \frac{4x+7}{9} - 1$ .      Ans.  $x = 21$ , or  $5$ .

8. Reduce  $\frac{x^3 - 10x^2 + 1}{x^2 - 6x + 9} = x - 3$ .      Ans.  $x = 1$ , or  $-23$ .
9. Reduce  $\frac{6}{x+1} + \frac{2}{x} = 3$ .      Ans.  $x = 2$ .
10. Reduce  $\frac{3x}{x+2} - \frac{x-1}{6} = x - 9$ .      Ans.  $x = 10$ .
11. Reduce  $\frac{x}{a} + \frac{a}{x} = \frac{2}{a}$ .      Ans.  $x = 1 \pm \sqrt{1 - a^2}$ .
12. Reduce  $x^4 + ax^2 = b$ .      Ans.  $x = \left( -\frac{a}{2} \pm \sqrt{b + \frac{a^2}{4}} \right)^{\frac{1}{2}}$ .
13. Reduce  $\frac{x^6}{2} - \frac{x^3}{4} = -\frac{1}{32}$ .      Ans.  $x = \sqrt{\frac{1}{4}}$ .
14. Reduce  $2x^{\frac{2}{3}} + 3x^{\frac{1}{3}} = 2$ .      Ans.  $x = \frac{1}{8}$ .
15. Reduce  $\frac{1}{2}x - \frac{1}{3}\sqrt{x} = 22\frac{1}{3}$ .      Ans.  $x = 49$ .
16. Reduce  $2x^4 - x^2 + 96 = 99$ .      Ans.  $x = \frac{1}{2}\sqrt{6}$ .
17. Reduce  $(10+x)^{\frac{1}{2}} - (10+x)^{\frac{1}{4}} = 2$ .      Ans.  $x = 6$ .
18. Reduce  $3x^{2n} - 2x^n = 8$ .      Ans.  $x = \sqrt[n]{2}$ .
19. Reduce  $2(1+x-x^2) - \sqrt{1+x-x^2} = -\frac{1}{3}$ .  
Ans.  $x = \frac{1}{2} + \frac{1}{6}\sqrt{33}$  41
20. Reduce  $\sqrt[3]{x^3 - a^3} = x - b$ .      Ans.  $x = \frac{b}{2} + \sqrt{\frac{4a^3 - b^3}{12b}}$ .
21. Reduce  $\frac{\sqrt{4x+2}}{4+\sqrt{x}} = \frac{4-\sqrt{x}}{\sqrt{x}}$ .      Ans.  $x = 4$ .
22. Reduce  $x^{\frac{5}{3}} + x^{\frac{3}{3}} = 756$ .      Ans.  $x = 243$ .
23. Reduce  $\sqrt{2x+1} + 2\sqrt{x} = \frac{21}{\sqrt{2x+1}}$ .      Ans.  $x = 4$ .
24. Reduce  $2\sqrt{x-a} + 3\sqrt{2x} = \frac{7a+5x}{\sqrt{x-a}}$ .      Ans.  $x = 9a$ .
25. Reduce  $x+16 - 7\sqrt{x+16} = 10 - 4\sqrt{x+16}$ .      Ans.  $x = 9$ .
26. Reduce  $\sqrt{x^5} + \sqrt{x^3} = 6\sqrt{x}$ .  
Dividing by  $\sqrt{x}$ ,       $x^2 + x = 6$       And  $x = 2$ .

27. Reduce  $\frac{4x-5}{x} - \frac{3x-7}{3x+7} = \frac{9x+23}{13x}$ .      Ans.  $x=2$ .

28. Reduce  $\frac{3}{6x-x^2} + \frac{6}{x^2+2x} = \frac{11}{5x}$ .      Ans.  $x=3$ .

29. Reduce  $(x-5)^3 - 3(x-5)^{\frac{3}{2}} = 40$ .      Ans.  $x=9$ .

30. Reduce  $x + \sqrt{x+6} = 2 + 3\sqrt{x+6}$ .      Ans.  $x=10$ .

**PROBLEMS PRODUCING QUADRATIC EQUATIONS.**

Prob. 1. A merchant has a piece of cotton cloth, and a piece of silk. The number of yards in both is 110; and if the square of the number of yards of silk be subtracted from 80 times the number of yards of cotton, the difference will be 400. How many yards are there in each piece?

Let  $x$  = the yards of silk.

Then  $110 - x$  = the yards of cotton.

By supposition       $400 = 80 \times (110 - x) - x^2$

Therefore       $x = -40 \pm \sqrt{10000} = -40 \pm 100$ .

The first value of  $x$ , is  $-40 + 100 = 60$ , the yards of silk;

And       $110 - x = 110 - 60 = 50$ , the yards of cotton.

The second value of  $x$ , is  $-40 - 100 = -140$ ; but as this is a negative quantity, it is not applicable to goods which a man has in his possession.

Prob. 2. The ages of two brothers are such, that their sum is 45 years, and their product 500. What is the age of each?

Ans. 25 and 20 years.

Prob. 3. To find two numbers such, that their difference shall be 4, and their product 117.

Let  $x$  = one number, and  $x + 4$  = the other.

By the conditions       $(x+4) \times x = 117$

This reduced, gives       $x = -2 \pm \sqrt{121} = -2 \pm 11$ .

One of the numbers therefore is 9, and the other 13.

Prob. 4. A merchant having sold a piece of cloth which cost him 30 dollars, found that if the price for which he sold it were multiplied by his gain, the product would be equal to the cube of his gain. What was his gain?

Let  $x$  = the gain.

Then  $30 + x$  = the price for which the cloth was sold.

By the statement  $x^2 = (30 + x) \times x$

Therefore  $x = \frac{1}{2} \pm \sqrt{\frac{1}{4} + 30} = \frac{1}{2} \pm \frac{11}{2}$

The first value of  $x$  is  $\frac{1}{2} + \frac{11}{2} = +6.$  }

The second value is  $\frac{1}{2} - \frac{11}{2} = -5.$  }

As the last answer is *negative*, it is to be rejected as inconsistent with the nature of the problem, (Art. 320,) for *gain* must be considered *positive*.

Prob. 5. To find two numbers, whose difference shall be 3, and the difference of their cubes 117.

Let  $x$  = the least number.

Then  $x + 3$  = the greatest.

By supposition  $(x + 3)^3 - x^3 = 117$

Expanding  $(x + 3)^3$  (Art. 217.)  $9x^2 + 27x = 117 - 27 = 90$

And  $x = -\frac{3}{2} \pm \sqrt{\frac{49}{4}} = -\frac{3}{2} \pm \frac{7}{2}.$

The two numbers, therefore, are 2 and 5.

Prob. 6. To find two numbers, whose difference shall be 12, and the sum of their squares 1424.

Ans. The numbers are 20 and 32.

Prob. 7. Two persons draw prizes in a lottery, the difference of which is 120 dollars, and the greater is to the less, as the less to 10. What are the prizes?

Ans. 40 and 160.

Prob. 8. What two numbers are those whose sum is 6, and the sum of their cubes 72?

Ans. 2 and 4.

Prob. 9. Divide the number 56 into two such parts, that their product shall be 640.

Putting  $x$  for one of the parts, we have,  $x = 28 \pm 12 = 40$  or 16.

In this case, the two values of the unknown quantity, are the two parts into which the given number was required to be divided.

Prob. 10. A gentleman bought a number of pieces of cloth for 675 dollars, which he sold again at 48 dollars by the piece, and gained by the bargain as much as one piece cost him. What was the number of pieces? Ans. 15.



Prob. 11. *A* and *B* started together, for a place 150 miles distant. *A*'s hourly progress was 3 miles more than *B*'s, and he arrived at his journey's end, 8 hours and 20 minutes before *B*. What was the hourly progress of each?

Ans. 9 and 6 miles.

Prob. 12. The difference of two numbers is 6; and if 47 be added to twice the square of the less, it will be equal to the square of the greater. What are the numbers?

Ans. 17 and 11.

Prob. 13. *A* and *B* distributed 1200 dollars each, among a certain number of persons. *A* relieved 40 persons more than *B*, and *B* gave to each individual 5 dollars more than *A*. How many were relieved by *A* and *B*?

Ans. 120 by *A*, and 80 by *B*.

Prob. 14. Find two numbers, whose sum is 10, and the sum of their squares 58?

Ans. 7 and 3.

Prob. 15. Several gentlemen made a purchase in company for 175 dollars. Two of them having withdrawn, the bill was paid by the others, each furnishing 10 dollars more than would have been his equal share, if the bill had been paid by the whole company. What was the number in the company at first?

Ans. 7.

Prob. 16. A merchant bought several yards of linen for 60 dollars, out of which he reserved 15 yards, and sold the remainder for 54 dollars, gaining 10 cents a yard. How many yards did he buy, and at what price?

Ans. 75 yards, at 80 cents a yard.

Prob. 17. *A* and *B* set out from two towns, which were 247 miles distant, and travelled the direct road till they met. *A* went 9 miles a day; and the number of days which they travelled before meeting, was greater by 3, than the number of miles which *B* went in a day. How many miles did each travel?

Ans. *A* went 117, and *B* 130 miles.

Prob. 18. A gentlemen bought two pieces of cloth, the finer of which cost 4 shillings a yard more than the other. The finer piece cost £18; but the coarser one, which was 2 yards longer than the finer, cost only £16. How many yards were there in each piece, and what was the price of a yard of each?

Ans. There were 18 yards of the finer piece, and 20 of the coarser; and the prices were 20 and 16 shillings.

Prob. 19. A merchant bought 54 gallons of Madeira wine, and a certain quantity of Teneriffe. For the former, he gave half as many shillings by the gallon, as there were gallons of Teneriffe, and for the latter, 4 shillings less by the gallon. He sold the mixture at 10 shillings by the gallon, and lost £28 16s. by his bargain. Required the price of the Madeira, and the number of gallons of Teneriffe.

Ans. The Madeira cost 18 shillings a gallon, and there were 36 gallons of Teneriffe.

Prob. 20. If the square of a certain number be taken from 40, and the square root of this difference be increased by 10, and the sum be multiplied by 2, and the product divided by the number itself, the quotient will be 4. What is the number?  
Ans. 6.

Prob. 21. A person being asked his age, replied, If you add the square root of it to half of it, and subtract 12, the remainder will be nothing. What was his age?

Ans. 16 years.

Prob. 22. Two casks of wine were purchased for 58 dollars, one of which contained 5 gallons more than the other, and the price by the gallon, was 2 dollars less, than  $\frac{1}{3}$  of the number of gallons in the smaller cask. Required the number of gallons in each, and the price by the gallon.

Ans. The numbers were 12 and 17, and the price by the gallon 2 dollars.

Prob. 23. In a parcel which contains 24 coins of silver and copper, each silver coin is worth as many cents as there are copper coins, and each copper coin is worth as many cents as there are silver coins; and the whole are worth 2 dollars and 16 cents. How many are there of each?

Ans. 6 of one, and 18 of the other.

Prob. 24. A person bought a certain number of oxen for 80 guineas. If he had received 4 more oxen for the same money, he would have paid one guinea less for each. What was the number of oxen?  
Ans. 16.

### SUBSTITUTION.

321. In the reduction of Quadratic Equations, as well as in other parts of algebra, a complicated process may be rendered much more simple, by introducing a new letter which

shall be made to represent several others. This is termed *substitution*. A letter may be put for a compound quantity as well as for a single number. Thus in the equation

$$x^2 - 2ax = \frac{3}{4} + \sqrt{86 - 64 + h},$$

we may substitute  $b$ , for  $\frac{3}{4} + \sqrt{86 - 64 + h}$ . The equation will then become  $x^2 - 2ax = b$ , and when reduced will be

$$x = a \pm \sqrt{a^2 + b}.$$

After the operation is completed, the compound quantity for which a single letter has been substituted, may be *restored*. The last equation, by restoring the value of  $b$ , will become

$$x = a \pm \sqrt{a^2 + \frac{3}{4} + \sqrt{86 - 64 + h}}.$$

Reduce the equation  $ax - 2x - d = bx - x^2 - x$

Transposing, &c.  $x^2 + (a - b - 1)x = d$

Substituting  $h$  for  $(a - b - 1)$ ,  $x^2 + hx = d$

Therefore  $x = -\frac{h}{2} \pm \sqrt{\frac{h^2}{4} + d}$

Restoring the value of  $h$ ,  $x = -\frac{a - b - 1}{2} \pm \sqrt{\frac{(a - b - 1)^2}{4} + d}$



SECTION XI.



SOLUTION OF PROBLEMS WHICH CONTAIN TWO OR MORE UNKNOWN QUANTITIES.

DEMONSTRATION OF THEOREMS.

ART. 322. **I**N the examples which have been given of the resolution of equations, in the preceding sections, each problem has contained only *one* unknown quantity. Or if, in some instances, there have been *two*, they have been so

related to each other, that they have both been expressed by means of the same letter. (Art. 195.)

But cases frequently occur in which *several* unknown quantities are introduced into the same calculation. And if the problem is of such a nature, as to admit of a determinate answer, there will arise from the conditions, as many equations independent of each other, as there are unknown quantities.

Equations are said to be *independent*, when they express different conditions; and *dependent*, when they express the same conditions under different forms. The former are not convertible into each other. But the latter may be changed from one form to the other, by the methods of reduction which have been considered. Thus  $b - x = y$ , and  $b = y + x$ , are dependent equations, because one is formed from the other by merely transposing  $x$ .

323. In solving a problem, it is necessary first to find the value of one of the unknown quantities, and then of the others in succession. To do this, we must derive from the equations which are given, a new equation, from which all the unknown quantities except one shall be excluded.

Suppose the following equations are given.

$$1. \quad x + y = 14$$

$$2. \quad x - y = 2.$$

If  $y$  be transposed in each, they will become

$$1. \quad x = 14 - y$$

$$2. \quad x = 2 + y.$$

Here the first member of each of the equations is  $x$ , and the second member of each is *equal* to  $x$ . But according to axiom 11th, quantities which are respectively equal to any other quantity are equal to each other; therefore,

$$2 + y = 14 - y.$$

Here we have a new equation, which contains only the unknown quantity  $y$ . Hence,

324 *Rule I.* To exterminate one of two unknown quantities, and deduce one equation from two; FIND THE VALUE OF ONE OF THE UNKNOWN QUANTITIES IN EACH OF THE EQUATIONS, AND FORM A NEW EQUATION BY MAKING ONE OF THESE VALUES EQUAL TO THE OTHER.

That quantity which is the least involved should be the one which is chosen to be exterminated.

For the convenience of referring to different parts of a solution, the several steps will, in future, be numbered. When an equation is formed from one *immediately preceding*, it will be unnecessary to specify it. In other cases, the number of the equation or equations from which a new one is derived will be referred to.

Prob. 1. To find two numbers such, that  
 Their sum shall be 24; and  
 The greater shall be equal to five times the less.

- |                                     |                            |
|-------------------------------------|----------------------------|
| Let $x$ = the greater;              | And $y$ = the less.        |
| 1. By the first condition,          | $x + y = 24$ }<br>}        |
| 2. By the second,                   | $x = 5y$ }<br>}            |
| 3. Transp. $y$ in the 1st equation, | $x = 24 - y$               |
| 4. Making the 2d and 3d equal,      | $5y = 24 - y$              |
| 5. And                              | $y = 4$ , the less number. |

Prob. 2. To find one of two quantities,  
 Whose sum is equal to  $h$ ; and  
 The difference of whose squares is equal to  $d$ .

- |                                      |                            |
|--------------------------------------|----------------------------|
| Let $x$ = the greater quantity;      | And $y$ = the less.        |
| 1. By the first condition,           | $x + y = h$ }<br>}         |
| 2. By the second,                    | $x^2 - y^2 = d$ }<br>}     |
| 3. Transp. $y^2$ in the 2d equation, | $x^2 = d + y^2$            |
| 4. By evolution, (Art. 297.)         | $x = \sqrt{d + y^2}$       |
| 5. Transp. $y$ in the 1st equation,  | $x = h - y$                |
| 6. Making the 4th and 5th equal      | $\sqrt{d + y^2} = h - y$   |
| 7. Therefore                         | $y = \frac{h^2 - d}{2h}$ . |

Prob. 3. Given  $ax + by = h$  }  
 And  $x + y = d$  } To find  $y$ . Ans.  $y = \frac{h - ad}{b - a}$ .

325. The rule given above may be generally applied, for the extermination of unknown quantities. But there are cases, in which other methods will be found more expeditious.

$$\left. \begin{array}{l} \text{Suppose } x = hy \\ \text{And } ax + bx = y^2 \end{array} \right\}$$

As in the first of these equations  $x$  is equal to  $hy$ , we may, in the second equation, *substitute* this value of  $x$  instead of  $x$  itself. The second equation will then be converted into  

$$ahy + bhy = y^2.$$

The equality of the two sides is not affected by this alteration, because we only change one quantity  $x$ , for another which is equal to it. By this means we obtain an equation which contains only one unknown quantity. Hence,

326. *Rule II.* To exterminate an unknown quantity, FIND THE VALUE OF ONE OF THE UNKNOWN QUANTITIES, IN ONE OF THE EQUATIONS; and then, in the other EQUATION, SUBSTITUTE THIS VALUE, FOR THE UNKNOWN QUANTITY ITSELF.

Prob. 4. A privateer in chase of a ship 20 miles distant, sails 8 miles, while the ship sails 7. How far must the privateer sail, before she overtakes the ship?

It is evident that the whole distance which the privateer sails during the chase, must be to the distance which the ship sails in the same time, as 8 to 7.

Let  $x$  = the distance which the privateer sails;  
 And  $y$  = the distance which the ship sails.

1. By the supposition,  $x = y + 20$  }  
 2. And also,  $x : y :: 8 : 7$  }  
 3. Art. 138,  $y = \frac{7}{8}x$   
 4. Substituting  $\frac{7x}{8}$  for  $y$  in the 1st equation,  $x = \frac{7}{8}x + 20$   
 5. Therefore  $x = 160$ .

Prob. 5. The ages of two persons  $A$  and  $B$  are such, that seven years ago,  $A$  was three times as old as  $B$ ; and seven years hence,  $A$  will be twice as old as  $B$ . What is the age of  $B$ ?

Let  $x$  = the age of  $A$ ;      And  $y$  = the age of  $B$ ;

Then  $x - 7$  was the age of  $A$ , 7 years ago;

And  $y - 7$  was the age of  $B$ , 7 years ago.

Also  $x + 7$  will be the age of  $A$ , 7 years hence;

And  $y + 7$  will be the age of  $B$ , 7 years hence.

1. By the first condition,  $x - 7 = 3 \times (y - 7) = 3y - 21$  }  
 2. By the second,  $x + 7 = 2 \times (y + 7) = 2y + 14$  }  
 3. Transp. 7 in the 1st equa.  $x = 3y - 14$   
 4. Subst.  $3y - 14$  for  $x$ , in the 2d,  $3y - 14 + 7 = 2y + 14$   
 5. Therefore  $y = 21$ , the age of  $B$ .

- Prob. 6. There are two numbers, of which  
 The greater is to the less, as 3 to 2; and  
 Their sum is the sixth part of their product.  
 What is the less number?      Ans. 10.

327. There is a *third* method of exterminating an unknown quantity from an equation, which, in many cases, is preferable to either of the preceding.

$$\begin{array}{l} \text{Suppose that } x+3y=a \} \\ \text{And that } \quad \quad x-3y=b \} \end{array}$$

If we *add together* the first members of these two equations, and also the second members, we shall have

$$2x=a+b$$

an equation which contains only the unknown quantity  $x$ . The other, having equal co-efficients with contrary signs, has disappeared. (Art. 77.) The equality of the sides is preserved, because we have only added equal quantities to equal quantities.

$$\begin{array}{l} \text{Again, suppose } 3x+y=h \} \\ \text{And } \quad \quad \quad 2x+y=d \} \end{array}$$

If we *subtract* the last equation from the first, we shall have

$$x=h-d$$

where  $y$  is exterminated, without affecting the equality of the sides.

$$\begin{array}{l} \text{Again, suppose } x-2y=a \} \\ \text{And } \quad \quad \quad x+4y=b \} \end{array}$$

$$\text{Multiplying the 1st by 2, } \quad 2x-4y=2a$$

$$\text{Then adding the 2d and 3d, } \quad 3x=b+2a. \quad \text{Hence,}$$

328. *Rule III.* To exterminate an unknown quantity,

MULTIPLY OR DIVIDE THE EQUATIONS, IF NECESSARY, IN SUCH A MANNER THAT THE TERM WHICH CONTAINS ONE OF THE UNKNOWN QUANTITIES SHALL BE THE SAME IN BOTH.

THEN SUBTRACT ONE EQUATION FROM THE OTHER, IF THE SIGNS OF THIS UNKNOWN QUANTITY ARE ALIKE, OR ADD THEM TOGETHER, IF THE SIGNS ARE UNLIKE.

It must be kept in mind that both members of an equation are always to be increased or diminished, multiplied or divided alike. (Art. 170.)

Prob. 7. The numbers in two opposing armies are such, that,

The sum of both is 21110 ; and

Twice the number in the greater army, added to three times the number in the less, is 52219.

What is the number in the greater army?

Let  $x$  = the greater.

And  $y$  = the less.

- |                                    |                   |   |
|------------------------------------|-------------------|---|
| 1. By the first condition,         | $x + y = 21110$   | } |
| 2. By the second,                  | $2x + 3y = 52219$ |   |
| 3. Multiplying the 1st by 3,       | $3x + 3y = 63330$ |   |
| 4. Subtracting the 2d from the 3d, | $x = 11111.$      |   |

Prob. 8. Given  $2x + y = 16$ , and  $3x - 3y = 6$ , to find the value of  $x$ .

- |                              |                |   |
|------------------------------|----------------|---|
| 1. By supposition,           | $2x + y = 16$  | } |
| 2. And                       | $3x - 3y = 6$  |   |
| 3. Multiplying the 1st by 3, | $6x + 3y = 48$ |   |
| 4. Adding the 2d and 3d,     | $9x = 54$      |   |
| 5. Dividing by 9,            | $x = 6.$       |   |

Prob. 9. Given  $x + y = 14$ , and  $x - y = 2$ , to find the value of  $y$ . Ans. 6.

In the succeeding problems, either of the three rules for exterminating unknown quantities will be made use of, as will in each case be most convenient.

329. When *one* of the unknown quantities is determined, the other may be easily obtained, by going back to an equation which contains both, and substituting, instead of that which is already found, its numerical value.

Prob. 10. The mast of a ship consists of two parts :

One third of the lower part, added to one sixth of the upper part, is equal to 28 ; and

Five times the lower part, diminished by six times the upper part, is equal to 12.

What is the height of the mast?



Let  $x$  = the lower part;      And  $y$  = the upper part.

- |                                       |                                      |
|---------------------------------------|--------------------------------------|
| 1. By the first condition,            | $\frac{1}{3}x + \frac{1}{6}y = 28$ } |
| 2. By the second,                     | $5x - 6y = 12$ }                     |
| 3. Multiplying the 1st by 6,          | $2x + y = 168$                       |
| 4. Dividing the 2d by 6,              | $\frac{5}{6}x - y = 2$               |
| 5. Adding the 3d and 4th,             | $2x + \frac{5}{6}x = 170$            |
| 6. Multiplying by 6,                  | $12x + 5x = 1020$                    |
| 7. Uniting terms, and dividing by 17, | $x = 60$ , the lower part.           |

Then by the 3d step,       $2x + y = 168$

That is, substituting 60 for  $x$ ,  $120 + y = 168$       [per part.

Transposing 120,       $y = 168 - 120 = 48$ , the up-

Prob. 11. To find a fraction such that,

If a unit be added to the numerator, the fraction will be equal to  $\frac{1}{3}$ ; but

If a unit be added to the denominator, the fraction will be equal to  $\frac{1}{4}$ .

Let  $x$  = the numerator,      And  $y$  = the denominator.

- |                            |                                 |
|----------------------------|---------------------------------|
| 1. By the first condition, | $\frac{x+1}{y} = \frac{1}{3}$ } |
| 2. By the second,          | $\frac{x}{y+1} = \frac{1}{4}$ } |
| 3. Therefore               | $x = 4$ , the numerator,        |
| 4. And                     | $y = 15$ , the denominator.     |

Prob. 12. What two numbers are those,

Whose *difference* is to their sum, as 2 to 3; and

Whose *sum* is to their product, as 3 to 5?

Ans. 10 and 2.

Prob. 13. To find two numbers such, that

The product of their sum and difference shall be 5, and

The product of the sum of their squares and the difference of their squares shall be 65.

Let  $x$  = the greater number;      And  $y$  the less.

1. By the first condition,  $(x+y) \times (x-y) = 5$  }
2. By the second,  $(x^2+y^2) \times (x^2-y^2) = 65$  }
3. Mult. the factors in the 1st, (Art. 235,)  $x^2 - y^2 = 5$
4. Dividing the 2d by the 3d, (Art. 118.)  $x^2 + y^2 = 13$
5. Adding the 3d and 4th,  $2x^2 = 18$
6. Therefore  $x = 3$ , the greater number.
7. And  $y = 2$ , the less.

In the 4th step, the first member of the 2d equation is divided by  $x^2 - y^2$ , and the second member by 5, which is equal to  $x^2 - y^2$ .

Prob. 14. To find two numbers, whose difference is 8, and product 240.

Prob. 15. To find two numbers,  
Whose difference shall be 12, and  
The sum of their squares 1424.

Let  $x =$  the greater ;      And  $y =$  the less.

1. By the 1st condition,  $x - y = 12$  }
2. By the second,  $x^2 + y^2 = 1424$  }
3. Transp.  $y$  in the 1st,  $x = y + 12$
4. Squaring both sides,  $x^2 = y^2 + 24y + 144$
5. Transp.  $y^2$  in the 2d,  $x^2 = 1424 - y^2$
6. Making the 4th and 5th equal,  $y^2 + 24y + 144 = 1424 - y^2$
7. Therefore,  $y = -6 \pm \sqrt{676} = -6 \pm 26$
8. And  $x = y + 12 = 20 + 12 = 32$ .

### EQUATIONS WHICH CONTAIN THREE OR MORE UNKNOWN QUANTITIES.

330. In the examples hitherto given, each has contained no more than *two* unknown quantities. And two independent equations have been sufficient to express the conditions of the question. But problems may involve three or more unknown quantities ; and may require for their solution as many independent equations.

Suppose  $x + y + z = 12$  }  
And  $x + 2y - 2z = 10$  } are given, to find  $x, y,$  and  $z,$   
And  $x + y - z = 4$  }

From these three equations, two others may be derived, which shall contain only *two* unknown quantities. One of the three in the original equations may be exterminated, in the same manner as when there are, at first, only two, by the rules in arts. 324, 6, 8.

In the equations given above, if we transpose  $y$  and  $z$ , we shall have,

$$\left. \begin{array}{l} \text{In the first, } x=12-y-z \\ \text{In the second, } x=10-2y+2z \\ \text{In the third, } x=4-y+z \end{array} \right\}$$

From these we may deduce two new equations, from which  $x$  shall be excluded.

$$\left. \begin{array}{l} \text{By making the 1st and 2d equal, } 12-y-z=10-2y+2z \\ \text{By making the 2d and 3d equal, } 10-2y+2z=4-y+z \\ \text{Reducing the 1st of these two, } y=3z-2 \\ \text{Reducing the second, } y=z+6 \end{array} \right\}$$

From these two equations, one may be derived containing only *one* unknown quantity,

$$\left. \begin{array}{l} \text{Making one equal to the other, } 3z-2=z+6 \\ \text{And } z=4. \text{ Hence,} \end{array} \right\}$$

331. To solve a problem containing *three* unknown quantities, and producing three independent equations,

FIRST, FROM THE THREE EQUATIONS DEDUCE TWO, CONTAINING ONLY TWO UNKNOWN QUANTITIES,

THEN, FROM THESE TWO DEDUCE ONE, CONTAINING ONLY ONE UNKNOWN QUANTITY.

For making these reductions, the rules already given are sufficient. (Art. 324, 6, 8.)

Prob. 16. Let there be given,

$$\left. \begin{array}{l} 1. \text{ The equation } x+5y+6z=53 \\ 2. \text{ And } x+3y+3z=30 \\ 3. \text{ And } x+y+z=12 \end{array} \right\} \text{ To find } x, y, \text{ and } z.$$

From these three equations to derive two, containing only two unknown quantities,

$$\left. \begin{array}{l} 4. \text{ Subtract the 2d from the 1st, } 2y+3z=23 \\ 5. \text{ Subtract the 3d from the 2d, } 2y+2z=18 \end{array} \right\}$$

From these two, to derive one,

$$6. \text{ Subtract the 5th from the 4th, } z=5.$$

To find  $x$  and  $y$ , we have only to take their values from the 3d and 5th equations. (Art. 329.)

7. Reducing the 5th,  $y=9-z=9-5=4$

8. Transposing in the 3d,  $x=12-z-y=12-5-4=3$ .

Prob. 17. To find  $x$ ,  $y$ , and  $z$ , from

- |                                      |   |   |
|--------------------------------------|---|---|
| 1. The equation                      | $x+y+z=12$                              | } |
| 2. And                               | $x+2y+3z=20$                            |   |
| 3. And                               | $\frac{1}{3}x+\frac{1}{2}y+z=6$         |   |
| 4. Multiplying the 1st by 3,         | $3x+3y+3z=36$                           |   |
| 5. Subtracting the 2d from the 4th,  | $2x+y=16$                               |   |
| 6. Subtracting the 3d from the 1st,  | $x-\frac{1}{3}x+y-\frac{1}{2}y=6$       |   |
| 7. Clearing the 6th of fractions,    | $4x+3y=36$                              | } |
| 8. Multiplying the 5th by 3,         | $6x+3y=48$                              |   |
| 9. Subtracting the 7th from the 8th, | $2x=12$ . And $x=6$ .                   |   |
| 10. Reducing the 7th,                | $y=\frac{36-4x}{3}=\frac{36-24}{3}=4$ . |   |
| 11. Reducing the 1st,                | $z=12-x-y=12-6-4=2$ .                   |   |

In this example all the reductions have been made according to the *third* rule for exterminating unknown quantities. (Art. 328.) But either of the three may be used at pleasure.

332. A calculation may often be very much abridged, by the exercise of judgment, in stating the question, in selecting the equations from which others are to be deduced, in simplifying fractional expressions, in avoiding radical quantities, &c. The skill which is necessary for this purpose, however, is to be acquired, not from a system of rules; but from practice, and a habit of attention to the peculiarities in the conditions of different problems, the variety of ways in which the same quantity may be expressed, the numerous forms which equations may assume, &c. In many of the examples in this and the preceding sections, the processes might have been shortened. But the object has been to illustrate general principles, rather than to furnish specimens of expeditious solutions. The learner will do well, as he passes along, to exercise his skill in abridging the calculations which are here given, or substituting others in their stead.

Prob. 18. Given,  $\left\{ \begin{array}{l} 1. x+y=a \\ 2. x+z=b \\ 3. y+z=c \end{array} \right\}$  To find  $x, y,$  and  $z.$

$$\text{Ans. } x = \frac{a+b-c}{2}. \text{ And } y = \frac{a+c-b}{2}. \text{ And } z = \frac{b+c-a}{2}.$$

Prob. 19. Three persons,  $A, B,$  and  $C,$  purchase a horse for 100 dollars, but neither is able to pay for the whole. The payment would require,

The whole of  $A$ 's money, together with half of  $B$ 's; or

The whole of  $B$ 's, with one third of  $C$ 's; or

The whole of  $C$ 's, with one fourth of  $A$ 's.

How much money had each?

Let  $x = A$ 's

$z = C$ 's

$y = B$ 's

$a = 100$

By the first condition,

$$x + \frac{1}{2}y = a$$

By the second,

$$y + \frac{1}{3}z = a$$

By the third,

$$z + \frac{1}{4}x = a$$

Therefore  $x = 64.$   $y = 72.$

$z = 34.$

333. The learner must exercise his own judgment, as to the choice of the quantity to be first exterminated. It will generally be best to begin with that which is most free from co-efficients, fractions, radical signs, &c.

Prob. 20. The sum of the distances which three persons,  $A, B,$  and  $C,$  have travelled is 62 miles;

$A$ 's distance is equal to 4 times  $C$ 's, added to twice  $B$ 's; and

Twice  $A$ 's added to 3 times  $B$ 's, is equal to 17 times  $C$ 's.

What are the respective distances?

Ans.  $A$ 's, 46 miles;  $B$ 's, 9; and  $C$ 's, 7.

Prob. 21. To find  $x, y,$  and  $z,$  from

The equation

$$\frac{1}{2}x + \frac{1}{3}y + \frac{1}{4}z = 62$$

And

$$\frac{1}{3}x + \frac{1}{4}y + \frac{1}{5}z = 47$$

And

$$\frac{1}{4}x + \frac{1}{5}y + \frac{1}{6}z = 38$$

And

$$x = 24. \quad y = 60. \quad z = 120.$$

Prob. 22. Given  $\left\{ \begin{array}{l} xy = 600 \\ xz = 300 \\ yz = 200 \end{array} \right\}$  To find  $x, y,$  and  $z.$

Ans.  $x = 30.$

$y = 20.$

$z = 10.$

334. The same method which is employed for the reduction of three equations, may be extended to 4, 5, or any number of equations, containing as many unknown quantities. The unknown quantities may be exterminated, one after another, and the number of equations may be reduced by successive steps, from five to four, from four to three, from three to two, &c.

Prob. 23. To find  $w$ ,  $x$ ,  $y$ , and  $z$ , from

- |                            |                                       |   |                      |
|----------------------------|---------------------------------------|---|----------------------|
| 1. The equation            | $\frac{1}{2}y + z + \frac{1}{2}w = 8$ | } | Four equations.      |
| 2. And                     | $x + y + w = 9$                       |   |                      |
| 3. And                     | $x + y + z = 12$                      |   |                      |
| 4. And                     | $x + w + z = 10$                      |   |                      |
| 5. Clear. the 1st of frac. | $y + 2z + w = 16$                     | } | Three equations.     |
| 6. Subtract. 2d from 3d,   | $z - w = 3$                           |   |                      |
| 7. Subtract. 4th from 3d,  | $y - w = 2$                           |   |                      |
| 8. Adding 5th and 6th,     | $y + 3z = 19$                         | } | Two equations.       |
| 9. Subtract. 7th from 6th, | $-y + z = 1$                          |   |                      |
| 10. Adding 8th and 9th,    | $4z = 20$ . Or $z = 5$                | } | Quantities required. |
| 11. Transp. in the 8th,    | $y = 19 - 3z = 4$                     |   |                      |
| 12. Transp. in the 3d,     | $x = 12 - y - z = 3$                  |   |                      |
| 13. Transp. in the 2d,     | $w = 9 - x - y = 2$                   |   |                      |

Prob. 24. Given  $\left\{ \begin{array}{l} w + 50 = x \\ x + 120 = 3y \\ y + 120 = 2z \\ z + 195 = 3w \end{array} \right\}$  To find  $w$ ,  $x$ ,  $y$ , and  $z$ .

Answer.  $w = 100$   $y = 90$   
 $x = 150$   $z = 105$ .

Prob. 25. There is a certain number consisting of two digits. The left-hand digit is equal to 3 times the right-hand digit; and if twelve be subtracted from the number itself, the remainder will be equal to the square of the left-hand digit. What is the number?

Let  $x$  = the left-hand digit, and  $y$  = the right-hand digit.

As the *local* value of figures increases in a tenfold ratio from right to left; the number required  $= 10x + y$

By the conditions of the problem  $x = 3y$   
 And  $10x + y - 12 = x^2$

The required number is, therefore, 93.

Prob. 26. If a certain number be divided by the product of its two digits, the quotient will be 2; and if 27 be added to the number, the digits will be inverted. What is the number?

Ans. 36.

Prob. 27. There are two numbers, such, that if the less be taken from 3 times the greater, the remainder will be 35; and if 4 times the greater be divided by 3 times the less + 1, the quotient will be equal to the less. What are the numbers?

Ans. 13 and 4.

Prob. 28. There is a certain fraction, such, that if 3 be added to the numerator, the value of the fraction will be  $\frac{1}{3}$ ; but if 1 be subtracted from the denominator, the value will be  $\frac{1}{5}$ . What is the fraction?

Ans.  $\frac{4}{21}$ .

Prob. 29. A gentleman has two horses, and a saddle which is worth 10 guineas. If the saddle be put on the *first* horse, the value of both will be *double* that of the *second* horse; but if the saddle be put on the *second* horse, the value of both will be less than that of the *first* horse by 13 guineas. What is the value of each horse?

Ans. 56 and 33 guineas.

Prob. 30. Divide the number 90 into 4 such parts, that the first *increased* by 2, the second *diminished* by 2, the third *multiplied* by 2, and the fourth *divided* by 2, shall all be equal.

If  $x$ ,  $y$ , and  $z$ , be three of the parts, the fourth will be  $90 - x - y - z$ . And by the conditions,

$$x + 2 = y - 2$$

$$x + 2 = 2z$$

$$2z = \frac{90 - x - y - z}{2}$$

The parts required are 18, 22, 10, and 40.

Prob. 31. Find three numbers, such, that the *first* with  $\frac{1}{2}$  the sum of the *second* and *third* shall be 120; the *second* with  $\frac{1}{5}$  the difference of the *third* and *first* shall be 70; and  $\frac{1}{2}$  the sum of the three numbers shall be 95.

Prob. 32. What two numbers are those, whose difference, sum, and product, are as the numbers 2, 3, and 5?

Ans. 10 and 2.

**Prob. 33.** A Vintner sold, at one time, 20 dozen of port wine, and 30 dozen of sherry; and for the whole received 120 guineas. At another time, he sold 30 dozen of port, and 25 dozen of sherry, at the same prices as before; and for the whole received 140 guineas. What was the price of a dozen of each sort of wine?

**Ans.** The port was 3 guineas, and the sherry 2 guineas a dozen.

**Prob. 34.** A merchant having mixed a certain number of gallons of brandy and water, found that, if he had mixed 6 gallons more of each, he would have put into the mixture 7 gallons of brandy for every 6 of water. But if he had mixed 6 less of each, he would have put in 6 gallons of brandy for every 5 of water. How many of each did he mix?

**Ans.** 78 gallons of brandy, and 66 of water.

**Prob. 35.** What fraction is that, whose numerator being doubled, and the denominator increased by 7, the value becomes  $\frac{2}{3}$ ; but the denominator being doubled, and the numerator increased by 2, the value becomes  $\frac{3}{5}$ ? **Ans.**  $\frac{4}{5}$ .

\* **Prob. 36.** A person expends 30 cents in apples and pears, giving a cent for four apples, and a cent for 5 pears. He afterwards parts with half his apples and one third of his pears, the cost of which was 13 cents. How many did he buy of each? **Ans.** 72 apples, and 60 pears.

---

335. If in the algebraic statement of the conditions of a problem, the original equations are more numerous than the unknown quantities; these equations will either be *contradictory*, or one or more of them will be *superfluous*.

Thus the equations  $\begin{cases} 3x=60 \\ \frac{1}{2}x=20 \end{cases}$  are contradictory.

For by the first,  $x=20$ , while by the second,  $x=40$ .

But if the latter be altered, so as to give to  $x$  the same value as the former, it will be useless, in the statement of a

\* For more examples of the solution of Problems by equations, see Euler's Algebra, Part. I. Sec. 4, Simpson's Algebra, Sec. II, Simpson's Exercises, Maclaurin's Algebra, Part I, Chap. 2 and 13, Emerson's Algebra, Book II, Sec. 1, Saunderson's Algebra, Book II and III, Dodson's Mathematical Repository, and Bland's Algebraical Problems.



problem. For nothing can be determined from the one, which can not be from the other.

Thus, of the equations  $\left\{ \begin{array}{l} 3x=60 \\ \frac{1}{2}x=10 \end{array} \right\}$  one is superfluous.

For either of them is sufficient to determine the value of  $x$ . They are not *independent* equations. (Art. 322.) One is convertible into the other. For if we divide the 1st by 6, it will become the same as the second.

Or if we multiply the second by 6, it will become the same as the first.

336. But if the number of independent equations produced from the conditions of a problem, is *less* than the number of unknown quantities, the subject is not sufficiently limited to admit of a definite answer. For each equation can limit but one quantity. And to enable us to find this quantity, all the others connected with it, must either be previously known, or be determined from other equations. If this is not the case, there will be a variety of answers which will equally satisfy the conditions of the question. If, for instance, in the equation

$$x+y=100,$$

$x$  and  $y$  are required, there may be fifty different answers. The values of  $x$  and  $y$  may be either 99 and 1, or 98 and 2, or 97 and 3, &c. For the sum of each of these pairs of numbers is equal to 100. But if there is a second equation which determines *one* of these quantities, the other may then be found from the equation already given. As  $x+y=100$ , if  $x=46$ ,  $y$  must be such a number as added to 46 will make 100, that is, it must be 54. No other number will answer this condition.

337. For the sake of abridging the solution of a problem, however, the number of independent equations actually put upon paper is frequently less, than the number of unknown quantities. Suppose we are required to divide 100 into two such parts, that the greater shall be equal to three times the less. If we put  $x$  for the greater, the less will be  $100-x$ . (Art. 195.)

Then by the supposition,

$$x=300-3x.$$

Transposing and dividing,

$$x=75, \text{ the greater.}$$

And

$$100-75=25, \text{ the less.}$$

Here, two unknown quantities are found, although there appears to be but one independent equation. The reason of this is, that a part of the solution has been omitted, because it is so simple, as to be easily supplied by the mind. To have a view of the whole, without abridging, let  $x$  = the greater number, and  $y$  = the less.

- |                                 |                                |   |
|---------------------------------|--------------------------------|---|
| 1. Then by the supposition,     | $x + y = 100$                  | } |
| 2. And                          | $3y = x$                       |   |
| 3. Transposing $x$ in the 1st,  | $y = 100 - x$                  |   |
| 4. Dividing the 2d by 3,        | $y = \frac{1}{3}x$             |   |
| 5. Making the 3d and 4th equal, | $\frac{1}{3}x = 100 - x$       |   |
| 6. Multiplying by 3,            | $x = 300 - 3x$                 |   |
| 7. Transposing and dividing,    | $x = 75$ , the greater.        |   |
| 8. By the 3d step,              | $y = 100 - x = 25$ , the less. |   |

By comparing these two solutions with each other, it will be seen that the first begins at the 6th step of the latter, all the preceding parts being omitted, because they are too simple to require the formality of writing down.

Prob. To find two numbers whose sum is 30, and the difference of their squares 120.

$$\text{Let } a = 30 \qquad b = 120$$

$$x = \text{the less number required.}$$

Then  $a - x =$  the greater. (Art. 195.)

And  $a^2 - 2ax + x^2 =$  the square of the greater. (Art. 214.)

From this subtract  $x^2$  the square of the less, and we shall have  $a^2 - 2ax =$  the difference of their squares.

$$\text{Therefore } x = \frac{a^2 - b}{2a} = \frac{(30)^2 - 120}{2 \times 30} = 13.$$

338. In most cases also, the solution of a problem which contains many unknown quantities, may be abridged, by particular artifices in *substituting* a single letter for several. (Art. 321.)

\* Suppose four numbers,  $u$ ,  $x$ ,  $y$ , and  $z$ , are required, of which

The sum of the three first is	13
The sum of the two first and last	17
The sum of the first and two last	18
The sum of the three last	21

\* Ludlam's Algebra, art. 161.c.

- Then
1.  $u+x+y=13$
  2.  $u+x+z=17$
  3.  $u+y+z=18$
  4.  $x+y+z=21$ .

Let  $S$  be substituted for the *sum* of the four numbers, that is, for  $u+x+y+z$ . It will be seen that, of these four equations,

The first contains all the letters except  $z$ , that is,  $S-z=13$   
 The second contains all except  $y$ , that is,  $S-y=17$   
 The third contains all except  $x$ , that is,  $S-x=18$   
 The fourth contains all except  $u$ , that is,  $S-u=21$ .

Adding all these equations together, we have

$$4S-z-y-x-u=69$$

Or  $4S-(z+y+x+u)=69$  (Art. 88.c.)

But  $S=(z+y+x+u)$  by substitution.

Therefore,  $4S-S=69$ , that is,  $3S=69$ , and  $S=23$ .

Then putting 23 for  $S$ , in the four equations in which it is first introduced, we have

$$\left. \begin{array}{l} 23-z=13 \\ 23-y=17 \\ 23-x=18 \\ 23-u=21 \end{array} \right\} \text{Therefore } \left\{ \begin{array}{l} z=23-13=10 \\ y=23-17=6 \\ x=23-18=5 \\ u=23-21=2. \end{array} \right.$$

Contrivances of this sort for facilitating the solution of particular problems, must be left to be furnished for the occasion, by the ingenuity of the learner. They are of a nature not to be taught by a system of rules.

339. In the resolution of equations containing several unknown quantities, there will often be an advantage in adopting the following method of notation.

The co-efficients of one of the unknown quantities are represented,

In the *first* equation, by a single letter, as  $a$ .

In the *second*, by the same letter marked with an accent, as  $a'$ ,

In the *third*, by the same letter with a *double* accent, as  $a''$ ,

&c.

The co-efficients of the other unknown quantities, are represented by other letters marked in a similar manner; as are also the terms which consist of *known* quantities only.

Two equations containing the two unknown quantities  $x$  and  $y$  may be written thus,

$$\begin{aligned} ax + by &= c \\ a'x + b'y &= c' \end{aligned}$$

Three equations containing  $x$ ,  $y$ , and  $z$ , thus,

$$\begin{aligned} ax + by + cz &= d \\ a'x + b'y + c'z &= d' \\ a''x + b''y + c''z &= d'' \end{aligned}$$

Four equations containing  $x$ ,  $y$ ,  $z$ , and  $u$ , thus

$$\begin{aligned} ax + by + cz + du &= e \\ a'x + b'y + c'z + d'u &= e' \\ a''x + b''y + c''z + d''u &= e'' \\ a'''x + b'''y + c'''z + d'''u &= e''' \end{aligned}$$

The same *letter* is made the co-efficient of the same unknown quantity, in different equations, that the co-efficients of the several unknown quantities may be distinguished, in any part of the calculation. But the letter is marked with different *accents*, because it actually stands for different quantities.

Thus we may put  $a=4$ ,  $a'=6$ ,  $a''=10$ ,  $a'''=20$ , &c.

To find the value of  $x$  and  $y$ .

- |  |                                     |
|--|-------------------------------------|
| 1. In the equation,                          | $ax + by = c$ }                     |
| 2. And                                       | $a'x + b'y = c'$ }                  |
| 3. Multiplying the 1st by $b'$ , (Art. 328.) | $ab'x + bb'y = cb'$                 |
| 4. Multiplying the 2d by $b$ ,               | $ba'x + bb'y = bc'$                 |
| 5. Subtracting the 4th from the 3d,          | $ab'x - ba'x = cb' - bc'$           |
| 6. Dividing by $ab' - ba'$ (Art. 121.)       | $x = \frac{cb' - bc'}{ab' - ba'}$ } |
|  | $y = \frac{ac' - ca'}{ab' - ba'}$ } |

By a similar process,

The symmetry of these expressions is well calculated to fix them in the memory. The denominators are the same in both; and the numerators are like the denominators, except a change of one of the letters in each term. But the particular advantage of this method is, that the expressions here obtained may be considered as *general solutions*, which give the values of the unknown quantities, in other equations, of a similar nature.

$$\begin{array}{l} \text{Thus if } 10x + 6y = 100 \\ \text{And } 40x + 4y = 200 \end{array} \left. \vphantom{\begin{array}{l} 10x + 6y = 100 \\ 40x + 4y = 200 \end{array}} \right\}$$

$$\begin{array}{lll} \text{Then putting } a=10 & b=6 & c=100 \\ & a'=40 & b'=4 & c'=200 \end{array}$$

$$\text{We have } x = \frac{cb' - bc'}{ab' - ba'} = \frac{100 \times 4 - 6 \times 200}{10 \times 4 - 6 \times 40} = 4.$$

$$\text{And } y = \frac{ac' - ca'}{ab' - ba'} = \frac{10 \times 200 - 100 \times 40}{10 \times 4 - 6 \times 40} = 10.$$

The equations to be resolved may, originally, consist of more than three terms. But if they are of the first degree, and have only two unknown quantities, each may be reduced to three terms by substitution.

$$\text{Thus the equation } dx - 4x + hy - 6y = m + 8$$

$$\text{Is the same, by art. 120, as } (d-4)x + (h-6)y = m + 8.$$

$$\text{And putting } a=d-4, \quad b=h-6 \quad c=m+8$$

$$\text{It becomes } ax + by = c.*$$

#### DEMONSTRATION OF THEOREMS.

340. Equations have been applied, in this and the preceding sections, to the solution of *problems*. They may be employed with equal advantage, in the demonstration of *theorems*. The principal difference, in the two cases, is in the order in which the steps are arranged. The operations themselves are substantially the same. It is essential to a demonstration, that complete certainty be carried through every part of the process. (Art. 11.) This is effected, in the reduction of equations, by adhering to the general rule, to make no alteration which shall affect the value of one of the members, without equally increasing or diminishing the other. In applying this principle, we are guided by the axioms laid down in art. 63. These axioms are as applicable to the demonstration of theorems, as to the solution of problems.

But the *order* of the steps will generally be different. In solving a problem, the object is to find the value of the unknown quantity, by disengaging it from all other quantities. But in conducting a demonstration, it is necessary to bring

\* For the application of this plan of notation to the solution of equations which contain more than two unknown quantities, see La Croix's Algebra, art. 85, Maclaurin's Algebra, Part I. Chap. 12, Fenn's Algebra, p. 57, and a paper of Laplace, in the Memoirs of the Academy of Sciences for 1772.

the equation to that particular form which will express, in algebraic terms, the proposition to be proved.

**Ex. 1. Theorem.** Four times the product of any two numbers, is equal to the square of their sum, diminished by the square of their difference.

Let  $x$  = the greater number,  $s$  = their sum,  
 $y$  = the less,  $d$  = their difference.

*Demonstration.*

- |                                     |                                |
|-------------------------------------|--------------------------------|
| 1. By the notation                  | $x + y = s$ }                  |
| 2. And                              | $x - y = d$ }                  |
| 3. Adding the two, (Ax. 1.)         | $2x = s + d$                   |
| 4. Subtracting the 2d from the 1st, | $2y = s - d$                   |
| 5. Mult. 3d and 4th, (Ax. 3.)       | $4xy = (s + d) \times (s - d)$ |
| 6. That is, (Art. 235.)             | $4xy = s^2 - d^2$              |

The last equation expressed in words is the proposition which was to be demonstrated. It will be easily seen that it is equally applicable to any two numbers whatever. For the particular values of  $x$  and  $y$  will make no difference in the nature of the proof.

$$\text{Thus } 4 \times 8 \times 6 = (8 + 6)^2 - (8 - 6)^2 = 192.$$

$$\text{And } 4 \times 10 \times 6 = (10 + 6)^2 - (10 - 6)^2 = 240.$$

$$\text{And } 4 \times 12 \times 10 = (12 + 10)^2 - (12 - 10)^2 = 480.$$

**Theorem 2.** The sum of the squares of any two numbers is equal to the square of their difference, added to twice their product.

Let  $x$  = the greater,  $d$  = their difference.  
 $y$  = the less,  $p$  = their product.

*Demonstration.*

- |                             |                         |
|-----------------------------|-------------------------|
| 1. By the notation          | $x - y = d$ }           |
| 2. And                      | $xy = p$ }              |
| 3. Squaring the first,      | $x^2 - 2xy + y^2 = d^2$ |
| 4. Multiplying the 2d by 2, | $2xy = 2p$              |
| 5. Adding the 3d and 4th,   | $x^2 + y^2 = d^2 + 2p.$ |

$$\text{Thus } 10^2 + 8^2 = (10 - 8)^2 + 2 \times 10 \times 8 = 164.$$

341. General propositions are also *discovered*, in an expeditious manner, by means of equations. The relations of quantities may be presented to our view, in a great variety

of ways, by the several changes through which a given equation may be made to pass. Each step in the process will contain a distinct proposition.

Let  $s$  and  $d$  be the sum and difference of two quantities  $x$  and  $y$ , as before.

- |                              |  |
|------------------------------|--|
| 1. Then                      | $s = x + y$ }  |
| 2. And                       | $d = x - y$ }  |
| 3. Dividing the 1st by 2,    | $\frac{1}{2}s = \frac{1}{2}x - \frac{1}{2}y$                     |
| 4. Dividing the 2d by 2,     | $\frac{1}{2}d = \frac{1}{2}x - \frac{1}{2}y$                     |
| 5. Adding the 3d and 4th,    | $\frac{1}{2}s + \frac{1}{2}d = \frac{1}{2}x + \frac{1}{2}x = x$  |
| 6. Sub. the 4th from the 3d, | $\frac{1}{2}s - \frac{1}{2}d = \frac{1}{2}y + \frac{1}{2}y = y.$ |

That is,

*Half the difference of two quantities, added to half their sum, is equal to the greater ; and*

*Half their difference subtracted from half their sum, is equal to the less.*



## SECTION XII.



### RATIO AND PROPORTION.\*

ART. 342. **T**HE design of mathematical investigations, is to arrive at the knowledge of particular quantities, by comparing them with other quantities, either *equal* to, or *greater*, or *less* than those which are the objects of inquiry. The end is most commonly attained by means of a series of *equations* and *proportions*. When we make use of equa-

\* Euclid's Elements, Book 5, 7, 8. Euler's Algebra, Part I, Sec. 3. Emerson on Proportion. Camus' Geometry, Book III. Ludlam's Mathematics. Wallis' Algebra, Chap. 19, 20. Saunderson's Algebra, Book 7. Barrow's Mathematical Lectures. Analyst for March, 1814.

tions, we determine the quantity sought, by discovering its *equality* with some other quantity or quantities already known.

We have frequent occasion, however, to compare the unknown quantity with others which are *not equal* to it, but either greater or less. Here, a different mode of proceeding becomes necessary. We may inquire, either *how much* one of the quantities is greater than the other; or *how many times* the one contains the other. In finding the answer to either of these inquiries, we discover what is termed a *ratio* of the two quantities. One is called *arithmetical*, and the other *geometrical* ratio. It should be observed, however, that both these terms have been adopted arbitrarily, merely for distinction sake. Arithmetical ratio, and geometrical ratio, are both of them applicable to arithmetic, and both to geometry.

As the whole of the extensive and important subject of proportion depends upon ratios, it is necessary that these should be clearly and fully understood.

343. ARITHMETICAL RATIO is the DIFFERENCE between two quantities or sets of quantities. The quantities themselves are called the *terms* of the ratio, that is, the terms between which the ratio exists. Thus two is the arithmetical ratio of 5 to 3. This is sometimes expressed, by placing two points between the quantities thus, 5 . . 3, which is the same as 5-3. Indeed the term arithmetical ratio, and its notation, by points are almost needless. For the one is only a substitute for the word *difference*, and the other for the sign -.

344. If both the terms of an arithmetical ratio be *multiplied* or *divided* by the same quantity, the *ratio* will, in effect, be multiplied or divided by that quantity.

Thus if

$$a - b = r$$

Then mult. both sides by  $h$ , (Ax. 3.)  $ha - hb = hr$

And dividing by  $h$ , (Ax. 4.)

$$\frac{a}{h} - \frac{b}{h} = \frac{r}{h}$$

345. If the terms of one arithmetical ratio be added to, or subtracted from, the corresponding terms of another, the ratio of their sum or difference will be equal to the sum or difference of the two ratios.



If  $a-b$  }  
And  $d-h$  } are the two ratios,

Then  $(a+d)-(b+h)=(a-b)+(d-h)$ . For each  $=a+d-b-h$ .  
And  $(a-d)-(b-h)=(a-b)-(d-h)$ . For each  $=a-d-b+h$ .

Thus the arith. ratio of  $11 \dots 4$  is  $7$  }  
And the arith. ratio of  $5 \dots 2$  is  $3$  }

The ratio of the sum of the terms  $16 \dots 6$  is  $10$ , the sum of the ratios.

The ratio of the difference of the terms  $6 \dots 2$  is  $4$ , the difference of the ratios.

346. GEOMETRICAL RATIO IS THAT RELATION BETWEEN QUANTITIES WHICH IS EXPRESSED BY THE QUOTIENT OF THE ONE DIVIDED BY THE OTHER.\*

Thus the ratio of 8 to 4, is  $\frac{8}{4}$  or 2. For this is the quotient of 8 divided by 4. In other words, it shows how often 4 is contained in 8.

In the same manner, the ratio of any quantity to another may be expressed by dividing the former by the latter, or, which is the same thing, making the former the numerator of a fraction, and the latter the denominator.

Thus the ratio of  $a$  to  $b$  is  $\frac{a}{b}$ .

The ratio of  $d+h$  to  $b+c$ , is  $\frac{d+h}{b+c}$ .

347. Geometrical ratio is also expressed by placing two points, one over the other, between the quantities compared.

Thus  $a:b$  expresses the ratio of  $a$  to  $b$ ; and  $12:4$  the ratio of 12 to 4. The two quantities together are called a *couplet*, of which the first term is the *antecedent*, and the last, the *consequent*.

348. This notation by points, and the other in the form of a fraction, may be exchanged the one for the other, as convenience may require; observing to make the antecedent of the couplet, the numerator of the fraction, and the consequent the denominator.

Thus  $10:5$  is the same as  $\frac{10}{5}$  and  $b:d$ , the same as  $\frac{b}{d}$ .

349. Of these three, the antecedent, the consequent, and the ratio, any two being given, the other may be found.

\* See Note H.

Let  $a$  = the antecedent,  $c$  = the consequent,  $r$  = the ratio.

By definition  $r = \frac{a}{c}$ ; that is, the ratio is equal to the antecedent divided by the consequent.

Multiplying by  $c$ ,  $a = cr$ , that is, the antecedent is equal to the consequent multiplied into the ratio.

Dividing by  $r$ ,  $c = \frac{a}{r}$ , that is, the consequent is equal to the antecedent divided by the ratio.

Cor. 1. If two couplets have their antecedents equal, and their consequents equal, their ratios must be equal.\*

Cor. 2. If, in two couplets the ratios are equal, and the antecedents equal, the consequents are equal: and if the ratios are equal and the consequents equal, the antecedents are equal.†

350. If the two quantities compared are *equal*, the ratio is a unit, or a ratio of equality. Thus the ratio of  $3 \times 6 : 18$  is a unit, for the quotient of any quantity divided by itself is 1.

If the antecedent of a couplet is *greater* than the consequent, the ratio is greater than a unit. For if a dividend is greater than its divisor, the quotient is greater than a unit. Thus the ratio of  $18 : 6$  is 3. (Art. 128. cor.) This is called a ratio of *greater inequality*.

On the other hand, if the antecedent is *less* than the consequent, the ratio is less than a unit, and is called a ratio of *less inequality*. Thus the ratio of  $2 : 3$ , is less than a unit, because the dividend is less than the divisor.

351. INVERSE OR RECIPROCAL RATIO IS THE RATIO OF THE RECIPROALS OF TWO QUANTITIES. See art. 49.

Thus the reciprocal ratio of 6 to 3, is  $\frac{1}{6}$  to  $\frac{1}{3}$ , that is  $\frac{1}{6} \div \frac{1}{3}$ .

The direct ratio of  $a$  to  $b$  is  $\frac{a}{b}$ , that is, the antecedent divided by the consequent.

The reciprocal ratio, is  $\frac{1}{a} : \frac{1}{b}$  or  $\frac{1}{a} \div \frac{1}{b} = \frac{1}{a} \times \frac{b}{1} = \frac{b}{a}$ , that is, the consequent  $b$  divided by the antecedent  $a$ .

\* Euclid 7. 5.

† Euc. 9. 5.

Hence a reciprocal ratio is expressed by *inverting the fraction* which expresses the direct ratio; or, when the notation is by points, by *inverting the order of the terms*.

Thus  $a$  is to  $b$ , inversely, as  $b$  to  $a$ .

352. COMPOUND RATIO IS THE RATIO OF THE PRODUCTS OF THE CORRESPONDING TERMS OF TWO OR MORE SIMPLE RATIOS.\*

Thus the ratio of	6 : 3, is 2
And the ratio of	12 : 4, is 3

The ratio compounded of these is	72 : 12 = 6.
----------------------------------	--------------

Here the compound ratio is obtained by multiplying together the two antecedents, and also the two consequents, of the simple ratios.

So the ratio compounded,

Of the ratio of	$a : b$
And the ratio of	$c : d$
And the ratio of	$h : y$

Is the ratio of	$ach : bdy = \frac{ach}{bdy}$ .
-----------------	---------------------------------

Compound ratio is not different in its *nature* from any other ratio. The term is used, to denote the origin of the ratio, in particular cases.

Cor. The compound ratio is equal to the product of the simple ratios.

The ratio of	$a : b$ , is $\frac{a}{b}$
--------------	----------------------------

The ratio of	$c : d$ , is $\frac{c}{d}$
--------------	----------------------------

The ratio of	$h : y$ , is $\frac{h}{y}$
--------------	----------------------------

And the ratio compounded of these is  $\frac{ach}{bdy}$ , which is the product of the fractions expressing the simple ratios. (Art. 155.)

353. If, in a series of ratios, the consequent of each preceding couplet, is the antecedent of the following one, *the*

\* See Note I.

ratio of the first antecedent to the last consequent, is equal to that which is compounded of all the intervening ratios.\*

Thus, in the series of ratios

$$\begin{array}{l} a : b \\ b : c \\ c : d \\ d : h \end{array}$$

the ratio of  $a : h$  is equal to that which is compounded of the ratios of  $a : b$ , of  $b : c$ , of  $c : d$ , of  $d : h$ . For the compound ratio, by the last article, is  $\frac{abcd}{bcdh} = \frac{a}{h}$ , or  $a : h$ . (Art. 145.)

In the same manner, all the quantities which are both antecedents and consequents will *disappear* when the fractional product is reduced to its lowest terms, and will leave the compound ratio to be expressed by the first antecedent and the last consequent.

354. A particular class of compound ratios is produced, by multiplying a simple ratio into *itself*, or into another *equal* ratio. These are termed *duplicate*, *triplicate*, *quadruplicate*, &c. according to the number of multiplications.

A ratio compounded of *two* equal ratios, that is, the *square* of the simple ratio, is called a *duplicate* ratio.

One compounded of *three*, that is, the *cube* of the simple ratio, is called *triplicate*, &c.

In a similar manner, the ratio of the *square roots* of two quantities, is called a *subduplicate* ratio; that of the *cube roots*, a *subtriplicate* ratio, &c.

Thus the simple ratio of  $a$  to  $b$ , is  $a : b$

The duplicate ratio of  $a$  to  $b$ , is  $a^2 : b^2$

The triplicate ratio of  $a$  to  $b$  is  $a^3 : b^3$

The subduplicate ratio of  $a$  to  $b$ , is  $\sqrt{a} : \sqrt{b}$

The subtriplicate of  $a$  to  $b$ , is  $\sqrt[3]{a} : \sqrt[3]{b}$ , &c.

The terms *duplicate*, *triplicate*, &c. ought not to be confounded with *double*, *triple*, &c.†

The ratio of 6 to 2 is  $6 : 2 = 3$

Double this ratio, that is, twice the ratio is  $12 : 2 = 6$  }

Triple the ratio, i. e. three times the ratio, is  $18 : 2 = 9$  }

\* This is the particular case of compound ratio which is treated of in the 5th book of Euclid. See the editions of Simson and Playfair.

† See Note K.

But the *duplicate* ratio, i. e. the *square* of the ratio, is  $6^2:2^2=9$  }  
 And the *triplicate* ratio, i. e. the *cube* of the ratio, is  $6^3:2^3=27$  }

355. That quantities may have a ratio to each other, it is necessary that they should be so far of the same nature, as that one can properly be said to be either equal to, or greater, or less than the other. A foot has a ratio to an inch, for one is twelve times as great as the other. But it can not be said that an hour is either longer or shorter than a rod; or that an acre is greater or less than a degree. Still, if these quantities are expressed by *numbers*, there may be a ratio between the numbers. There is a ratio between the number of minutes in an hour, and the number of rods in a mile.

356. Having attended to the *nature* of ratios, we have next to consider in what manner they will be affected, by varying one or both of the terms between which the comparison is made. It must be kept in mind that, when a direct ratio is expressed by a fraction, the *antecedent* of the couplet is always the *numerator*, and the *consequent*, the *denominator*. It will be easy, then, to derive from the properties of fractions, the changes produced in ratios by variations in the quantities compared. For the ratio of the two quantities is the same as the *value* of the fractions, each being the *quotient* of the numerator divided by the denominator. (Arts. 135, 346.) Now it has been shown, (Art. 137.) that multiplying the numerator of a fraction by any quantity, is multiplying the *value* by that quantity; and that dividing the numerator is dividing the value. Hence,

357. *Multiplying the antecedent of a couplet by any quantity, is multiplying the ratio by that quantity; and dividing the antecedent is dividing the ratio.*

Thus the ratio of  $6:2$  is 3  
 And the ratio of  $24:2$  is 12.

Here the antecedent and the ratio, in the last couplet, are each four times as great as in the first.

The ratio of  $a:b$  is  $\frac{a}{b}$   
 And the ratio of  $na:b$  is  $\frac{na}{b}$ .

Cor. With a given consequent, the greater the *antecedent*, the greater the *ratio*; and on the other hand, the greater the *ratio*, the greater the *antecedent*.\* See art. 137. cor.

358. *Multiplying the consequent of a couplet by any quantity is, in effect, dividing the ratio by that quantity; and dividing the consequent is multiplying the ratio.* For multiplying the denominator of a fraction, is dividing the value; and dividing the denominator is multiplying the value. (Art. 138.)

Thus the ratio of 12:2, is 6

And the ratio of 12:4, is 3.

Here the consequent, in the second couplet, is *twice* as great, and the ratio only *half* as great, as in the first.

The ratio of  $a:b$  is  $\frac{a}{b}$

And the ratio of  $a:nb$ , is  $\frac{a}{nb}$ .

Cor. With a given antecedent, the greater the consequent, the less the ratio; and the greater the ratio, the less the consequent.† See art. 138. cor.

359. From the two last articles, it is evident that *multiplying the antecedent* of a couplet, by any quantity, will have the same effect on the ratio, as *dividing the consequent*, by that quantity; and *dividing the antecedent* will have the same effect as *multiplying the consequent*. See art. 139.

Thus the ratio of 3:4, is 2

Mult. the antecedent by 2, the ratio of 16:4, is 4

Divid. the consequent by 2, the ratio of 8:2, is 4.

Cor. Any *factor* or *divisor* may be transferred, from the antecedent of a couplet to the consequent, or from the consequent to the antecedent, without altering the ratio.

It must be observed that, when a factor is thus transferred from one term to the other, it becomes a divisor; and when a divisor is transferred, it becomes a factor.

Thus the ratio of  $3 \times 6 : 9 = 2$  } the same ratio.  
Transferring the factor 3,  $6 : \frac{9}{3} = 2$  }

\* Euclid 8 and 10. 5. The first part of the propositions.

† Euclid 8 and 10. 5. The last part of the propositions.

The ratio of	$\frac{ma}{y} : b = \frac{ma}{y} \div b = \frac{ma}{by}$	}
Transferring $y$ ,	$ma : by = ma \div by = \frac{ma}{by}$	
Transferring $m$ ,	$a : \frac{by}{m} = a \div \frac{by}{m} = \frac{ma}{by}$	

360. It is farther evident, from arts. 357 and 358, that if THE ANTECEDENT AND CONSEQUENT BE BOTH MULTIPLIED, OR BOTH DIVIDED, BY THE SAME QUANTITY, THE RATIO WILL NOT BE ALTERED.\* See art. 140.

Thus the ratio of	$8 : 4 = 2$	}
Mult. both terms by 2,	$16 : 8 = 2$	
Divid. both terms by 2,	$4 : 2 = 2$	

the same ratio.

The ratio of	$a : b = \frac{a}{b}$	}
Multiplying both terms by $m$ ,	$ma : mb = \frac{ma}{mb} = \frac{a}{b}$	
Dividing both terms by $n$ ,	$\frac{a \cdot b}{n \cdot n} = \frac{an}{bn} = \frac{a}{b}$	

Cor. 1. The ratio of two *fractions* which have a common denominator, is the same as the ratio of their *numerators*.

The ratio of  $\frac{a}{n} : \frac{b}{n}$ , is the same as that of  $a : b$ .

Cor. 2. The *direct* ratio of two fractions which have a common numerator, is the same as the reciprocal ratio of their *denominators*.

Thus the ratio of  $\frac{a}{m} : \frac{a}{n}$ , is the same as  $\frac{1}{m} : \frac{1}{n}$  or  $n : m$ .

361. From the last article, it will be easy to determine the ratio of any two fractions. If each term be multiplied by the two denominators, the ratio will be assigned in integral expressions. Thus, multiplying the terms of the couplet  $\frac{a}{b} : \frac{c}{d}$  by  $bd$ , we have  $\frac{abd}{b} : \frac{bcd}{d}$ , which becomes  $ad : bc$ , by cancelling equal quantities from the numerators and denominators.

\* Euclid 15. 5.

361.b. A ratio of *greater inequality*, compounded with another ratio, *increases* it.

Let the ratio of greater inequality be that of  $1+n:1$   
 And any given ratio, that of  $a:b$

The ratio compounded of these, (Art. 352,) is  $a+na:b$   
 Which is greater than that part of  $a:b$  (Art. 356, cor.)

But a ratio of *lesser inequality*, compounded with another ratio, *diminishes* it.

Let the ratio of lesser inequality be that of  $1-n:1$   
 And any given ratio, that of  $a:b$

The ratio compounded of these is  $a-na:b$   
 Which is less than that of  $a:b$ .

362. If to or from the terms of any couplet, there be ADDED or SUBTRACTED two other quantities having the same ratio, the sums or remainders will also have the same ratio.\*

Let the ratio of  $a:b$  }  
 Be the same as that of  $c:d$  }

Then the ratio of the *sum* of the antecedents, to the sum of the consequents, viz. of  $a+c$  to  $b+d$ , is also the same.

$$\text{That is } \frac{a+c}{b+d} = \frac{c}{d} = \frac{a}{b}.$$

*Demonstration.*

1. By supposition,

$$\frac{a}{b} = \frac{c}{d}$$

2. Multiplying by  $b$  and  $d$ ,

$$ad=bc$$

3. Adding  $cd$  to both sides

$$ad+cd=bc+cd$$

4. Dividing by  $d$ ,

$$a+c = \frac{bc+cd}{d}$$

5. Dividing by  $b+d$ ,

$$\frac{a+c}{b+d} = \frac{c}{d} = \frac{a}{b}.$$

The ratio of the *difference* of the antecedents, to the difference of the consequents, is also the same.

$$\text{That is } \frac{a-c}{b-d} = \frac{c}{d} = \frac{a}{b}.$$

\* Euclid 5 and 6. 5.



*Demonstration.*

- |                                      |   |
|--------------------------------------|---|
| 1. By supposition, as before,        | $\frac{a}{b} = \frac{c}{d}$                         |
| 2. Multiplying by $b$ and $d$ ,      | $ad = bc$   |
| 3. Subtracting $cd$ from both sides, | $ad - cd = bc - cd$                                 |
| 4. Dividing by $d$ ,                 | $a - c = \frac{bc - cd}{d}$                         |
| 5. Dividing by $b - d$               | $\frac{a - c}{b - d} = \frac{c}{d} = \frac{a}{b}$ . |

Thus the ratio of	15 : 5 is 3	}
And the ratio of	9 : 3 is 3	}

Then adding and subtracting the terms of the two couplets,

The ratio of	15 + 9 : 5 + 3 is 3	}
And the ratio of	15 - 9 : 5 - 3 is 3	}

Here the terms of only *two* couplets have been added together. But the proof may be extended to *any number* of couplets, where the ratios are equal. For, by the addition of the two first, a *new* couplet is formed, to which, upon the same principle, a third may be added, a fourth, &c. Hence,

363. If, in several couplets, the ratios are equal, THE SUM OF ALL THE ANTECEDENTS HAS THE SAME RATIO TO THE SUM OF ALL THE CONSEQUENTS, WHICH ANY ONE OF THE ANTECEDENTS HAS, TO ITS CONSEQUENT.\*

Thus the ratio  $\left\{ \begin{array}{l} 12 : 6 = 2 \\ 10 : 5 = 2 \\ 8 : 4 = 2 \\ 6 : 3 = 2 \end{array} \right.$

Therefore the ratio of  $(12 + 10 + 8 + 6) : (6 + 5 + 4 + 3) = 2$ .

363.b. A ratio of *greater inequality* is *diminished*, by adding the *same quantity* to both the terms.

Let the given ratio be that of  $a + b : a$  or  $\frac{a + b}{a}$

Adding  $x$  to both terms, it becomes  $a + b + x : a + x$  or  $\frac{a + b + x}{a + x}$

Reducing them to a common denominator,

The first becomes  $\frac{a^2 + ab + ax + bx}{a(a+x)}$

And the latter  $\frac{a^2 + ab + ax}{a(a+x)}$

As the latter numerator is manifestly less than the other, the *ratio* must be less. (Art. 356. cor.)

But a ratio of *lesser inequality* is increased, by adding the same quantity to both terms.

Let the given ratio be that of  $a-b : a$ , or  $\frac{a-b}{a}$

Adding  $x$  to both terms, it becomes  $a-b+x : a+x$  or  $\frac{a-b+x}{a+x}$

Reducing them to a common denominator,

The first becomes  $\frac{a^2 - ab + ax - bx}{a(a+x)}$

And the latter,  $\frac{a^2 - ab + ax}{a(a+x)}$

As the latter numerator is greater than the other, the *ratio* is greater.

If the same quantity, instead of being added, is *subtracted* from both terms, it is evident the effect upon the ratio must be reversed.

### Examples.

1. Which is the greatest, the ratio of 11 : 9, or that of 44 : 35?

2. Which is the greatest, the ratio of  $a+3 : \frac{1}{3}a$ , or that of  $2a+7 : \frac{1}{3}a$ ?

3. If the antecedent of a couplet be 65, and the ratio 13, what is the consequent?

4. If the consequent of a couplet be 7, and the ratio 18, what is the antecedent?

5. What is the ratio compounded of the ratios of 3 : 7, and  $2a : 5b$ , and  $7x+1 : 3y-2$ ?

6. What is the ratio compounded of  $x+y : b$ , and  $x-y : a+b$ , and  $a+b : h$ ?  
 Ans.  $x^2 - y^2 : bh$ .

7. If the ratios of  $5x+7:2x-3$ , and  $x+2:\frac{1}{2}x+3$  be compounded, will they produce a ratio of greater inequality, or of lesser inequality?

Ans. A ratio of greater inequality.

8. What is the ratio compounded of  $x+y:a$ , and  $x-y:b$ , and  $b:\frac{x^2-y^2}{a}$ ?

Ans. A ratio of equality.

9. What is the ratio compounded of  $7:5$ , and the duplicate ratio of  $4:9$ , and the triplicate ratio of  $3:2$ ?

Ans.  $14:15$ .

10. What is the ratio compounded of  $3:7$ , and the triplicate ratio of  $x:y$ , and the subduplicate ratio of  $49:9$ ?

Ans.  $x^3:y^3$ .

PROPORTION.

363. An accurate and familiar acquaintance with the doctrine of ratios, is necessary to a ready understanding of the principles of *proportion*, one of the most important of all the branches of the mathematics. In considering ratios, we compare two *quantities*, for the purpose of finding either their difference, or the quotient of the one divided by the other. But in proportion, the comparison is between two *ratios*. And this comparison is limited to such ratios as are *equal*. We do not inquire how much one ratio is *greater* or *less* than another, but whether they are the *same*. Thus the numbers 12, 6, 8, 4, are said to be proportional, because the ratio of  $12:6$  is the same as that of  $8:4$ .

364. PROPORTION, then, is an equality of ratios. It is either *arithmetical* or *geometrical*. Arithmetical proportion is an equality of arithmetical ratios, and geometrical proportion is an equality of geometrical ratios.\* Thus the numbers 6, 4, 10, 8, are in *arithmetical* proportion, because the *difference* between 6 and 4 is the same as the difference between 10 and 8. And the numbers 6, 2, 12, 4, are in *geometrical* proportion, because the *quotient* of 6 divided by 2 is the same, as the quotient of 12 divided by 4.

365. Care must be taken not to confound *proportion* with *ratio*. This caution is the more necessary, as in common discourse, the two terms are used indiscriminately, or rather,

\* See Note L.

proportion is used for both. The expenses of one man are said to bear a greater proportion to his income, than those of another. But according to the definition which has just been given, one proportion is neither greater nor less than another. For *equality* does not admit of degrees. One *ratio* may be greater or less than another. The ratio of 12 : 2 is greater than that of 6 : 2, and less than that of 20 : 2. But these differences are not applicable to *proportion*, when the term is used in its technical sense. The loose signification which is so frequently attached to this word, may be proper enough in *familiar language*: For it is sanctioned by general usage. But, for scientific purposes, the distinction between proportion and ratio, should be clearly drawn, and cautiously observed.

366. The equality between two ratios, as has been stated, is called proportion. The word is sometimes applied also to the series of terms among which this equality of ratios exists. Thus the two couplets 15 : 5 and 6 : 2 are, when taken together, called a proportion.

367. Proportion may be expressed, either by the common sign of equality, or by four points between the two couplets.

Thus  $\left\{ \begin{array}{l} 8 \cdot 6 = 4 \cdot 2, \text{ or } 8 \cdot 6 :: 4 \cdot 2 \\ a \cdot b = c \cdot d, \text{ or } a \cdot b :: c \cdot d \end{array} \right\}$  are arithmetical proportions.

And  $\left\{ \begin{array}{l} 12 : 6 = 8 : 4, \text{ or } 12 : 6 :: 8 : 4 \\ a : b = d : h, \text{ or } a : b :: d : h \end{array} \right\}$  are geometrical proportions.

The latter is read, 'the ratio of  $a$  to  $b$  equals the ratio of  $d$  to  $h$ ;' or more concisely, ' $a$  is to  $b$ , as  $d$  to  $h$ .'

368. The first and last terms are called the *extremes*, and the other two the *means*. *Homologous* terms are either the two antecedents or the two consequents. *Analogous* terms are the antecedent and consequent of the same couplet.

369. As the ratios are equal, it is manifestly immaterial which of the two couplets is placed first.

If  $a : b :: c : d$ , then  $c : d :: a : b$ . For if  $\frac{a}{b} = \frac{c}{d}$  then  $\frac{c}{d} = \frac{a}{b}$ .

370. The number of terms must be, at least, four. For the equality is between the ratios of *two couplets*; and each couplet must have an antecedent and a consequent. There may be a proportion, however, among three *quantities*. For

one of the quantities may be *repeated*, so as to form two terms. In this case, the quantity repeated is called the *middle term*, or a *mean proportional* between the two other quantities, especially if the proportion is geometrical.

Thus the numbers 8, 4, 2, are proportional. That is,  $8 : 4 :: 4 : 2$ . Here 4 is both the consequent in the first couplet, and the antecedent in the last. It is therefore a mean proportional between 8 and 2.

The *last* term is called a *third proportional* to the two other quantities. Thus 2 is a third proportional to 8 and 4.

371. *Inverse* or *reciprocal* proportion is an equality between a *direct* ratio and a *reciprocal* ratio.

Thus  $4 : 2 :: \frac{1}{3} : \frac{1}{6}$ ; that is, 4 is to 2, *reciprocally*, as 3 to 6. Sometimes also, the order of the terms in one of the couplets is inverted, without writing them in the form of a fraction. (Art. 351.)

Thus  $4 : 2 :: 3 : 6$  inversely. In this case, the *first* term is to the *second*, as the *fourth* to the *third*; that is, the first divided by the second, is equal to the fourth divided by the third.

372. When there is a series of quantities, such that the ratios of the first to the second, of the second to the third, of the third to the fourth, &c. are *all equal*; the quantities are said to be in *continued proportion*. The consequent of each preceding ratio is, then, the antecedent of the following one. Continued proportion is also called *progression*, as will be seen in a following section.

Thus the numbers 10, 8, 6, 4, 2, are in continued *arithmetical* proportion. For  $10 - 8 = 8 - 6 = 6 - 4 = 4 - 2$ .

The numbers 64, 32, 16, 8, 4, are in continued *geometrical* proportion. For  $64 : 32 :: 32 : 16 :: 16 : 8 :: 8 : 4$ .

If *a*, *b*, *c*, *d*, *h*, &c. are in continued geometrical proportion; then  $a : b :: b : c :: c : d :: d : h$ , &c.

One case of continued proportion is that of *three* proportional quantities. (Art. 370.)

373 As an *arithmetical* proportion is, generally, nothing more than a very simple equation, it is scarcely necessary to give the subject a separate consideration.

The proportion

$$a \cdot b :: c \cdot d$$

Is the same as the equation

$$a - b = c - d.$$

It will be proper, however, to observe that, if *four* quantities are in arithmetical proportion, *the sum of the extremes is equal to the sum of the means.*

Thus if  $a \cdot \cdot b :: h \cdot \cdot m$ , then  $a + m = b + h$

For by supposition,  $a - b = h - m$

And transposing  $-b$  and  $-m$   $a + m = b + h$

So in the proportion,  $12 \cdot \cdot 10 :: 11 \cdot \cdot 9$ , we have  $12 + 9 = 10 + 11$ .

Again, if *three* quantities are in arithmetical proportion, *the sum of the extremes is equal to double the mean.*

If  $a \cdot \cdot b :: b \cdot \cdot c$ , then  $a - b = b - c$

And transposing  $-b$  and  $-c$ ,  $a + c = 2b$ .

### GEOMETRICAL PROPORTION.

374. But if four quantities are in *geometrical* proportion, *the product of the extremes is equal to the product of the means.*

If  $a : b :: c : d$ ,  $ad = bc$

For by supposition, (Arts. 346, 364.)  $\frac{a}{b} = \frac{c}{d}$

Multiplying by  $bd$ , (Ax. 3.)  $\frac{abd}{b} = \frac{cbd}{d}$

Reducing the fractions,  $ad = bc$

Thus  $12 : 3 :: 15 : 10$ , therefore  $12 \times 10 = 3 \times 15$ .

Cor. Any *factor* may be transferred from one mean to the other, or from one extreme to the other, without affecting the proportion. If  $a : mb : x : y$ , then  $a : b :: mx : y$ . For the product of the means is, in both cases, the same. And if  $na : b :: x : y$ , then  $a : b :: x : ny$ .

375. On the other hand, if the product of two quantities is equal to the product of two others, the four quantities will form a proportion, when they are so arranged, that those on one side of the equation shall constitute the means, and those on the other side, the extremes.

If  $my = nh$ , then  $m : n :: h : y$ , that is  $\frac{m}{n} = \frac{h}{y}$

For by dividing  $my = nh$  by  $ny$ , we have  $\frac{my}{ny} = \frac{nh}{ny}$

And reducing the fractions,  $\frac{m}{n} = \frac{h}{y}$ .

Cor. The same must be true of *any factors* which form the two sides of an equation.

If  $(a+b) \times c = (d-m) \times y$ , then  $a+b : d-m :: y : c$ .

376. If *three* quantities are proportional, the product of the extremes is equal to the *square* of the mean. For this mean proportional is, at the same time, the consequent of the first couplet, and the antecedent of the last. (Art 370.) It is therefore to be multiplied *into itself*, that is, it is to be *squared*.

If  $a : b :: b : c$ , then mult. extremes and means,  $ac = b^2$ .

Hence, a *mean proportional* between two quantities may be found, by *extracting the square root of their product*.

If  $a : x :: x : c$ , then  $x^2 = ac$ , and  $x = \sqrt{ac}$ . (Art. 297.)

377. It follows, from art. 374, that in a proportion, either extreme is equal to the product of the means, divided by the other extreme; and either of the means is equal to the product of the extremes, divided by the other mean.

- |                                |   |
|--------------------------------|---|
| 1. If $a : b :: c : d$ , then  | $ad = bc$                                       |
| 2. Dividing by $d$ ,           | $a = \frac{bc}{d}$                              |
| 3. Dividing the first by $c$ , | $b = \frac{ad}{c}$                              |
| 4. Dividing it by $b$ ,        | $c = \frac{ad}{b}$                              |
| 5. Dividing it by $a$ ,        | $d = \frac{bc}{a}$ ; that is, the <i>fourth</i> |
- term is equal to the *product of the second and third divided by the first*.

On this principle is founded the rule of simple proportion in arithmetic, commonly called the *Rule of Three*. Three numbers are given to find a fourth, which is obtained by multiplying together the second and third, and dividing by the first.

378. The propositions respecting the products of the means, and of the extremes, furnish a very simple and convenient criterion for determining whether any four quantities are proportional. We have only to multiply the means together, and also the extremes. If the two products are equal, the quantities are proportional. If the two products are not equal, the quantities are not proportional,

379. In mathematical investigations, when the relations of several quantities are given, they are frequently stated in the form of a proportion. But it is commonly necessary that this first proportion should pass through a number of transformations, before it brings out distinctly the unknown quantity, or the proposition which we wish to demonstrate. It may undergo any change which will not affect the equality of the ratios; or which will leave the product of the means equal to the product of the extremes.

It is evident, in the first place, that any alteration in the *arrangement*, which will not affect the equality of these two products, will not destroy the proportion. Thus, if  $a:b::c:d$ , the order of these four quantities may be varied, in any way which will leave  $ad=bc$ . Hence,

380. If four quantities are proportional, THE ORDER OF THE MEANS, OR OF THE EXTREMES, OR OF THE TERMS OF BOTH COUPLETS, MAY BE INVERTED, WITHOUT DESTROYING THE PROPORTION.

If  $a:b::c:d$  }  
And  $12:8::6:4$  } then,

1. *Inverting the means,\**

$a:c::b:d$  } that is, { The *first*, is to the *third*,  
 $12:6::8:4$  } { As the *second*, to the *fourth*,

In other words, the ratio of the *antecedents* is equal to the ratio of the *consequents*.

This inversion of the means is frequently referred to by geometers under the name of *Alternation*.†

2. *Inverting the extremes.*

$d:b::c:a$  } that is, { The *fourth*, is to the *second*,  
 $4:3::6:12$  } { As the *third*, to the *first*.

3. *Inverting the terms of each couplet.*

$b:a::d:c$  } that is, { The *second*, is to the *first*,  
 $3:12::4:6$  } { As the *fourth*, to the *third*.

This is technically called *Inversion*.

Each of these may also be varied, by changing the order of the *two couplets*. (Art. 369.)

Cor. The order of the *whole proportion* may be inverted.

If  $a:b::c:d$ , then  $d:c::b:a$ .

\* See Note M. † Euclid 16. 5.



In each of these cases, it will be at once seen that, by taking the products of the means, and of the extremes, we have  $ad=bc$ , and  $12 \times 4 = 8 \times 6$ .

If the terms of only *one* of the couplets are inverted, the proportion becomes *reciprocal*. (Art. 371.)

If  $a : b :: c : d$ , then  $a$  is to  $b$ , reciprocally, as  $d$  to  $c$ .

381. A difference of arrangement is not the *only* alteration which we have occasion to produce, in the terms of a proportion. It is frequently necessary to multiply, divide, involve, &c. In all cases, the art of conducting the investigation consists in so ordering the several changes, as to maintain a constant equality, between the ratio of the two first terms, and that of the two last. As in resolving an equation, we must see that the *sides* remain equal; so in varying a proportion, the equality of the *ratios* must be preserved. And this is effected, either by keeping the ratios the *same*, while the *terms* are altered; or by increasing or diminishing *one* of the ratios, *as much as the other*. Most of the succeeding proofs are intended to bring this principle distinctly into view, and to make it familiar. Some of the propositions might be demonstrated, in a more simple manner, perhaps, by multiplying the extremes and means. But this would not give so clear a view of the *nature* of the several changes in the proportions.

It has been shown that, if *both* the terms of a couplet be multiplied or divided by the same quantity, the ratio will remain the same; (Art. 360.) that multiplying the *antecedent* is, in effect, multiplying the ratio, and dividing the antecedent, is dividing the ratio; (Art. 357.) and farther, that multiplying the *consequent* is, in effect, dividing the ratio, and dividing the consequent is multiplying the ratio. (Art. 358.) As the ratios in a proportion are equal, if they are both multiplied, or both divided, by the same quantity, they will still be equal. (Ax. 3.) One will be increased or diminished as much as the other. Hence,

382. If four quantities are proportional, TWO ANALOGOUS, OR TWO HOMOLOGOUS TERMS MAY BE MULTIPLIED OR DIVIDED BY THE SAME QUANTITY, WITHOUT DESTROYING THE PROPORTION.

If *analogous* terms be multiplied or divided, the ratios will not be altered. (Art. 360.) If *homologous* terms be multiplied or divided, both ratios will be equally increased or diminished. (Arts. 357, 8.)

If  $a : b :: c : d$ , then,

- |                                      |                                      |
|--------------------------------------|--------------------------------------|
| 1. Multiplying the two first terms,  | $ma : mb :: c : d$                   |
| 2. Multiplying the two last terms,   | $a : b :: mc : md$                   |
| 3. Multiplying the two antecedents,* | $ma : b :: mc : d$                   |
| 4. Multiplying the two consequents,  | $a : mb :: c : md$                   |
| 5. Dividing the two first terms,     | $\frac{a}{m} : \frac{b}{m} :: c : d$ |
| 6. Dividing the two last terms,      | $a : b :: \frac{c}{m} : \frac{d}{m}$ |
| 7. Dividing the two antecedents,     | $\frac{a}{m} : b :: \frac{c}{m} : d$ |
| 8. Dividing the two consequents,     | $a : \frac{b}{m} :: c : \frac{d}{m}$ |

Cor. 1. *All* the terms may be multiplied or divided by the same quantity.†

$$ma : mb :: mc : md \quad \frac{a}{m} : \frac{b}{m} :: \frac{c}{m} : \frac{d}{m}$$

Cor. 2. In any of the cases in this article, multiplication of the consequent may be substituted for division of the antecedent in the same couplet, and division of the consequent, for multiplication of the antecedent. (Art. 359, cor.)

Thus for  $\left\{ \begin{array}{l} ma : b :: mc : d \\ \frac{a}{m} : b :: \frac{c}{m} : d \end{array} \right\}$  may be put  $\left\{ \begin{array}{l} a : \frac{b}{m} :: mc : d \\ a : mb :: \frac{c}{m} : d \end{array} \right\}$  or  $\left\{ \begin{array}{l} ma : b :: c : \frac{d}{m} \\ \frac{a}{m} : b :: c : md. \end{array} \right.$

383. It is often necessary, not only to alter the terms of a proportion, and to vary the arrangement, but to *compare one proportion with another*. From this comparison will frequently arise a *new* proportion, which may be requisite in solving a problem, or in carrying forward a demonstration. One of the most important cases is that in which two of the terms in one of the proportions compared, are the *same* with two in the other. The similar terms may be made to *disappear*, and a new proportion may be formed of the four remaining terms. For,

\* Euclid 3. 5.

† Euclid 4. 5.

384. IF TWO RATIOS ARE RESPECTIVELY EQUAL TO A THIRD, THEY ARE EQUAL TO EACH OTHER.\*

This is nothing more than the 11th axiom applied to ratios.

1. If  $a : b :: m : n$   
 - And  $c : d :: m : n$  } then  $a : b :: c : d$ , or  $a : c :: b : d$ . (Art. 380.)

2. If  $a : b :: m : n$   
 And  $m : n :: c : d$  } then  $a : b :: c : d$ , or  $a : c :: b : d$ .

Cor. If  $a : b :: m : n$   
 $m : n > c : d$  } then  $a : b > c : d$ .†

For if the ratio of  $m : n$  is greater than that of  $c : d$ , it is manifest that the ratio of  $a : b$ , which is *equal* to that of  $m : n$ , is also greater than that of  $c : d$ .

385. In these instances, the terms which are alike, in the two proportions, are the two *first* and the two *last*. But this arrangement is not essential. The order of the terms may be changed, in various ways, without affecting the equality of the ratios.

1. The similar terms may be the two *antecedents*, or the two *consequents*, in each proportion. Thus,

If  $m : a :: n : b$   
 And  $m : c :: n : d$  } then { By alternation,  $m : n :: a : b$   
 { And  $m : n :: c : d$

Therefore  $a : b :: c : d$ , or  $a : c :: b : d$ , by the last article.

2. The *antecedents* in one of the proportions, may be the same as the *consequents* in the other.

If  $m : a :: n : b$   
 And  $c : m :: d : n$  } then { By inver. and altern.  $a : b :: m : n$   
 { By alternation,  $c : d :: m : n$

Therefore  $a : b$ , &c. as before.

3. Two *homologous* terms, in one of the proportions, may be the same, as two *analogous* terms in the other.

If  $a : m :: b : n$   
 And  $c : d :: m : n$  } then { By alternation,  $a : b :: m : n$   
 { And  $c : d :: m : n$

Therefore  $a : b$ , &c.

All these are instances of an *equality*, between the ratios in one proportion, and those in another. In geometry, the

\* Euclid 11. 5.

† Euclid 13. 5.

proposition to which they belong is usually cited by the words "*ex aequo*," or "*ex aequali*."\* The second case in this article is that which, in its form, most obviously answers to the explanation in Euclid. But they are all upon the same principle, and are frequently referred to, without discrimination.

336. Any number of proportions may be compared, in the same manner, if the two first or the two last terms in each preceding proportion, are the same with the two first or the two last in the following one.\*

$$\left. \begin{array}{l} \text{Thus if } a:b::c:d \\ \text{And } c:d::h:l \\ \text{And } h:l::m:n \\ \text{And } m:n::x:y \end{array} \right\} \text{ then } a:b::x:y.$$

That is, the two first terms of the first proportion have the same ratio, as the two last terms of the last proportion. For it is manifest that the ratio of *all* the couplets is the same.

And if the terms do not stand in the same order as here, yet if they can be *reduced* to this form, the same principle is applicable.

$$\left. \begin{array}{l} \text{Thus if } a:c::b:d \\ \text{And } c:h::d:l \\ \text{And } h:m::l:n \\ \text{And } m:x::n:y \end{array} \right\} \text{ then by alternation } \left\{ \begin{array}{l} a:b::c:d \\ c:d::h:l \\ h:l::m:n \\ m:n::x:y \end{array} \right.$$

Therefore  $a:b::x:y$ , as before.

In all the examples in this, and the preceding articles, the two terms in one proportion which have equals in another, are neither the two *means*, nor the two *extremes*, but one of the means, and one of the extremes; and the resulting proportion is uniformly *direct*.

337. But if the two means, or the two extremes, in one proportion, be the same with the means, or the extremes, in another, the four remaining terms will be *reciprocally proportional*.

$$\left. \begin{array}{l} \text{If } a:m::n:b \\ \text{And } c:m::n:d \end{array} \right\} \text{ then } a:c::\frac{1}{b}:\frac{1}{d}, \text{ or } a:c::d:b.$$

$$\left. \begin{array}{l} \text{For } ab=mn \\ \text{And } cd=mn \end{array} \right\} \text{ (Art. 374.) Therefore } ab=cd, \text{ and } a:c::d:b.$$

\* Euclid 22. 5.

In this example, the two means in one proportion, are like those in the other. But the principle will be the same, if the *extremes* are alike, or if the extremes in one proportion are like the means in the other.

If  $m : a :: b : n$  }  
 And  $m : c :: d : n$  } then  $a : c :: d : b$ .

Or if  $a : m :: n : b$  }  
 And  $m : c :: d : n$  } then  $a : c :: d : b$ .

The proposition in geometry which applies to this case, is usually cited by the words "*ex aequo perturbate.*"\*

388. Another way in which the terms of a proportion may be varied, is by *addition* or *subtraction*.

IF TO OR FROM TWO ANALOGOUS OR TWO HOMOLOGOUS TERMS OF A PROPORTION, TWO OTHER QUANTITIES HAVING THE SAME RATIO BE ADDED OR SUBTRACTED, THE PROPORTION WILL BE PRESERVED.†

For a ratio is not altered, by adding to it, or subtracting from it, the terms of another *equal* ratio. (Art. 362.)

If  $a : b :: c : d$  }  
 And  $a : b :: m : n$  }

Then by adding to, or subtracting from  $a$  and  $b$ , the terms of the equal ratio  $m : n$ , we have,

$$a + m : b + n :: c : d \quad \text{and} \quad a - m : b - n :: c : d.$$

And by adding and subtracting  $m$  and  $n$ , to and from  $c$  and  $d$  we have,

$$a : b :: c + m : d + n, \quad \text{and} \quad a : b :: c - m : d - n.$$

Here the addition and subtraction are to and from *analogous* terms. But by alternation, (Art. 380,) these terms will become *homologous*, and we shall have,

$$a + m : c :: b + n : d, \quad \text{and} \quad a - m : c :: b - n : d.$$

Cor. 1. This addition may, evidently, be extended to *any number* of equal ratios.‡

Thus if  $a : b ::$   $\left. \begin{array}{l} c : d \\ h : l \\ m : n \\ x : y \end{array} \right\}$

Then  $a : b :: c + h + m + x : d + l + n + y$ .

\* Euclid 23, 5.

† Euclid 2, 5.

‡ Euclid 2, 5. Cor.

Cor. 2. If  $a:b::c:d$  }  
 And  $m:b::n:d$  } then  $a+m:b::c+n:d$ .\*

For by alternation  $a:c::b:d$  } there- {  $a+m:c+n::b:d$   
 And  $m:n::b:d$  } fore { or  $a+m:b::c+n:d$ .

389. From the last article it is evident that if, in any proportion, the terms be added to, or subtracted from *each other*, that is,

IF TWO ANALOGOUS OR HOMOLOGOUS TERMS BE ADDED TO, OR SUBTRACTED FROM THE TWO OTHERS, THE PROPORTION WILL BE PRESERVED.

Thus, if  $a : b :: c : d$ , and  $12 : 4 :: 6 : 2$ , then,

1. *Adding* the two *last* terms, to the two *first*.

$$\begin{array}{ll} a+c:b+d::a:b & 12+6:4+2::12:4 \\ \text{and } a+c:b+d::c:d & 12+6:4+2::6:2 \\ \text{or } a+c:a::b+d:b & 12+6:12::4+2:4 \\ \text{and } a+c:c::b+d:d. & 12+6:6::4+2:2. \end{array}$$

2. *Adding* the two *antecedents*, to the two *consequents*.

$$\begin{array}{ll} a+b:b::c+d:d & 12+4:4::6+2:2 \\ a+b:a::c+d:c, \&c. & 12+4:12::6+2:6, \&c. \end{array}$$

This is called *Composition*.†

3. *Subtracting* the two *first* terms, from the two *last*.

$$\begin{array}{l} c-a:a::d-b:b \\ c-a:c::d-b:d, \&c. \end{array}$$

4. *Subtracting* the two *last* terms, from the two *first*.

$$\begin{array}{l} a-c:b-d::a:b\dagger \\ a-c:b-d::c:d, \&c. \end{array}$$

5. *Subtracting* the *consequents*, from the *antecedents*.

$$\begin{array}{l} a-b:b::c-d:d \\ a:a-b::c:c-d, \&c. \end{array}$$

The alteration expressed by the last of these forms is called *Conversion*.

6. *Subtracting* the *antecedents*, from the *consequents*.

$$\begin{array}{l} b-a:a::d-c:c \\ b:b-a::d:d-c, \&c. \end{array}$$

\* Euclid 24, 5.

† Euclid 18, 5.

‡ Euclid 19, 5.

7. Adding and subtracting,

$$a+b : a-b :: c+d : c-d.$$

That is, the sum of the two first terms, is to their difference, as the sum of the two last, to their difference.

Cor. If any compound quantities, arranged as in the preceding examples, are proportional, the simple quantities of which they are compounded are proportional also.

Thus, if  $a+b : b :: c+d : d$ , then  $a : b :: c : d$ .

This is called *Division*.\*

390. IF THE CORRESPONDING TERMS OF TWO OR MORE RANKS OF PROPORTIONAL QUANTITIES BE MULTIPLIED TOGETHER, THE PRODUCTS WILL BE PROPORTIONAL.

This is *compounding* ratios, (Art. 352,) or *compounding proportions*. It should be distinguished from what is called *composition*, which is an *addition* of the terms of a ratio. (Art. 389. 2.)

$$\begin{array}{l} \text{If } a : b :: c : d \\ \text{And } h : l :: m : n \end{array} \left. \vphantom{\begin{array}{l} a : b :: c : d \\ h : l :: m : n \end{array}} \right\} \begin{array}{l} 12 : 4 :: 6 : 2 \\ 10 : 5 :: 8 : 4 \end{array}$$

$$\text{Then } \underline{ah : bl :: cm : dn.} \qquad \underline{120 : 20 :: 48 : 8.}$$

For, from the nature of proportion, the two ratios in the first rank are equal, and also the ratios in the second rank. And multiplying the corresponding terms is multiplying the *ratios*, (Art. 352. cor.) that is, multiplying *equals by equals*; (Ax. 3.) so that the ratios will still be equal, and therefore the four products must be proportional.

The same proof is applicable to any number of proportions.

$$\text{If } \left\{ \begin{array}{l} a : b :: c : d \\ h : l :: m : n \\ p : q :: x : y \end{array} \right.$$

$$\text{Then } ahp : blq :: cmx : dny.$$

From this it is evident, that if the terms of a proportion be multiplied, each into *itself*, that is, if they be *raised to any power*, they will still be proportional.

$$\begin{array}{l} \text{If } a : b :: c : d \\ \quad a : b :: c : d \end{array} \qquad \begin{array}{l} 2 : 4 :: 6 : 12 \\ 2 : 4 :: 6 : 12 \end{array}$$

$$\text{Then } \underline{a^2 : b^2 :: c^2 : d^2} \qquad \underline{4 : 16 :: 36 : 144}$$

\* Euclid 17. 5. See Note N.

Proportionals will also be obtained, by reversing this process, that is, by extracting the roots of the terms.

If  $a:b::c:d$ , then  $\sqrt{a}:\sqrt{b}::\sqrt{c}:d$ .

For, taking the product of extr. and means,  $ad=bc$

And extracting both sides,

$$\sqrt{ad}=\sqrt{bc}$$

That is, (Arts. 259, 375.)

$$\sqrt{a}:\sqrt{b}::\sqrt{c}:\sqrt{d}.$$

Hence,

391. If several quantities are proportional, THEIR LIKE POWERS OR LIKE ROOTS ARE PROPORTIONAL.\*

If  $a:b::c:d$

Then  $a^n:b^n::c^n:d^n$ . and  $\sqrt[m]{a}:\sqrt[m]{b}::\sqrt[m]{c}:\sqrt[m]{d}$ .

And  $\sqrt[m]{a^n}:\sqrt[m]{b^n}::\sqrt[m]{c^n}:\sqrt[m]{d^n}$ , that is,  $a^{\frac{n}{m}}:b^{\frac{n}{m}}::c^{\frac{n}{m}}:d^{\frac{n}{m}}$ .

392. If the terms in one rank of proportionals be divided by the corresponding terms in another rank, the quotients will be proportional.

This is sometimes called the resolution of ratios.

If  $a:b::c:d$  }  $12:6::18:9$  }  
 And  $h:l::m:n$  }  $6:2::9:3$  }

Then  $\frac{a}{h}:\frac{b}{l}::\frac{c}{m}:\frac{d}{n}$   $\frac{12}{6}:\frac{6}{2}::\frac{18}{9}:\frac{9}{3}$

This is merely reversing the process in art. 390, and may be demonstrated in a similar manner.

This should be distinguished from what geometers call division, which is a subtraction of the terms of a ratio. (Art. 339. cor.)

When proportions are compounded by multiplication, it will often be the case, that the same factor will be found in two analogous or two homologous terms.

Thus if  $a:b::c:d$  }  
 And  $m:a::n:c$  }

$$\underline{am:ab::cn:cd}$$

Here  $a$  is in the two first terms, and  $c$  in the two last. Dividing by these, (Art. 382,) the proportion becomes

$$m:b::n:d. \quad \text{Hence,}$$

\* It must not be inferred from this, that quantities have the same ratio, as their like powers or like roots. See art. 354.



393. In compounding proportions, *equal factors*, or *divisors* in two analogous or homologous terms may be *rejected*.

$$\text{If } \begin{cases} a : b :: c : d \\ b : h :: d : l \\ h : m :: l : n \end{cases} \qquad \begin{cases} 12 : 4 :: 9 : 3 \\ 4 : 8 :: 3 : 6 \\ 8 : 20 :: 6 : 15 \end{cases}$$

$$\text{Then } \underline{a : m :: c : n} \qquad \underline{12 : 20 :: 9 : 15}$$

This rule may be applied to the cases, to which the terms "*ex aequo*," and "*ex aequo perturbate*" refer. See arts. 385 and 387. One of the methods may serve to verify the other.

394. The changes which may be made in proportions, without disturbing the equality of the ratios, are so numerous, that they would become burdensome to the memory, if they were not reducible to a few general principles. They are mostly produced,

1. By inverting the *order* of the terms, Art. 380.
2. By *multiplying* or *dividing* by the *same quantity*, Art. 382.
3. By comparing proportions which have *like terms*, Art. 384, 5, 6, 7.
4. By *adding* or *subtracting* the terms of equal ratios, Art. 388, 9.
5. By *multiplying* or *dividing* one proportion by another, Art. 390, 2, 3.
6. By *involving* or *extracting the roots* of the terms, Art. 391.

395. When four quantities are proportional, if the *first* be greater than the *second*, the *third* will be greater than the *fourth*; if equal, equal; if less, less.

For, the ratios of the two couplets being the same, if one is a ratio of *equality*, the other is also, and therefore the antecedent in each is *equal* to its consequent; (Art. 350.) if one is a ratio of *greater inequality*, the other is also, and therefore the antecedent in each is *greater* than its consequent; and if one is a ratio of *lesser inequality*, the other is also, and therefore the antecedent in each is *less* than its consequent.

$$\text{Let } a : b :: c : d ; \text{ then if } \begin{cases} a = b, c = d \\ a > b, c > d \\ a < b, c < d. \end{cases}$$

Cor. 1. If the *first* be greater than the *third*, the *second* will be greater than the *fourth*; if equal, equal; if less, less.\*

For by alternation,  $a : b :: c : d$  becomes  $a : c :: b : d$ , without any alteration of the quantities. Therefore, if  $a = b$ ,  $c = d$ , &c. as before.

Cor. 2. If  $a : m :: c : n$  }  
And  $m : b :: n : d$  } then if  $a = b$ ,  $c = d$ , &c.†

For, by equality of ratios, (Art. 385. 2.) or compounding ratios, (Arts. 390, 393.)

$a : b :: c : d$ . Therefore, if  $a = b$ ,  $c = d$ , &c. as before.

Cor. 3. If  $a : m :: n : d$  }  
And  $m : b :: c : n$  } then if  $a = b$ ,  $c = d$ , &c.‡

For, by compounding ratios, (Arts. 390, 393.)

$a : b :: c : d$ . Therefore, if  $a = b$ ,  $c = d$ , &c.

395.b. If four quantities are proportional, their *reciprocals* are proportional; and v. v.

If  $a : b :: c : d$ , then  $\frac{1}{a} : \frac{1}{b} :: \frac{1}{c} : \frac{1}{d}$ .

For in each of these proportions, we have, by reduction,  $ad = bc$ .

### CONTINUED PROPORTION.

396. When quantities are in continued proportion, *all* the ratios are *equal*. (Art. 372.) If

$$a : b :: b : c :: c : d :: d : e,$$

the ratio of  $a : b$  is the same, as that of  $b : c$ , of  $c : d$ , or of  $d : e$ . The ratio of the *first* of these quantities to the *last*, is equal to the *product* of all the intervening ratios; (Art. 353,) that is, the ratio of  $a : e$  is equal to

$$\frac{a}{b} \times \frac{b}{c} \times \frac{c}{d} \times \frac{d}{e}.$$

But as the intervening ratios are all *equal*, instead of multiplying them into each other, we may multiply any one of them into *itself*; observing to make the number of factors

\* Euclid 14. 5.

† Euclid 20. 5.

‡ Euclid 21. 5.

equal to the number of intervening ratios. Thus the ratio of  $a : e$ , in the example just given, is equal to

$$\frac{a}{b} \times \frac{a}{b} \times \frac{a}{b} \times \frac{a}{b} = \frac{a^4}{b^4}.$$

When several quantities are in continued proportion, the number of couplets, and of course, the number of ratios, is *one less* than the number of quantities. Thus the five proportional quantities  $a, b, c, d, e$ , form four couplets containing four ratios; and the ratio of  $a : e$  is equal to the ratio of  $a^4 : b^4$ , that is, the ratio of the fourth power of the first quantity, to the fourth power of the second. Hence,

397. If *three* quantities are proportional, *the first is to the third, as the square of the first, to the square of the second*; or as the square of the second, to the square of the third. In other words, the first has to the third, a *duplicate* ratio of the first to the second. And conversely, if the first of three quantities is to the third, as the square of the first to the square of the second; the three quantities are proportional.

If  $a : b :: b : c$ , then  $a : c :: a^2 : b^2$ . Universally,

398. If several quantities are in continued proportion, the ratio of the first to the last is equal to one of the intervening ratios raised to a power whose index is one less than the number of quantities.

If there are *four* proportionals  $a, b, c, d$ , then  $a : d :: a^3 : b^3$ .

If there are *five*  $a, b, c, d, e$ ,  $a : e :: a^4 : b^4$ , &c.

399. If several quantities are in continued proportion, they will be proportional when the order of the whole is *inverted*. This has already been proved, with respect to *four* proportional quantities. (Art. 380. cor.) It may be extended to any number of quantities.

Between the numbers,	64, 32, 16, 8, 4,
The ratios are	2, 2, 2, 2,
Between the same inverted	4, 8, 16, 32, 64,
The ratios are	$\frac{1}{2}$ , $\frac{1}{2}$ , $\frac{1}{2}$ , $\frac{1}{2}$ .

So if the order of any proportional quantities be inverted, the ratios in one series will be the *reciprocals* of those in the other. For, by the inversion, each antecedent becomes a consequent, and *v. v.* and the ratio of a consequent to its antecedent is the reciprocal of the ratio of the antecedent

to the consequent. (Art. 351.) That the reciprocals of equal quantities are themselves equal, is evident from ax. 4.

*Examples.*

1. Divide the number 49 into two such parts, that the greater increased by 6, may be to the less diminished by 11; as 9 to 2.

Let  $x =$  the greater, and  $49 - x =$  the less.

By the conditions proposed,  $x + 6 : 38 - x :: 9 : 2$

Adding terms, (Art. 389, 2.)  $x + 6 : 44 :: 9 : 11$

Dividing the consequents, (Art. 382, 8.)  $x + 6 : 4 :: 9 : 1$

Multiplying extremes and means,  $x + 6 = 36$ . And  $x = 30$ .

2. What number is that, to which if 1, 5, and 13, be severally added, the first sum shall be to the second, as the second to the third?

Let  $x =$  the number required.

By the conditions,  $x + 1 : x + 5 :: x + 5 : x + 13$

Subtracting terms, (Art. 389, 6.)  $x + 1 : 4 :: x + 5 : 8$

Therefore  $8x + 8 = 4x + 20$ . And  $x = 3$ .

3. Find two numbers, the greater of which shall be to the less, as their sum to 42; and as their difference to 6.

Let  $x$  and  $y =$  the numbers.

By the conditions  $x : y :: x + y : 42$

And  $x : y :: x - y : 6$

By equality of ratios  $x + y : 42 :: x - y : 6$

Inverting the means  $x + y : x - y :: 42 : 6$

Adding and subtracting terms, (Art. 389, 7,)  $2x : 2y :: 48 : 36$

Dividing terms, (Art. 382.)  $x : y :: 4 : 3$

Therefore  $3x = 4y$  And  $x = \frac{4y}{3}$ .

From the 2d proportion,  $6x = y \times (x - y)$

Substituting  $\frac{4y}{3}$  for  $x$ ,  $y = 24$ . And  $x = 32$ .

4. Divide the number 18 into two such parts, that the squares of those parts may be in the proportion of 25 to 16.

Let  $x =$  the greater part, and  $18 - x =$  the less.

By the conditions,  $x^2 : (18-x)^2 :: 25 : 16$   
 Extracting, (Art. 391.)  $x : 18-x :: 5 : 4$   
 Adding terms,  $x : 18 :: 5 : 9$   
 Dividing terms,  $x : 2 :: 5 : 1$   
 Therefore,  $x=10$ .

5. Divide the number 14 into two such parts, that the quotient of the greater divided by the less, shall be to the quotient of the less divided by the greater, as 16 to 9.

Let  $x =$  the greater part, and  $14-x =$  the less.

By the conditions,  $\frac{x}{14-x} : \frac{14-x}{x} :: 16 : 9$   
 Multiplying terms,  $x^2 : (14-x)^2 :: 16 : 9$   
 Extracting,  $x : 14-x :: 4 : 3$   
 Adding terms,  $x : 14 :: 4 : 7$   
 Dividing terms,  $x : 2 :: 4 : 1$   
 Therefore,  $x=8$ .

6. If the number 20 be divided into two parts, which are to each other in the *duplicate* ratio of 3 to 1, what number is a mean proportional between those parts?

Let  $x =$  the greater part, and  $20-x =$  the less.

By the conditions,  $x : 20-x :: 3^2 : 1^2 :: 9 : 1$   
 Adding terms  $x : 20 :: 9 : 10$   
 Therefore  $x=18$ . And  $20-x=2$   
 A mean propor. between 18 and 2 (Art. 376.)  $=\sqrt{2 \times 18}=6$ .

7. There are two numbers whose product is 24, and the difference of their cubes, is to the cube of their difference, as 19 to 1. What are the numbers?

Let  $x$  and  $y$  be equal to the two numbers.

1. By supposition  $xy=24$  }  
 2. And  $x^3 - y^3 : (x-y)^3 :: 19 : 1$  }  
 3. Or, (Art. 217.)  $x^3 - y^3 : x^3 - 3x^2y + 3xy^2 - y^3 :: 19 : 1$   
 4. Therefore, (Art. 389, 5)  $3x^2y - 3xy^2 : (x-y)^3 :: 18 : 1$   
 5. Dividing by  $x-y$  (Art. 382, 5)  $3xy : (x-y)^2 :: 18 : 1$   
 6. Or, as  $3xy=3 \times 24=72$ ,  $72 : (x-y)^2 :: 18 : 1$   
 7. Multiplying extremes and means,  $(x-y)^2=4$   
 8. Extracting,  $x-y=2$  }  
 9. By the first condition, we have  $xy=24$  }

Reducing these two equations, we have  $x=6$ , and  $y=4$ .

8. It is required to prove that  $a : x :: \sqrt{2a-y} : \sqrt{y}$   
 on supposition that  $(a+x)^2 : (a-x)^2 :: x+y : x-y$ .\*

1. Expanding,  $a^2 + 2ax + x^2 : a^2 - 2ax + x^2 :: x+y : x-y$
2. Adding and subtracting terms,  $2a^2 + 2x^2 : 4ax :: 2x : 2y$
3. Dividing terms,  $a^2 + x^2 : 2ax :: x : y$
4. Transf. the factor  $x$ , (Art. 374. cor.)  $a^2 + x^2 : 2a :: x^2 : y$
5. Inverting the means,  $a^2 + x^2 : x^2 :: 2a : y$
6. Subtracting terms,  $a^2 : x^2 :: 2a - y : y$
7. Extracting  $a : x :: \sqrt{2a-y} : \sqrt{y}$ .

9. It is required to prove that  $dx = cy$ , if  $x$  is to  $y$  in the triplicate ratio of  $a : b$ , and  $a : b :: \sqrt[3]{c+x} : \sqrt[3]{d+y}$ .

1. Involving terms.  $a^3 : b^3 :: c+x : d+y$
2. By the first supposition,  $a^3 : b^3 :: x : y$
3. By equality of ratios,  $c+x : d+y :: x : y$
4. Inverting the means,  $c+x : x :: d+y : y$
5. Subtracting terms,  $c : x :: d : y$
6. Therefore,  $dx = cy$ .

10. There are two numbers whose product is 135, and the difference of their squares, is to the square of their difference, as 4 to 1. What are the numbers?

Ans. 15 and 9.

11. What two numbers are those, whose difference, sum, and product, are as the numbers 2, 3, and 5, respectively?

Ans. 10 and 2.

12. Divide the number 24 into two such parts, that their product shall be to the sum of their squares, as 3 to 10.

Ans. 18 and 6.

13. In a mixture of rum and brandy, the difference between the quantities of each, is to the quantity of brandy, as 100 is to the number of gallons of rum; and the same difference is to the quantity of rum, as 4 to the number of gallons of brandy. How many gallons are there of each?

Ans. 25 of rum, and 5 of brandy.

14. There are two numbers which are to each other as 3 to 2. If 6 be added to the greater, and subtracted from

\* Bridge's Algebra.

the less, the sum and remainder will be to each other, as 3 to 1. What are the numbers?      Ans. 24 and 16.

15. There are two numbers whose product is 320; and the difference of their cubes, is to the cube of their difference, as 61 to 1. What are the numbers?

Ans. 20 and 16.

16. There are two numbers, which are to each other, in the duplicate ratio of 4 to 3; and 24 is a mean proportional between them. What are the numbers?

Ans. 32 and 18.



### SECTION XIII.



## DIVISION BY COMPOUND DIVISORS.

ART. 400. **I**N the section on division, the case in which the divisor is a compound quantity was omitted, because the operation, in most instances, requires some knowledge of the nature of *powers*; a subject which had not been previously explained.

Division by a compound divisor is performed by the following rule, which is substantially the same, as the rule for division in arithmetic :

To obtain the first term of the quotient, divide the first term of the dividend, by the first term of the divisor.

Multiply the whole divisor, by the term placed in the quotient; subtract the product from a part of the dividend; and to the remainder bring down as many of the following terms, as shall be necessary to continue the operation :

Divide again by the first term of the divisor, and proceed as before, till all the terms of the dividend are brought down.

Ex. 1. Divide  $ac + bc + ad + bd$ , by  $a + b$ .

$$a + b)ac + bc + ad + bd(c + d$$

$ac + bc$ , the first subtrahend.

$$\begin{array}{r} * \quad * \\ \hline ad + bd \end{array}$$

$ad + bd$ , the second subtrahend.

$$\begin{array}{r} * \quad * \\ \hline * \quad * \end{array}$$

Here  $ac$ , the first term of the dividend, is divided by  $a$ , the first term of the divisor, (Art. 116.) which gives  $c$  for the first term of the quotient. Multiplying the whole divisor by this; we have  $ac + bc$  to be subtracted from the two first terms of the dividend. The two remaining terms are then brought down, and the first of them is divided by the first term of the divisor, as before. This gives  $d$  for the second term of the quotient. Then multiplying the divisor by  $d$ , we have  $ad + bd$  to be subtracted, which exhausts the whole dividend, without leaving any remainder.

The rule is founded on this principle, that the product of the divisor into the several parts of the quotient, is equal to the dividend. (Art. 115.) Now by the operation, the product of the divisor into the *first* term of the quotient is subtracted from the dividend; then the product of the divisor in the *second* term of the quotient; and so on, till the product of the divisor into each term of the quotient, that is, the product of the divisor into the *whole* quotient, (Art. 100.) is taken from the dividend. If there is no remainder, it is evident that this product is *equal* to the dividend. If there *is* a remainder, the product of the divisor and quotient is equal to the whole of the dividend *except* the remainder. And this remainder is not included in the parts subtracted from the dividend, by operating according to the rule.

401. Before beginning to divide, it will generally be expedient to make some preparation in the *arrangement of the terms*.

The letter which is in the first term of the divisor, should be in the first term of the dividend also. And the *powers* of this letter should be arranged in order, both in the divisor and in the dividend; the highest power standing first, the next highest next, and so on.


Ex. 2. Divide  $2a^2b + b^3 + 2ab^2 + a^3$ , by  $a^2 + b^2 + ab$ .

Here if we take  $a^2$  for the first term of the divisor, the



other terms should be arranged according to the powers of  $a$ , thus,

$$\begin{array}{r} (a^2 + ab + b^2)a^3 + 2a^2b + 2ab^2 + b^3(a + b) \\ a^3 + a^2b + ab^2 \\ \hline a^2b + ab^2 + b^3 \\ a^2b + ab^2 + b^3 \\ \hline * \quad * \quad * \end{array}$$



In these operations, particular care will be necessary in the management of *negative quantities*. Constant attention must be paid to the rules for the signs in subtraction, multiplication and division. (Arts. 32, 105, 123.)

Ex. 3. Divide  $2ax - 2a^2x - 3a^3xy + 6a^3x + axy - xy$ , by  $2a - y$ .

If the terms be arranged according to the powers of  $a$ , they will stand thus ;

$$\begin{array}{r} 2a - y) 6a^3x - 3a^2xy - 2a^2x + axy + 2ax - xy(3a^2x - ax + x. \\ 6a^3x - 3a^2xy \end{array}$$

$$\begin{array}{r} * \quad * \quad -2a^2x + axy \\ -2a^2x + axy \\ \hline * \quad * \quad +2ax - xy \\ +2ax - xy \\ \hline \end{array}$$

402. In multiplication, some of the terms, by balancing each other, may be lost in the product. (Art. 110.) These may *re-appear* in division, so as to present terms in the course of the process, different from any which are in the dividend.

Ex. 4.

$$\begin{array}{r} (a + x)a^3 + x^3(a^2 - ax + x^2) \\ a^3 + a^2x \\ \hline * \quad -a^2x + x^3 \\ -a^2x - ax^2 \\ \hline * \quad ax^2 + x^3 \\ ax^2 + x^3 \\ \hline \end{array}$$

Ex. 5.

$$\begin{array}{r}
 a^2 - 2ax + 2x^2 \quad a^4 + 4x^4 (a^2 + 2ax + 2x^2) \\
 \underline{a^4 - 2a^3x + 2a^2x^2} \\
 * \quad + 2a^3x - 2a^2x^2 + 4x^4 \\
 \quad \quad + 2a^3x - 4a^2x^2 + 4ax^3 \\
 \hline
 * \quad \quad + 2a^2x^2 - 4ax^3 + 4x^4 \\
 \quad \quad + 2a^2x^2 - 4ax^3 + 4x^4 \\
 \hline
 \end{array}$$

If the learner will take the trouble to multiply the quotient into the divisor, in the two last examples, he will find, in the partial products, the several terms which appear in the process of dividing. But most of them, by balancing each other, are lost in the general product.

Ex. 6. Divide  $a^3 + a^2 + a^2b + ab + 3ac + 3c$ , by  $a + 1$ .  
 Quotient.  $a^2 + ab + 3c$ .

Ex. 7. Divide  $a + b - c - ax - bx + cx$ , by  $a + b - c$ .  
 Quotient.  $1 - x$ .

Ex. 8. Divide  $2a^4 - 13a^3x + 11a^2x^2 - 8ax^3 + 2x^4$ , by  $2a^2 - ax + x^2$ .  
 Quotient.  $a^2 - 6ax + 2x^2$ .

403. When there is a *remainder* after all the terms of the dividend have been brought down, this may be placed over the divisor and added to the quotient, as in arithmetic.

Ex. 9.

$$\begin{array}{r}
 (a+b)ac + bc + ad + bd + x(c+d + \frac{x}{a+b}) \\
 \underline{ac + bc} \\
 * \quad * \quad ad + bd \\
 \quad \quad ad + bd \\
 \hline
 * \quad * \quad x \\
 \hline
 \end{array}$$

Ex. 10.

$$\begin{array}{r}
 d-h)ad-ah+bd-bh+y(a+b+\frac{y}{d-h}) \\
 \underline{ad-ah} \\
 * \quad * \quad bd-bh \\
 \quad \quad \quad bd-bh \\
 \hline
 \quad \quad * \quad * \quad y \\
 \hline
 \end{array}$$

It is evident that  $a+b$  is the quotient belonging to the whole of the dividend, *excepting* the remainder  $y$ . (Art. 400. And  $\frac{y}{d-h}$  is the quotient belonging to this remainder. (Art. 124.)

Ex. 11. Divide  $6ax+2xy-3ab-by+3ac+cy+h$ , by  $3a+y$ .  
 Quotient.  $2x-b+c+\frac{h}{3a+y}$ .

Ex. 12. Divide  $a^2b-3a^2+2ab-6a-4b+22$ , by  $b-3$ .  
 Quotient.  $a^2+2a-4+\frac{10}{b-3}$ .

Ex. 13. See art. 283.

$$\begin{array}{r}
 a+\sqrt{b})ac+c\sqrt{b}+a\sqrt{d}+\sqrt{bd}(c+\sqrt{d}) \\
 \underline{ac+c\sqrt{b}} \\
 * \quad * \quad a\sqrt{d}+\sqrt{bd} \\
 \quad \quad \quad a\sqrt{d}+\sqrt{bd} \\
 \hline
 \end{array}$$

Ex. 14. Divide  $a+\sqrt{y}+ar\sqrt{y}+ry$ , by  $a+\sqrt{y}$ .  
 Quotient.  $1+r\sqrt{y}$ .

15. Divide  $x^3-3ax^2+3a^2x-a^3$ , by  $x-a$ .

16. Divide  $2y^3-19y^2+26y-17$ , by  $y-8$ .

17. Divide  $x^6-1$ , by  $x-1$ .

18. Divide  $4x^4-9x^2+6x-3$ , by  $2x^2+3x-1$ .

19. Divide  $a^4+4a^2b+3b^4$ , by  $a+2b$ .

20. Divide  $x^4-a^2x^2+2a^2x-a^4$ , by  $x^2-ax+a^2$ .

## SECTION XIV.

## EVOLUTION OF COMPOUND QUANTITIES.

Art. 404. **T**HE roots of compound quantities may be extracted by the following general rule :

After arranging the terms according to the powers of one of the letters, so that the highest power shall stand first, the next highest next, &c.

Take the root of the first term, for the first term of the required root :

Subtract the power from the given quantity, and divide the first term of the remainder, by the first term of the root involved to the next inferiour power, and multiplied by the index of the given power ;† the quotient will be the next term of the root.

Subtract the power of the terms already found from the given quantity, and, using the same divisor, proceed as before.

This rule verifies itself. For the root, whenever a new term is added to it, is involved, for the purpose of subtracting its power from the given quantity : and when the power is equal to this quantity, it is evident the true root is found.

Ex. 1. Extract the cube root of

$$a^6 + 3a^5 - 3a^4 - 11a^3 + 6a^2 + 12a - 8(a^2 + a - 2)$$

$a^6$ , the first subtrahend.

$$\underline{3a^4) * \quad 3a^5, \text{ \&c. the first remainder.}}$$

$$\underline{\quad \quad \quad a^6 + 3a^5 + 3a^4 + a^3, \text{ the 2d subtrahend.}}$$

$$\underline{3a^4) * \quad * \quad -6a^4, \text{ \&c. the 2d remainder.}}$$

$$\underline{\quad \quad \quad a^6 + 3a^5 - 3a^4 - 11a^3 + 6a^2 + 12a - 8.}$$

† By the given power is meant a power of the same name with the required root. As powers and roots are correlative, any quantity is the square of its square root, the cube of its cube root, &c.

Here  $a^3$ , the cube root of  $a^6$ , is taken for the first term of the required root. The power  $a^6$  is subtracted from the given quantity. For a divisor, the first term of the root is squared, that is, raised to the next inferior power, and multiplied by 3, the index of the given power.

By this, the first term of the remainder  $3a^5$ , &c. is divided, and the quotient  $a$  is added to the root. Then  $a^2 + a$ , the part of the root now found, is involved to the cube, for the second subtrahend, which is subtracted from the whole of the given quantity. The first term of the remainder  $-6a^4$ , &c. is divided by the divisor used above, and the quotient  $-2$  is added to the root. Lastly, the whole root is involved to the cube, and the power is found to be exactly equal to the given quantity.

It is not necessary to write the remainders at length, as, in dividing, the first term only is wanted.

2. Extract the fourth root of

$$\begin{array}{r}
 a^4 + 3a^3 + 24a^2 + 32a + 16(a + 2 \\
 a^4 \\
 \hline
 4a^3)^* \quad 3a^3, \text{ \&c.} \\
 \hline
 a^4 + 3a^3 + 24a^2 + 32a + 16. \\
 \hline
 \hline
 \end{array}$$

3. What is the 5th root of

$$a^5 + 5a^4b + 10a^3b^2 + 10a^2b^3 + 5ab^4 + b^5 ?$$

Ans.  $a + b$ .

4. What is the cube root of

$$a^3 - 6a^2b + 12ab^2 - 8b^3 ?$$

Ans.  $a - 2b$ .

5. What is the square root of

$$4a^2 - 12ab + 9b^2 + 16ah - 24bh + 16h^2 (2a - 3b + 4h)$$

$$4a^2$$


---


$$4a)^* \quad -12ab, \text{ \&c.}$$

---


$$4a^2 - 12ab + 9b^2$$


---

$$4a)^* \quad * \quad * \quad 16ah, \text{ \&c.}$$


---

$$4a^2 - 12ab + 9b^2 + 16ah - 24bh + 16h^2.$$

In finding the divisor here, the term  $2a$  in the root is not involved, because the power next below the square is the first power.

405. But the square root is more commonly extracted by the following rule, which is of the same nature, as that which is used in arithmetic.

After arranging the terms according to the powers of one of the letters, take the root of the first term, for the first term of the required root, and subtract the power from the given quantity.

Bring down two other terms for a dividend. Divide by double the root already found, and add the quotient, both to the root, and to the divisor. Multiply the divisor thus increased, into the term last placed in the root, and subtract the product from the dividend.

Bring down two or three additional terms, and proceed as before.

Ex. 1. What is the square root of

$$a^2 + 2ab + b^2 + 2ac + 2bc + c^2 (a+b+c)$$

$a^2$ , the first subtrahend.

$$\begin{array}{r} 2a+b) * \quad 2ab+b^2 \\ \text{Into } b = \quad 2ab+b^2, \text{ the 2d subtrahend.} \\ \hline \end{array}$$

$$\begin{array}{r} 2a+2b+c) * \quad * \quad 2ac+2bc+c^2 \\ \text{Into } c = \quad 2ac+2bc+c^2, \text{ the 3d subtrahend.} \\ \hline \hline \end{array}$$

Here it will be seen, that the several subtrahends are successively taken from the given quantity, till it is exhausted. If then, these subtrahends are together equal to the square of the terms placed in the root, the root is truly assigned by the rule.

The *first* subtrahend is the square of the first term of the root.

The *second* subtrahend is the product of the second term of the root, into itself, and into twice the preceding term.

The *third* subtrahend is the product of the third term of the root, into itself, and into twice the sum of the two preceding terms, &c.

That is, the subtrahends are equal to

$$a^2 + (2a+b) \times b + (2a+2b+c) \times c, \text{ \&c.}$$

and this expression is equal to the square of the root.

For  $(a+b)^2 = a^2 + 2ab + b^2 = a^2 + (2a+b) \times b$ . (Art. 120.)

And putting  $h = a+b$ , the square  $h^2 = a^2 + (2a+b) \times b$ .

And  $(a+b+c)^2 = (h+c)^2 = h^2 + (2h+c) \times c$ ;

that is, restoring the values of  $h$  and  $h^2$ ,

$$(a+b+c)^2 = a^2 + (2a+b) \times b + (2a+2b+c) \times c.$$

In the same manner it may be proved, that, if another term be added to the root, the power will be increased, by the product of that term, into itself, and into twice the sum of the preceding terms.

The demonstration will be substantially the same, if some of the terms be *negative*.

2. What is the square root of

$$\begin{array}{r} 1 - 4b + 4b^2 + 2y - 4by + y^2 \quad (1 - 2b + y \\ 1 \end{array}$$

$$\underline{\hspace{1.5cm}} \\ 2 - 2b) \quad * \quad -4b + 4b^2$$

Into  $-2b = -4b + 4b^2$

$$\begin{array}{r} 2 - 4b + y) \quad * \quad * \quad 2y - 4by + y^2 \\ \text{Into} \quad y = \quad \quad \quad \underline{\hspace{1.5cm}} \\ \quad \quad \quad \quad \quad \quad \quad 2y - 4by + y^2. \end{array}$$

3. What is the square root of

$$a^6 - 2a^5 + 3a^4 - 2a^3 + a^2 ? \quad \text{Ans.} \quad a^3 - a^2 + a.$$

4. What is the square root of

$$a^4 + 4a^2 b + 4b^2 - 4a^2 - 3b + 4 ? \quad \text{Ans.} \quad a^2 + 2b - 2.$$

406. It will frequently facilitate the extraction of roots, to consider the index as composed of two or more *factors*.

Thus  $a^{\frac{1}{4}} = a^{\frac{1}{2}} \times \frac{1}{2}$ . (Art. 253.) And  $a^{\frac{1}{6}} = a^{\frac{1}{3}} \times \frac{1}{2}$ . That is,

The fourth root is equal to the square root of the square root;

The sixth root is equal to the square root of the cube root;

The eighth root is equal to the square root of the fourth root, &c.

To find the sixth root, therefore, we may first extract the cube root, and then the square root of this.

1. Find the square root of  $x^4 - 4x^3 + 6x^2 - 4x + 1$ .
2. Find the cube root of  $x^6 - 5x^5 + 15x^4 - 20x^3 + 15x^2 - 6x + 1$ .
3. Find the square root of  $4x^6 - 4x^4 + 13x^2 - 6x + 9$ .
4. Find the fourth root of  

$$16a^4 - 96a^3x + 216a^2x^2 - 216ax^3 + 81x^4.$$
5. Find the 5th root of  $x^5 + 5x^4 + 10x^3 + 10x^2 + 5x + 1$ .
6. Find the sixth root of  

$$a^6 - 6a^5b + 15a^4b^2 - 20a^3b^3 + 15a^2b^4 - 6ab^5 + b^6.$$



## SECTION XV.

## INFINITE SERIES.

ART. 407. **I**T is frequently the case, that, in attempting to extract the root of a quantity, or to divide one quantity by another, we find it impossible to assign the quotient or root with exactness. But, by continuing the operation, one term after another may be added, so as to bring the result nearer and nearer to the value required. When the number of terms is supposed to be extended beyond any determinate limits, the expression is called an *infinite series*. The *quantity*, however, may be finite, though the number of terms be unlimited.

A fraction may often be expanded into an infinite series, by dividing the numerator by the denominator. For the value of a fraction is equal to the quotient of the numerator divided by the denominator. (Art. 135.) When this quotient can not be expressed, in a limited number of terms, it may be represented by an infinite series.

Ex. 1. To reduce the fraction  $\frac{1}{1-a}$  to an infinite series, divide 1 by  $1-a$ , according to the rule in art. 400.



$$\begin{array}{r}
 1-a) 1 \quad (1+a+a^2+a^3, \&c. \\
 \underline{1-a} \\
 * \quad a \\
 \quad \underline{a-a^2} \\
 * \quad a^2 \\
 \quad \underline{a^2-a^3} \\
 * \quad a^3, \&c. \\
 \underline{\hspace{1.5cm}}
 \end{array}$$

By continuing the operation, we obtain the terms

$1+a+a^2+a^3+a^4+a^5+a^6, \&c.$  which are sufficient to show that the series, after the first term, consists of the powers of  $a$ , rising regularly one above another.

2. If  $\frac{1}{1+a}$  be expanded, the series will be the same as that from  $\frac{1}{1-a}$ , except that the terms which consist of the *odd* powers of  $a$  will be *negative*.

So that  $\frac{1}{1+a} = 1 - a + a^2 - a^3 + a^4 - a^5 + a^6 \&c.$

3. Reduce  $\frac{h}{a-b}$  to an infinite series.

$$\begin{array}{r}
 a-b) h \quad \left( \frac{h}{a} + \frac{bh}{a^2} + \frac{b^2h}{a^3} \&c. \right. \\
 \underline{h - \frac{bh}{a}} \\
 * \quad \frac{bh}{a} \\
 \quad \underline{\frac{bh}{a} - \frac{b^2h}{a^2}} \\
 \quad \quad \underline{\frac{b^2h}{a^2} \&c.}
 \end{array}$$

Here  $h$  divided by  $a$ , gives  $\frac{h}{a}$  for the first term of the quotient. (Art. 124.) This is multiplied into  $a-b$ , and the

product is  $h - \frac{bh}{a}$ ; (Arts. 159, 153.) which subtracted from  $h$  leaves  $\frac{bh}{a}$ . This divided by  $a$ , gives  $\frac{bh}{a^2}$  (Art. 163.) for the second term of the quotient. If the operation be continued in the same manner, we shall obtain the series,

$$\frac{h}{a} + \frac{bh}{a^2} + \frac{b^2h}{a^3} + \frac{b^3h}{a^4} + \frac{b^4h}{a^5} \&c.$$

in which the exponents of  $b$  and of  $a$  increase regularly by 1.

4. Reduce  $\frac{1+a}{1-a}$  to an infinite series.

$$\text{Ans. } 1 + 2a + 2a^2 + 2a^3 + 2a^4 \&c.$$

403. An infinite series may be produced, by extracting the root of a compound surd.

Ex. 1. Reduce  $\sqrt{a^2 + b^2}$  to an infinite series, by extracting the square root, according to the rule in art. 405.

$$a^2 + b^2 \left( a + \frac{b^2}{2a} - \frac{b^4}{8a^3} + \frac{b^6}{16a^5} \&c. \right)$$

$$\frac{a^2}{2a + \frac{b^2}{2a}} \quad *$$

$$b^2 + \frac{b^4}{4a^2}$$

$$2a + \frac{b^2}{a} - \frac{b^4}{8a^3} \left) - \frac{b^4}{4a^2} \&c.$$

$$2. \sqrt{a^2 - b^2} = a - \frac{b^2}{2a} - \frac{b^4}{8a^3} - \frac{b^6}{16a^5} \&c.$$

$$3. \sqrt{2} = \sqrt{1+1} = 1 + \frac{1}{2} - \frac{1}{8} + \frac{1}{16} \&c.$$

## NOTES.

[Notes A and B are omitted in the abridgment.]

### NOTE C. Page 32.

**I**T is common to define multiplication, by saying that 'it is finding a product which has the same ratio to the multiplicand, that the multiplier has to a unit.' This is strictly and universally true. But the objection to it, *as a definition*, is, that the idea of ratio, as the term is understood in arithmetic and algebra, seems to imply a *previous* knowledge of multiplication, as well as of division. In this work at least, geometrical ratio is made to depend on division, and division, on multiplication. Ratio, therefore, could not be properly introduced into the definition of multiplication.

[Note D omitted in the abridgment.]

### NOTE E. p. 86.

As the *direct* powers of an integral quantity have *positive* indices, while the *reciprocal* powers have *negative* indices; it is common to call the former *positive powers*, and the latter *negative powers*. But this language is ambiguous, and may lead to mistake. For the same terms are applied to powers with positive and negative signs *prefixed*. Thus  $+8a^4$  is called a positive power; while  $8-a^4$  is called a negative one. It may occasion perplexity, to speak of the latter as being both positive and negative at the same time; positive, because it has a positive *index*, and negative, because it has a negative co-efficient. This ambiguity may be avoided, by using the terms direct and reciprocal; meaning, by the former, powers with positive exponents, and, by the latter, powers with negative exponents.

[Note F omitted.]

### NOTE G. p. 146.

Every affected quadratic equation may be reduced to one of the three following forms.

$$\begin{array}{l} 1. x^2 + ax = b \\ 2. x^2 - ax = b \\ 3. x^2 - ax = -b \end{array} \left. \vphantom{\begin{array}{l} 1. \\ 2. \\ 3. \end{array}} \right\}$$

These, when they are resolved, become

$$\left. \begin{array}{l} 1. x = -\frac{1}{2}a \pm \sqrt{\frac{1}{4}a^2 + b} \\ 2. x = \frac{1}{2}a \pm \sqrt{\frac{1}{4}a^2 + b} \\ 3. x = \frac{1}{2}a \pm \sqrt{\frac{1}{4}a^2 - b} \end{array} \right\}$$

In the two first of these forms, the roots are never imaginary. For the terms under the radical sign are both positive. But, in the third form, whenever  $b$  is greater than  $\frac{1}{4}a^2$ , the expression  $\frac{1}{4}a^2 - b$  is negative, and therefore its root is impossible.

[Note H omitted.]

NOTE I. p. 177.

This definition of compound ratio is more comprehensive than the one which is given in Euclid. That is included in this, but is limited to a particular case, which is stated in art. 353. It may answer the purposes of geometry, but is not sufficiently general for algebra.

NOTE K. p. 178.

It is not denied, that very respectable writers use these terms indiscriminately. But it appears to be without any necessity. The ratio of 6 to 2 is 3. There is certainly a difference between *twice* this ratio, and the *square* of it, that is, between twice three, and the square of three. All are agreed to call the latter a *duplicate* ratio. What occasion is there, then, to apply to it the term *double* also? This is wanted, to distinguish the other ratio. And if it is confined to that, it is used according to the common acceptance of the word, in familiar language.

NOTE L. p. 185.

The definition here given is meant to be applicable to quantities of every description. The subject of proportion, as it is treated of in Euclid, is embarrassed by the means which are taken to provide for the case of *incommensurable* quantities. But this difficulty is avoided by the algebraic notation, which may represent the ratio even of incommensurables.

NOTE M. p. 190.

The inversion of the means can be made, with strict propriety, in those cases only in which all the terms are quanti-

ties of the same kind. For, if the two last be different from the two first, the antecedent of each couplet, after the inversion, will be different from the consequent, and therefore, there can be no ratio between them. (Art. 355.)

This distinction, however, is of little importance in practice. For, when the several quantities are expressed in *numbers*, there will always be a ratio between the numbers. And when two of them are to be multiplied together, it is immaterial which is the multiplier, and which the multiplicand. Thus, in the Rule of Three in arithmetic, a change in the order of the two middle terms will make no difference in the result.

NOTE N. p. 197.

The terms *composition* and *division* are derived from geometry, and are introduced here, because they are generally used by writers on proportion. But they are calculated rather to perplex, than to assist, the learner. The objection to the word *composition* is, that its meaning is liable to be mistaken for the composition or compounding of *ratios*. (Art. 390.) The two cases are entirely different, and ought to be carefully distinguished. In one, the terms are *added*, in the other, they are *multiplied* together. The word compound has a similar ambiguity in other parts of the mathematics. The expression  $a+b$ , in which  $a$  is *added* to  $b$ , is called a compound quantity. The fraction  $\frac{1}{2}$  of  $\frac{2}{3}$ , or  $\frac{1}{2} \times \frac{2}{3}$ , in which  $\frac{1}{2}$  is *multiplied* into  $\frac{2}{3}$ , is called a compound fraction.

The term *division*, as it is used here, is also exceptionable. The alteration to which it is applied, is effected by *subtraction*, and has nothing of the nature of what is called division in arithmetic and algebra. But there is another case, (Art. 392.) totally distinct from this, in which the change in the terms of the proportion is actually produced by division.

THE END.

$a + b + c = 35 - 72 \times 3$  or  $512$   
 $2a + 2b + 2c = 372 - 512$   
 $2a + 2b + 2c = 140$   
 $a + b + c = 70$   
 $z = 70 - 1$


O mathematicians, Oh me  
 mathematicians "All's me"  
 I say the fact

"Empire Lead"

All the Mathematics  
of mathematicians "hem."

P. and S.







Miss

Handwritten text at the top of the page, possibly a title or header, which is mostly illegible due to fading and bleed-through.

Handwritten mark or symbol, possibly a stylized letter or a signature element.

Handwritten text, possibly a date or a name, located in the middle section of the page.

Large, decorative handwritten flourish or signature element on the left side of the bottom section.

Handwritten text, possibly a name or a signature, located to the right of the large flourish.

Handwritten text at the bottom of the page, possibly a date or a signature, which is partially obscured by the flourish above it.

# 14 DAY USE

RETURN TO DESK FROM WHICH BORROWED

LOAN DEPT.

This book is due on the last date stamped below, or  
on the date to which renewed.

Renewed books are subject to immediate recall.

**STACK DEAD**

JUL 21 1969 9 3

RECEIVED

JUL 9 '69 -6 PM

LOAN DEPT.

LD 21A-50m-9,'58  
(6889s10)476B

General Library  
University of California  
Berkeley

